

DIRECTIONS

USDA Reg. No. 5253-13

Lawns

Japanese Beetle, Oriental Beetle, Asiatic Beetle Larvae and White Grubs - Use 1 pound per 250 square feet.

Sod Webworms - Use 1 to 2 1/2 lbs. per 1000 square feet

Ants and Earwigs - Use 1 pound per 1000 square feet.

Note: Keep children and pets off treated areas until treated areas have been watered and have dried completely.

Cabbage

Cabbage Looper, Harlequin Cabbage Bug, Imported Cabbage Worm - Use 1 pound per 2000 square feet

Cabbage Maggot - Use 1 pound per 1000 square feet. apply to the soil or place around plants at the time of planting.

Strawberry

Strawberry Weevil and Thrips - Use 1 pound per 1000 square feet

Do not apply to strawberries after the blossoms open

Manufactured for
AGRICULTURAL PRODUCTS CO
WEST KINGSTON, R. I.

APCO

ACCEPTED

1-30-71

PER THE FEDERAL IN-
SPECTION SERVICE
OFFICE OF PEST CONTROL
REGISTERED PRODUCT
NUMBER 5253-13

6%

**CHLORDANE
DUST**

**CAUTION
KEEP OUT OF REACH
OF CHILDREN**

**SEE SIDE
PANEL FOR
ADDITIONAL
CAUTIONS**

**JAPANESE BEETLE GRUBS
SOD WEBWORMS • ANTS**

ACTIVE INGREDIENT:
Technical Chlordane 6.00
INERT INGREDIENTS 94.00
Equivalent to 0.6 Octachloro 1,7 Methano-
trahydroindane and 2.4 related compounds on
a weight basis.

NET WEIGHT ONE POUND

NET WEIGHT ONE POUND

APCO Chlordane 6% Dust may be applied any time of the day on either dry or moist foliage. Apply dust at frequent intervals beginning soon after pests appear. Both lower and upper sides of plant growth should be thoroughly dusted. For best results keep foliage covered.

Soil Insects - Following treatment on lawns, sprinkle the lawns thoroughly with water to wash the insecticide into the turf.

Surface Insects - Water lawn before application and not for 24 to 48 hours after application.

CAUTION

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS.

Avoid inhalation.

Avoid contact with skin, eyes and clothing. In case of skin contact, wash with soap and water. For eyes, get medical attention.

Avoid contamination of feed and feedstuffs.

Do not use or apply to cabbage and similar crops after edible parts have formed.

Do not use on crops to be taken to dairy animals or to animals going to market for slaughter.

Remove any family residues at harvest of vegetables and fruits, such as by washing.

Do not reuse empty container. Destroy it by burning with waste or burning. Stay away from smoke or fumes.

FISH and WILDLIFE CAUTION

Toxic to fish and wildlife. Birds feeding on lawn may be killed. Do not apply when weather conditions favor drift from areas treated. Do not contaminate any body of water, nor any other conditions favor runoff from areas treated.