

50 U.S. GALLONS

APPLY THIS PRODUCT ONLY
AS SPECIFIED ON LABEL

GENERAL INFORMATION

For information on the use of this product, consult the label. The label contains important information regarding the use of this product. The label also contains information regarding the safety of this product. The label is the most important part of the product. It is the responsibility of the user to read the label carefully and to follow the instructions on the label.

DIRECTIONS FOR USE

As a treatment for control of the carrier of Dutch elm disease.

FOR DORMANT SPRAY

IN HYDRAULIC SPRAYER:

Apply 2 to 4 gallons per acre of a 2% solution of this product to the bark of the tree. The solution should be applied to the bark of the tree in a series of vertical stripes. The stripes should be applied to the bark of the tree in a series of vertical stripes. The stripes should be applied to the bark of the tree in a series of vertical stripes.

IN MIST BLOWERS

Apply 2 to 4 gallons per acre of a 2% solution of this product to the bark of the tree. The solution should be applied to the bark of the tree in a series of vertical stripes. The stripes should be applied to the bark of the tree in a series of vertical stripes. The stripes should be applied to the bark of the tree in a series of vertical stripes.

FOR FOLIAR SPRAY

After leaves have bud or full grown.

IN HYDRAULIC SPRAYER:

Apply 2 to 4 gallons per acre of a 2% solution of this product to the leaves of the tree. The solution should be applied to the leaves of the tree in a series of vertical stripes. The stripes should be applied to the leaves of the tree in a series of vertical stripes. The stripes should be applied to the leaves of the tree in a series of vertical stripes.

IN MIST BLOWERS

Apply 2 to 4 gallons per acre of a 2% solution of this product to the leaves of the tree. The solution should be applied to the leaves of the tree in a series of vertical stripes. The stripes should be applied to the leaves of the tree in a series of vertical stripes. The stripes should be applied to the leaves of the tree in a series of vertical stripes.

WHEN TO APPLY

Apply this product to the bark of the tree in a series of vertical stripes. The stripes should be applied to the bark of the tree in a series of vertical stripes. The stripes should be applied to the bark of the tree in a series of vertical stripes.

MICHLIN CHEMICAL CORP
DETROIT MICH 48211

ACCEPTED
May 27 1971
FEDERAL AGENCY FOR THE PROTECTION OF THE ENVIRONMENT
U.S. DEPARTMENT OF AGRICULTURE
WASHINGTON, D.C. 20250

MICHLIN "MX-2"

25% TECHNICAL

METHOXYCHLOR

EMULSIFIABLE SOLUTION

CONTAINING

2 LBS. ACTUAL METHOXYCHLOR PER GALLON

ELM TREE SPRAY

INSECTICIDE

FOR CONTROL OF INSECT PESTS OF LIVESTOCK,
CROPS, STORED GRAIN, AND TREES

ACTIVE INGREDIENTS

Methoxychlor, 25%
Xylene

BY WEIGHT

25.0
2.0

INERT INGREDIENTS

73.0

100.0 = 100%

The above information is for informational purposes only. It is not intended to be used as a substitute for the label. The label is the most important part of the product. It is the responsibility of the user to read the label carefully and to follow the instructions on the label.

IMPORTANT: Users should observe the usual precautions concerning application of oil-containing products to crops during high temperatures since many crops are then more susceptible to oil injury.

WARNING!

KEEP OUT OF REACH OF CHILDREN

1. **HARMFUL IF SWALLOWED.**
2. Inhalation and skin contact may cause toxic symptoms. Avoid breathing spray mist. Avoid contact with the skin.
3. In case of spillage on skin, wash with soap and water.
4. Avoid contamination of feed and foodstuffs.
5. Do not spray house pets.
6. Do not feed treated plants to dairy animals or to animals to be slaughtered for slaughter.
7. Do not spray inside dairy barns to prevent contamination of milk.
8. This product is toxic to fish.
9. **FLAMMABLE MIXTURE:** Keep away from heat or open flame. Do not spray into or near fire or open flame.
10. Store at temperature above 40°F. If exposed to temperatures below 40°F, warm storage area to 40°F. Do not store in a cold storage area.

INSECTICIDE PRODUCTS DIVISION

MICHLIN MX-2 METHOXY-CHLOR INSECTICIDE FOR
USE AS A MOSQUITO LARVICIDE LIMITED TO NORTH
EAST AND MIDWEST CONTINENTAL UNITED STATES

THE RATES ARE ONLY FOR USE AS FOGS, SPRAYS IN WATER AND AERIAL APPLICATIONS
SHOULD NOT EXCEED SPECIFIED LIMITATIONS



HOW TO APPLY

For air-rattling fogging, use 2 to 4 lb. actual methoxy-chlor per acre.

Prehatch This is the best time to apply the insecticide to prevent the hatch of the larvae. Apply at the rate of 1 to 2 quarts per acre.

Larvicide This is the best time to apply the insecticide to control the larvae. Apply at the rate of 1 to 2 quarts per acre.

NOTE: Do not apply to water bodies where the water is used for drinking or for the raising of livestock. Do not apply to water bodies where the water is used for the raising of livestock.

OTHER RECOMMENDATIONS AND DIRECTIONS

FORAGE CROPS Do not apply to forage crops where the forage is used for the raising of livestock.

POTATOES

LIVESTOCK

Use as a spray only. Do not use in dipping vats.

HOUSING

SELF-OPERATOR FLY CONTROL ON CATTLE

GRAIN STORAGE BINS

CONTACT AND SPRAY PROTECTION

CONTACT

CONTACT

