

UNITED STATES FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
AND R.N.O. SUBJECT
TO ATTACHED COMMENTS

"BAY-GO" #101

RESIDUAL SURFACE SPRAY

READY-TO-USE

o-Isopropoxyphenyl methylcarbamate†	1.00%
2-Butoxyethanol	9.00%
† Petroleum Distillate	90.00%

This material used under license from Chemagro Corporation.

³ Baygon is the registered trademark of Farbenfabriken Bayer. Chemagro Corporation licensee.

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER PRECAUTIONS
ON THE BACK/SIDE PANEL



INSECTICIDE PRODUCTS DIVISION
MICHIGAN CHEMICAL CORPORATION
9045 VINCENT
DETROIT, MICHIGAN 48211

Michlin Formula "DKN-1" (20% Dieldrin) is a ready-to-use formulation for control of the pests listed below, including those resistant to certain other hydrocarbon and phosphate insecticides. This product drives insects from their hiding places, gives rapid kill and long residual activity.

KILLS: Ants, Brown Dog Ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Earwigs, Firebrats, Fleas, Flies, Gnats, Millipedes, Mosquitoes, Paperlice, Punkies, Sandflies, Scorpions, Silverfish, Sowbugs and Spiders. Exposed stages of Caddises, Cigarette Beetles, Dermestids, Drugstore Beetles, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Saw-toothed Grain Beetles and Weevils.

GENERAL: Apply with a low-pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run-off. Apply to all insect breeding sites and harborages. Also treat adjacent exposed surfaces where insects may crawl when they come out of hiding. Repeat treatment as necessary.

Cockroaches — Thoroughly treat the infested area. Spray baseboards, cracks and crevices; around window and door frames; the under surfaces of shelves, drawers and work tables; behind and beneath cabinets, refrigerators, sinks and stoves; in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or rest.

Ants – Apply to trails and places where ants enter premises such as window and door sills. Spray hills and runways outdoors.

Caddisflies, Cigarette Beetles, Dermestids, Drugstore Beetles, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Saw-toothed Grain Beetles and Weevils (exposed stages) – Thoroughly treat floors and the walls of the infested area with attention to cracks, crevices, corners and perimeters. Sanitation and dust control are of primary importance. Fumigate or destroy infested products.

Silverfish, Firebrats and Paperlice — Apply to localized areas infested by these insects.

Brown Dog Ticks — Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks, and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. *Do not spray animals.*

Clover Mites, Crickets, Earwigs, Millipedes, Centipedes, Sowbugs, Scorpions and Spiders – Thoroughly spray the interior and exterior perimeter of the infested premises. For Clover Mites give special attention to window sills.

Fleas, Flies, Gnats, Mosquitoes, Punkies and Sandflies (outdoors only) – Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios and other areas where insects may enter.

FOR USE IN MEAT PACKING AND FOOD PROCESSING PLANTS: Use only in the inedible product areas of meat packing plants, including those that are federally inspected. Any exposed food product, equipment, or packaging materials must be removed or covered before spraying. Rinse all equipment with water before resuming operations. Treat food processing plants early in the season before canning or other processing begins. Repeat applications, if necessary, when plant is not in operation. Spraying unloading docks and outside areas where insects congregate will aid in preventing re-infestation. Good sanitation and proper disposal of wastes is essential in any effective insect control program.

INSECT CONTROL IN RESEARCH FACILITIES: The use of Residual Spray is used around research animal quarters for insect control will not induce changes in liver microsomal enzyme activity in these animals.

RESTRICTIONS: Do not use as a space spray. Provide adequate ventilation of area being treated. Do not allow children to contact treated areas. Wash hands thoroughly after use. Do not store near food or freshly mowing equipment, as well as handling surfaces. Do not apply to humans. Remove pets and livestock from area before spraying. Do not spray plants.

CAUTION: May be harmful if swallowed, inhaled, or absorbed, except in the skin. Do not get in eyes, on skin, or on clothing. In case of contact, immediately wash with water. Wash hands, arms, and face thoroughly with soap and water. If swallowed, do not induce vomiting. Drink milk or water and get medical attention.

TO: Mr. [redacted] - Campaign office; [redacted]

...and the **best** place **away from** the city.

NET CONTENTS

MECHANICAL,
IMPLIED,
AND OTHER

nominal 100% in accord with the statements made on this subject.