

Camera man's Note  
Poor Copy

NET WEIGHT 22 LBS.



**Nu-turf**  
**20-10-5**  
**FEED & WEED**

MINIMUM GUARANTEED ANALYSIS

LAWN FOOD	
Total Nitrogen (N)	20.00%
4.0% Ammoniacal Nitrogen	
16.0% Urea Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	10.00%
From Phosphoric Acid	
Soluble Potash (K <sub>2</sub> O)	5.00%
From Murates of Potash	
Potential acidity 900 lbs. calcium carbonate equivalent per ton.	
HERBICIDAL	
ACTIVE INGREDIENTS	
Dimethylamine Salts of	
2,4-Dichlorophenoxyacetic Acid	0.24%
Isocetyl Ester of Silvex	
2,2,4-Tribromophenoxy Propionic Acid	0.58%
INERT INGREDIENTS	
	99.18%
Total	100.00%

\* Analyzed to 100% 2,4-Dichlorophenoxyacetic Acid  
\* Analyzed to 41% Silvex

A Product of  
**FEDERAL CHEMICAL COMPANY**  
LOUISVILLE, KENTUCKY 40202  
A Subsidiary of TEXACO INC.

**INSTRUCTIONS**


This product is a 20-10-5 formula lawn food for feeding grasses while controlling broad leaf weeds such as dandelion, plantain, buckhorn, pigweed, ragweed, sourgrass and many others. Can be used in Spring, Summer and Fall. Best results will be obtained when weeds are growing vigorously.

1. Spread the contents of this bag over 4,400 sq. ft.
2. Apply when grass is wet from dew or sprinkling.
3. Do not apply when rain is predicted and do not water the lawn for 24 hours after application.
4. Do not treat new lawns until after third mowing.
5. Do not apply on windy days to avoid drift to surrounding plants.

Do not apply to trees, shrubs, and other desirable plants.

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
This product contains 2,4-Dichlorophenoxyacetic Acid which is a weed killer. It is highly toxic to fish, birds, and other animals.  
**PRECAUTIONS AND DIRECTIONS**  
Do not use on lawns until after third mowing. Do not use on trees or shrubs.  
Labels may be printed on other bags of this product.

THE Quality Line



**Nu-turf**



**Feeds Lawns and Controls Weeds**  
such as PLANTAIN-DANDELION-BUCKHORN

**COVERS 4,400 SQ. FT. (44'x100')**

**INGREDIENT STATEMENT**  
NUTRITIONAL VALUE  
Total Nitrogen (N) 20.00%  
4.0% Ammoniacal Nitrogen  
16.0% Urea Nitrogen  
Available Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>) 10.00%  
From Phosphoric Acid  
Soluble Potash (K<sub>2</sub>O) 5.00%  
From Murates of Potash  
Potential acidity 900 lbs. calcium carbonate equivalent per ton.  
HERBICIDAL  
ACTIVE INGREDIENTS  
Dimethylamine Salts of  
2,4-Dichlorophenoxyacetic Acid 0.24%  
Isocetyl Ester of Silvex  
2,2,4-Tribromophenoxy Propionic Acid 0.58%  
INERT INGREDIENTS 99.18%  
Total 100.00%  
\* Analyzed to 100% 2,4-Dichlorophenoxyacetic Acid  
\* Analyzed to 41% Silvex

ACCEPTED

20-10-5




FEED

& WEED

WEED


FEED



20-10-5

302 MAY 1974  
REGISTERED IN U.S. PAT. OFFICE  
PESTICIDE REG. NO. 250


20-10-5



FEED & WEED

&

FEED & WEED



Nu-turf



Feeds Lawns and Controls Weeds such as PLANTAIN-DANDELION-BUCKHORN

COVERS 4,400 SQ. FT. (44'x100')

INGREDIENT STATEMENT... ACTIVE INGREDIENTS... DIMETHYLAMINE SALTS OF 2,4-DICHLOROPHENOXYACETIC ACID... 0.24%... 2,4,5-TRICHLOROPHENOXY PROPIONIC ACID... 0.58%... NET WEIGHT 22 LBS.

ACCEPTED JUL 29 1974

FEED & WEED

FEED & WEED



20-10-5



MINIMUM GUARANTEED ANALYSIS

Table with 2 columns: Component and Percentage. Includes Total Nitrogen (20.00%), Available Phosphoric Acid (10.00%), Soluble Potash (5.00%), and Potential acidity.

INGREDIENT STATEMENT

Table with 2 columns: Ingredient and Percentage. Includes Dimethylamine Salts of 2,4-Dichlorophenoxyacetic Acid (0.24%), 2,4,5-Trichlorophenoxy Propionic Acid (0.58%), and Inert Ingredients (98.48%).

A Product of FEDERAL CHEMICAL COMPANY LOUISVILLE, KENTUCKY 40202 A Subsidiary of TEXACO INC.


INSTRUCTIONS

This product is a 20-10-5 formula lawn food for feeding grasses while controlling broad leaf weeds such as dandelion, plantain, buckhorn, pigweed, ragweed, sourgrass and many others.

- 1. Spread the contents of this bag over 4,400 sq. ft.
2. Apply when grass is wet from dew or sprinkling.
3. Do not apply when rain is predicted and do not water the lawn for 24 hours after application.
4. Do not treat new lawns until after third mowing.
5. Do not apply on windy days to avoid drift to surrounding plants.
6. Avoid application around or near evergreen, shrubs and other flowering plants.

CAUTION: Keep out of reach of children... SPECIAL FORMULATION FOR... E.P.A. NO. 102881

Table with 3 columns: Area, Suggested Spreader Settings, and Rate. Includes areas like Lawns, Parks, and Golf Courses.



Company Name

20-10-5




FEED & WEED

FEED & WEED

FEED & WEED

THE Quality Line



Nu-turf

FEED & WEED

feeds lawns controls weeds

20-10-5

Feeds Lawns and Controls Weeds such as PLANTAIN-DANDELION-DUCKHOORN

COVERS 4,400 SQ. FT. (44'x100')

INGREDIENT STATEMENT  
HERBICIDAL  
ACTIVE INGREDIENTS  
Diquat Dibromide 0.74%  
2,4-Dichlorophenoxyacetic Acid 0.18%  
2,4,5-Tri-chlorophenoxyacetic Acid 0.08%  
2,4,6-Trichlorophenoxyacetic Acid 0.00%

CAUTION: See label for details of use and other information.  
NET WEIGHT 22 LBS.


ACCEPTED  
JUL 22 1974

FEED & WEED



20-10-5

NET WEIGHT 22 LBS



Nu-turf  
20-10-5  
FEED & WEED

MINIMUM GUARANTEED ANALYSIS  
Total Nitrogen N 20.00%  
Total Phosphorus P 10.00%  
Total Potash K<sub>2</sub>O 5.00%

INGREDIENT STATEMENT  
HERBICIDAL  
ACTIVE INGREDIENTS  
Diquat Dibromide 0.74%  
2,4-Dichlorophenoxyacetic Acid 0.18%  
2,4,5-Tri-chlorophenoxyacetic Acid 0.08%  
2,4,6-Trichlorophenoxyacetic Acid 0.00%

FEDERAL CHEMICAL COMPANY  
1005 HILL, ST. LOUIS, MO. 63102  
A Subsidiary of TEXACO INC.

INSTRUCTIONS  
This product is a 20-10-5 formula lawn feed for feeding grasses while controlling broad leaf weeds such as dandelion, plantain, buckhorn, ground, ragweed, sorgrass and many others. Can be used in Spring, Summer and Fall. Best results will be obtained when weeds are growing vigorously.

- 1. Spread the contents of this bag over 4,400 sq. ft.
- 2. Apply when grass is wet from dew or sprinkling.
- 3. Do not apply when rain is predicted and do not water the lawn for 24 hours after application.
- 4. Do not treat new lawns until after third mowing.
- 5. Do not apply on windy days to avoid drift to surrounding plants.
- 6. Avoid application around or near evergreens, shrubs and other flowering plants.

PRECAUTIONS  
Keep out of reach of children. Avoid contact with eyes, nose, mouth, and clothing. Wash hands after use. Do not eat, drink, or smoke while using this product. Do not use in areas where children or pets play.

DO NOT REAPPLY UNTIL AFTER 48 HOURS IF RAINING AT THE TIME DO NOT REAPPLY  
SUGGESTED SPRINKLER SETTINGS  
HEADS 8 FEET 10 MIN  
HEADS 12 FEET 15 MIN  
HEADS 16 FEET 20 MIN  
HEADS 20 FEET 25 MIN  
HEADS 24 FEET 30 MIN  
HEADS 28 FEET 35 MIN  
HEADS 32 FEET 40 MIN  
HEADS 36 FEET 45 MIN  
HEADS 40 FEET 50 MIN  
HEADS 44 FEET 55 MIN  
HEADS 48 FEET 60 MIN  
HEADS 52 FEET 65 MIN  
HEADS 56 FEET 70 MIN  
HEADS 60 FEET 75 MIN  
HEADS 64 FEET 80 MIN  
HEADS 68 FEET 85 MIN  
HEADS 72 FEET 90 MIN  
HEADS 76 FEET 95 MIN  
HEADS 80 FEET 100 MIN

U.S. DEPARTMENT OF AGRICULTURE  
 NATIONAL BUREAU OF PLANT INDUSTRY  
 WASHINGTON, D. C.

**ANALYSIS**

**Active Ingredients**

Dichloroacetic Salt of 2, 4-Dichlorophenoxyacetic Acid.....	2.2%
Isopropyl Ester of Silvex (2-(2, 4, 6-Trichlorophenoxy) Propionic Acid) .....	.7%
<b>Inert Ingredients</b> (other than herbicidal) Versalene .....	97.1%
<b>Total</b> .....	<b>100.0%</b>

- \* Equivalent to 1.82% 2, 4-Dichlorophenoxyacetic Acid
- \*\* Equivalent to 0.52% Silvex

U.S.D.A. Registration No. 5208-L

**CAUTION:** Keep out of reach of children. Do not contaminate waters used for domestic or irrigation purposes. Avoid inhalation and contact with skin, eyes and clothing, or contamination of food stuffs. Wash hands after using. Do not store near fertilizers, insecticides, or fungicides. Do not allow dust to drift onto susceptible plants such as ornamentals, vegetables, etc. Vapors from this product may cause injury to susceptible plants growing in the immediate vicinity. Do not apply to stoloniferous grasses (those with underground stems) such as fescues, Bermuda or St. Augustine. Do not use on clover if you desire it to remain in the lawn. Certain desirable grasses may exhibit slight browning but will recover after a short period if the product has been used as directed.

(Back of Container)  
**Directions for Use**

**M. N. A. WEED KLEEN** is a combination of 2, 4-D Acid and Silvex on a light weight inert carrier. Its function is to control broadleaf weeds such as:

Biantin	Dandelion	Wild Carrot
Buckhorn	Common Chickweed	Wild Onion
Thistle	Houseleaved Chickweed	Pig Weed
Rag Weed	Sour Grass	Lambs Quarters
White Clover	Clover	and many others

The combination of ingredients is very effective, economical and easy to use and due to the dry granular form is safer to use than a spray.

**M. N. A. WEED KLEEN** Can be used in Spring, Summer or Fall. Best results are obtained when weeds are growing vigorously. When using this product strive to distribute it uniformly and to retain the granules on the leaves of weeds. The following suggestions will assist:

1. The contents of this bag are to cover a 5,000 sqft area (50x100)
2. This can best be accomplished using a spreader. Apply between mowings to provide sufficient leaf surface to be treated.
3. Sprinkle lawn prior to application or apply after a light rainfall.
4. Delay application when rain is predicted.
5. Do not water lawn for 24 hours after application.
6. Avoid application on windy days to avoid drift to surrounding plants
7. Do not treat newly seeded lawns until grasses are well established or at least until after third mowing.

