

ACCEPTED
MAR 3, 1971
5298-11

50 LBS. NET GRANULAR CRAB CONTROL

GRANULAR CRABGRASS KILLER CONTAINING TUPERSAN®

GENERAL INFORMATION: "Granular Crab Control" is an easy to apply granular product for selective pre-emergence control of certain annual weed grasses in turf areas such as golf course fairways, lawns, parks, roadsides and turf grown for grass seed or sod production. It is highly effective for the control of crabgrass (smooth and hairy) and may be used on newly seeded areas as well as on established turf. Because of its high degree of selectivity, "Granular Crab Control" may be applied at time of seeding without causing injury to germinating seeds of many common temperate region grasses.

DIRECTIONS: "Granular Crab Control" is recommended for pre-emergence control of annual weed grasses, such as crabgrass (smooth and hairy) fox-tail and barnyard grass, in newly seeded or established plantings of the following grasses only: bluegrass, fescue, red top, smooth brome, perennial ryegrass, orchard grass, zoysia, and the following strains of bentgrass: Penncross, Seaside, Highland, Astoria, Nimisila, C-1, C-7 and C-19. Do not use on other bentgrass strains nor on Bermudagrass, as injury to turf may result. The effect of "Granular Crab Control" on golf course greens may vary with turf management practices and environmental conditions; therefore, its use on golf greens is not recommended, as injury to turf may result. Treatment will not control annual bluegrass (*Poa annua*), clover, or most broadleaf weeds.

CAUTION

KEEP OUT OF REACH OF CHILDREN

May irritate eyes, nose, throat and skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash after using.

"Granular Crab Control" may be applied at any stage of turf development, including new seedings. At least 1/2 inch of water must be provided by rainfall or irrigation within 3 days after treatment in order to carry the chemical into the root zone of germinating weeds.

AMOUNT TO USE — WHEN TO APPLY

NEW SPRING PLANTINGS: After soil has been seeded apply at a rate of 5 lbs. per 1,000 sq./ft. For best results do not work the soil after application. At least 1/2 inch of water must be provided by rainfall or sprinkler within 3 days after treatment in order to carry the chemicals into the root zone of germinating weeds.

Manufactured by
OLD FOX AGRICULTURAL SALES, INC.
East Providence, Rhode Island

Printed in U.S.A. Made in U.S.A.

NEW FALL PLANTINGS AND ESTABLISHED TURF: Apply at a rate of 10 lbs. per 1,000 sq./ft. and provide at least 1/2 inch of water to carry the chemical into the root zone of germinating weeds. This product can be applied anytime in April or May but prior to the expected emergence of crabgrass.

Note: Use only on turf areas; do not apply to crops used for food or feed purposes.

ACTIVE INGREDIENTS:

Siduron (1-(2-Methylcyclohexyl)-3-phenylurea) 2.3%

INERT INGREDIENTS 97.7%

TOTAL 100.0%

WARRANTY: The manufacturer makes no representation or warranty except as appears on this label, and shall not be responsible for any injury to person or property resulting from improper use.

U.S.D.A. Registration No. 5298-11