



ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl
N-methylcarbamate*)9.5%
Malathion**8.0%
Folpet N-(Trichloromethyl-
thio) phthalimide14.5%
1,1 Bis(chlorophenyl)-
2,2,2-trichloroethanol***1.5%

INERT INGREDIENTS:66.5%
TOTAL 100.0%

*Otherwise known as SEVIN®
Union Carbide Corp.

**O,O-Dimethyl dithiophosphate
of diethyl mercaptosuccinate.

***Otherwise known as KELTHANE®
Rohm & Haas Co.

EPA Reg. No. 5330-4

KILLS GARDEN PESTS

Such as
JAPANESE BEETLES • APHIDS • MITES • CERTAIN OTHERS
CONTROLS MANY GARDEN DISEASES

Such as
POWDERY MILDEW • BLACK SPOT • CERTAIN OTHERS

STERN'S
MIRA-GARD®
MULTI-PURPOSE
ROSE & GARDEN
SPRAY

Trade Name

INSECTICIDE • FUNGICIDE • MITICIDE

Net Weight 12 oz. Av.

For Use On
ROSES • FLOWERS • VEGETABLES
FRUITS • TREES • SHRUBS

CAUTION: KEEP OUT OF REACH OF CHILDREN
see back panel for complete caution statement

Satisfaction Guaranteed By
STERN'S NURSERIES, INC.
Geneva, New York 14456

CAUTION:

KEEP OUT OF REACH OF CHILDREN. HAZARDOUS IF INHALED OR SWALLOWED. May cause irritation of nose, throat and skin. Avoid breathing or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after using. Store away from domestic animals. Avoid contamination of feed and foodstuffs. In case of contact, flush with plenty of water; for eye contact, seek medical attention.

This product is highly toxic to bees exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift into areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product as specified on this label.

Do not apply to Fruit Trees within 7 days of harvest. Do not apply to Vegetables within 2 days of harvest — especially on Tomatoes. Not for use on Pecans, Cotton, Artichokes and Avocados.

NOTICE: Seller's guarantee shall be limited to the contents of the label, and subject thereto the buyer assumes all risk to persons or property arising out of use or handling and accepts this product on these conditions.

Do not reuse container. Destroy when empty.

CAUTION:

KEEP OUT OF REACH OF CHILDREN. HARMFUL IF INHALED OR SWALLOWED. May cause irritation of nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after using. Store away from domestic animals. Avoid contamination of feed and foodstuffs. In case of contact, flush with plenty of water; for eyes, get medical attention.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not apply to Fruit Trees within 7 days of harvest. Do not apply to Vegetables within 2 days of harvest — 5 days on Tomatoes. Not for use on Pecans, Cotton, Artichokes and Avocados.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts this product on these conditions.

Do not reuse container. Destroy when empty.

Before using read carefully
Complete directions inside package

DIRECTIONS FOR USE

APPLY ON — Roses, Flowers, Shrubs, certain Fruit Trees and certain Vegetables by spraying at 7 to 10 day intervals and after heavy rains. It is best to start before insects become numerous and before diseases become apparent. For timing of applications on Fruit Trees, follow Fruit Tree Spray Schedule enclosed in this package or consult your State Agricultural Experiment Station or State Extension Service Specialists.

DILUTION RATE — Mix 3 tablespoons of spray material per gallon of water. (Use handy measuring spoon enclosed.)

SPRAYING — For Hand Sprayers, Tank-Type Sprayers and Sprinkling Can, follow dilution instruction above. Agitate while spraying to insure even coverage. For Hose-End Jar Type Sprayer units, mix the above rate of spray material with the level of water recommended by the manufacturer of your sprayer. (Example — for two gallons of finished spray, mix 6 level tablespoons with the level of water recommended and indicated on your sprayer.) Then attach to your hose and spray. Agitate while spraying to insure even coverage.

* * * * *

GUARANTEED BY STERN'S NURSERIES

Satisfaction
or
Money Back

At any time within one year, if you are not satisfied with results obtained from this product, return the unused portion to STERN'S NURSERIES, INC., Geneva, New York. Your purchase price will be promptly refunded.

RECOMMENDED USES

ROSES & FLOWERS

For control of Japanese Beetles, Aphids, Spider Mites, Lace Bugs, Scale, Leafhoppers, and other insects. For control of Black Spot, Powdery Mildew, and other diseases. For control of Japanese Beetles, Aphids, Spider Mites, Lace Bugs, Scale, Leafhoppers, and other insects. For control of Black Spot, Powdery Mildew, and other diseases.

DISEASES—For control of Black Spot, Powdery Mildew, and other diseases. For control of Japanese Beetles, Aphids, Spider Mites, Lace Bugs, Scale, Leafhoppers, and other insects.

SHRUBS

INSECTS—For control of Japanese Beetles, Aphids, Spider Mites, Lace Bugs, Scale, Leafhoppers, and other insects. For control of Black Spot, Powdery Mildew, and other diseases.

CAUTION

May be harmful if swallowed or inhaled. May cause irritation of nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after using. Store away from domestic animals. Avoid contamination of feed and foodstuffs. In case of contact, flush with plenty of water; for eyes, get medical attention. **KEEP OUT OF REACH OF CHILDREN.**

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

STERN'S NURSERIES, INC.
Geneva, New York

ACTIVE INGREDIENTS

Carbaryl (1-Naphthyl N-methylcarbamate*) 9.5%
Malathion** 8.0%
Folpet N-(Trichloromethylthio) phthalimide 14.5%
1,1-Bis(chlorophenyl)-2,2,2-trichloroethanol*** 1.5%

INERT INGREDIENTS 66.5%
TOTAL 100.0%

*Otherwise known as SEVIN® Union Carbide Corp.

**O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate

***Otherwise known as KELTHANE® Rohm & Haas Co.

EPA Reg. No. 5330-4

INSECTS—For control of Spider Mites, Scale, Leafhoppers, Holly Leaf Miners, European Spruce Sawfly, Tent Caterpillars, Apple Periodical Cicada, Elm Leaf Miner, and Oak Leaf Miners on PINE, SPRUCE, HOLLY, ELMS, OAKS, MAGNOLIAS, MAPLES and SYCAMORES.

TREES

INSECTS—For control of Spider Mites, Scale, Leafhoppers, Holly Leaf Miners, European Spruce Sawfly, Tent Caterpillars, Apple Periodical Cicada, Elm Leaf Miner, and Oak Leaf Miners on PINE, SPRUCE, HOLLY, ELMS, OAKS, MAGNOLIAS, MAPLES and SYCAMORES.

FRUIT TREES

INSECTS—For control of Spider Mites, Scale, Leafhoppers, Holly Leaf Miners, European Spruce Sawfly, Tent Caterpillars, Apple Periodical Cicada, Elm Leaf Miner, and Oak Leaf Miners on PINE, SPRUCE, HOLLY, ELMS, OAKS, MAGNOLIAS, MAPLES and SYCAMORES.

Mites, Pear Psylla, Tent Caterpillars, Apple Mites, and Banded Leaf-Roller. Periodical Cicada on APPLES and SOUR CHERRIES.

DISEASES—Apple Scab, Spot, Bitter Rot, Leaf Spot on APPLES and CHERRIES.

SUGGESTED SPRAY SCHEDULE

FRUIT

Delayed Dormant—(From first green on bud to 1/2 inch leaves on Apple).

Pre-bloom—(From first pink to first bloom on APPLE and CHERRY).

Petal Fall—(When last petals have fallen on all fruits listed above).

Cover Sprays—(2 to 4 applications). Begin 10 days after petal fall on APPLE.

Post Harvest Spray—For Cherry Leaf Spot apply immediately after harvest and 10 days later.

CAUTIONS: Do not apply to fruit trees within 7 days of harvest and on Sour Cherries make one application and then not again for 30 days.

May be injurious to some varieties of apples. Not for use on Pecans, Cotton, Artichokes and Avocados.

VEGETABLES

TOMATOES—For the control of worms, Aphids, Spider Mites, Beetles, Colorado Potato Beetle, Anthracnose.

CUCUMBERS — SUMMER
For control of Pickleworms, Cucumber Beetles, Bugs, Flea Beetles, Downy Mildew and Anthracnose.

CAUTION: Do not apply to within 2 days of harvest-Tomatoes.



STERN'S
MIRA-GARD
MULTI-PURPOSE
ROSE & GARDEN
SPRAY

Ⓢ Trade Name

FOR USE ON

ROSES • FLOWERS • VEGETABLES • FRUITS • TREES •

APPLY ON—Roses, Flowers, Shrubs, certain Fruit Trees and certain Vegetables by spraying at 7 to 10 day intervals and after heavy rains. It is best to start before insects become numerous and before diseases become apparent. For timing of applications on Fruit Trees follow Fruit Tree Spray Schedule or consult your State Agricultural Experiment Station or State Extension Service Specialists.

DILUTION RATE—Mix 3 tablespoons of spray material per gallon of water.

SPRAYING—For Hand Spray Type Sprayers and Sprinklers use low dilution instructions and spray while spraying to insure coverage. For Hose-End Sprayer units, mix the above spray material with the level recommended by the manufacturer (Example—four gallons of finished spray, mix with 12 tablespoons with the level recommended and indicate on sprayer). Then attach to hose and spray. Agitate while spraying to insure even coverage.

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROBOTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO.

POISON SIGNS (Symptoms)

[illegible]

SPECIM

**KILLS
FLIES
FAST!**

WARNING:

KEEP OUT OF REACH OF CHILDREN

See side panel for additional precautions



**FLY
FIGHTER**
Double Strength

FLY BAIT

Active Ingredient:	% By Wt.
2,2-dichlorovinyl dimethyl phosphate	0.93
Related compounds07
Inert Ingredients	99.00
	<hr/>
	100.00%

FLIES FEED AND DIE

5481-9

Accepted