

SEP 19 1967
UNDER THE FEDERAL
FUNGICIDE ACT
FOR ECONOMIC
FE UNDER 5362-B
TO ATTACH

DIRECTIONS

Wash and rinse prior to sanitizing. To sanitize milk utensils, equipment use 1 oz. Solventol #310 in 7 gallons water (100 ppm available chlorine). Brush, immerse, spray or circulate (CIP). Rinse with water. If no facilities are available to determine a reduction to less than 50 ppm. available chlorine in the sanitizing solutions, start with a 200 ppm available chlorine solution.

NOTE: Contact time. The laws of some states require that utensils and equipment remain in contact with the germicidal solution for at least 2 minutes. Solventol #310 fulfills the criteria of Appendix F as revised March 12, 1966, the Milk Ordinance and Code, 1953 recommendations of the U.S. Public Health Service when tested by Chambers method. Replace cover store in cool, dry place.

Solventol. **SOLO SAN** **310**

CHLORINE - SANITIZER

GLASSES - DISHES - UTENSILS

CAUTION:

KEEP OUT OF REACH OF CHILDREN
(SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS)

NET WEIGHT

SOLVENTOL CHEMICAL PRODUCTS

ACTIVE INGREDIENT

[(MONOTRICHLO
(MONOPOTA

PENTA-ISOCYA
K₄CL₁₁ (C

INERT INGREDIENT

CAUTION

KEEP OUT OF

HARMFUL IF S
CONTAMINATI

IN CASE OF C
EYES, FLUSH

U.S.D.A

SEP 19 1967

UNDER THE FEDERAL FOOD, DRUG & COSMETIC ACT
FUNGICIDE
FOR ECONOMIC PEACE
FD UNDER 5362-13
TO ATTACH

Solventol.

SOLO SAN

310

CHLORINE - SANITIZER

GLASSES - DISHES - UTENSILS

CAUTION:

KEEP OUT OF REACH OF CHILDREN

(SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS)


NET WEIGHT

SOLVENTOL CHEMICAL PRODUCTS

ACTIVE INGREDIENTS

[(MONOTRICHLORO) - TETRA
(MONOPOTASSIUM DICHLORO)]

PENTA-ISOCYANURATE



16%

INERT INGREDIENTS

84%

100%

CAUTION:

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED, AVOID
CONTAMINATION OF FOOD.

IN CASE OF CONTACT WITH SKIN OR
EYES, FLUSH WITH PLENTY OF WATER

U.S.D.A. REG. #5362-13

ACCEPTED
 SEP 19 1967
 UNDER THE FEDERAL FOOD, DRUG & COSMETIC ACT
 FUNGICIDE
 FOR ECOLOGICAL CONTROL
 ED UNDER 362B
 TO ATTACH

DIRECTIONS

Wash and rinse prior to sanitizing. To sanitize milk utensils, equipment use 1 oz. Solventol #310 in 7 gallons water (100 ppm available chlorine). Brush, immerse, spray or circulate (CIP). Rinse with water. If no facilities are available to determine a reduction to less than 50 ppm. available chlorine in the sanitizing solutions, start with a 200 ppm available chlorine solution.

NOTE: Contact time. The laws of some states require that utensils and equipment remain in contact with the germicidal solution for at least 2 minutes. Solventol #310 fulfills the criteria of Appendix F as revised March 12, 1966, the Milk Ordinance and Code, 1953 recommendations of the U.S. Public Health Service when tested by Chambers method. Replace cover, store in cool, dry place.

Solventol.
SOLO SAN
310
CHLORINE - SANITIZER

GLASSES - DISHES - UTENSILS

CAUTION:
KEEP OUT OF REACH OF CHILDREN
 (SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS)

NET WEIGHT
 SOLVENTOL CHEMICAL PRODUCTS, Inc.
 Romulus, Michigan

ACTIVE INGREDIENTS

[(MONOTRICHORO) - TETRA
 (MONOPOTASSIUM DICHORO)]

PENTA-ISOCYANURATE
 $K_4CL_{11}(CNO)_{15}$ 16%

INERT INGREDIENTS 84%
 100%

CAUTION:

KEEP OUT OF REACH OF CHILDREN

**HARMFUL IF SWALLOWED, AVOID
 CONTAMINATION OF FOOD.**

**IN CASE OF CONTACT WITH SKIN OR
 EYES, FLUSH WITH PLENTY OF WATER**

U.S.D.A. REG. #5362-13

8