DIRECTIONS FOR USE

EQUIPMENT SANITATION

Preclean equipment with cold or lukewarm water. Fill with solution 2 oz. Solo San I20 to IO gal. water. (Provides 25 ppm titratable iodine.) When ready to use equipment, drain off Solo San I20 solution and flush with water.

TANK AND VAT SANITATION

Preclean with water (not hot). Add solution 2 oz. Solo San 120 to 10 gal. water. (Provides 25 ppm titratable iodine). After five minutes, drain off solution and flush with water.

SANITATION OF WALLS, FLOORS, CONVEYORS, ETC.

Clean daily with solution 2 oz. Solo San 120 to 10 gal. water (25 ppm titratable iodine.)
Clean weekly or at batch run end with solution 4 oz. Solo San 120 to 10 gal. water (50 ppm titratable iodine) and rinse.

SANITATION OF BEVERAGE CONTAINERS

Follow local Health Department regulations, which usually suggest use-solutions in the range of the lower ppm levels shown in the Dilution Table presented on label.

SOLVENTOL®

SOLO SAN 120

lodophor Germicidal Detergent

ACTIVE INGREDIENTS

Butoxy Polypropoxy Ethanol- 12.54%

Iodine Complex

(1.6% Titratable Iodine)

INERT INGREDIENTS

87.46%

100.00%

CAUTION:

Keep out of reach of children. (See side panel for warnings)

Manufactured by:

SOLVENTOL CHEMICAL PRODUCTS, INC. Romulus, Michigan

USDA REG. NO._____

1 GALLON

DIRECTIONS FOR DILUTION

Titratable lodine

¹/₂ oz. Solo San 120 per 10 gal. water = 6¼ppm

1 oz. Solo San 120 per 10 gal. water = 12½ppm

2 oz. Solo San 120 per 10 gal. water = 25ppm

4 oz. Solo San 120 per 10 gal. water = 50ppm

WARNING:

Do not allow concentrated Solo San 120 to come into contact with food.

Harmful if swallowed.

Avoid contact with skip and eyes. In case

Avoid contact with skin and eyes. In case of contact with eyes, flush immediately with water; get medical attention.

ACT. ZDI

1194 5,093

536274