

FOR USE

ELEVATORS ATTENTION

For Best Results Follow Directions

HOW TO APPLY

Apply fumigant when bin is being filled during grain filling. Pour fumigant rapidly directly onto the grain surface. Apply at intervals, so that total application is made once for each ten bushels of grain. Add an extra 2 to 4 gallons per 1,000 bushels entering the bin. Seal the bin after treatment. If possible, move the grain to a new bin at least 72 hours. Best results are obtained if treated grain is left in bin as long as possible. **AVOID TOXIC GASES. DO NOT ENTER BIN DURING OR AFTER TREATMENT.** Best treated grain may be used for human consumption.

AVERAGE DOSAGE TABLE

Gallons Per 1,000 Bushels Grain			
Type Bin	Wheat	Rye	Oats
Steel and Concrete	2	5	5
Wooden	4	7	7

TREATMENT OF RICE

For treatment of rice, use 4 gallons per 1,000 bushels of grain. Apply fumigant directly to the grain surface. Seal the bin after treatment. If possible, move the grain to a new bin at least 72 hours. Best results are obtained if treated grain is left in bin as long as possible. **AVOID TOXIC GASES. DO NOT ENTER BIN DURING OR AFTER TREATMENT.** Best treated grain may be used for human consumption.

INCREASE DOSAGES

- For Slow Bin Filling Time
- For High Moisture Grain
- For High Dockage
- For High Foreign Matter

BOXCAR TREATMENT


Apply fumigant when grain is being filled during grain filling. Pour fumigant rapidly directly onto the grain surface. Apply at intervals, so that total application is made once for each ten bushels of grain. Add an extra 2 to 4 gallons per 1,000 bushels entering the bin. Seal the bin after treatment. If possible, move the grain to a new bin at least 72 hours. Best results are obtained if treated grain is left in bin as long as possible. **AVOID TOXIC GASES. DO NOT ENTER BIN DURING OR AFTER TREATMENT.** Best treated grain may be used for human consumption.

NOTICE TO BUYER

Buyer should inspect grain for signs of infestation before purchase. If infestation is found, the grain should be treated with a suitable fumigant before use. The use of this fumigant is recommended for the treatment of grain infested with weevils, flour beetles, and other insects.


5 GAL. NET

Vulcan



TIME-TESTED FORMULA FORTIFIED WITH 7.2% ETHYLENE DIBROMIDE

KILLS ADULT AND EGG STAGES of Grain Weevil, Rice Weevil, Lesser Grain Borer, Confused Flour Beetle, Angoumois Grain Moth Larva, Indian Meal Moth Larva, Mediterranean Flour Moth Larva, Cadelle, Black Carpet Beetle, Sawtooth Grain Beetle and certain other insects attacking stored cereal grains.



Vulcan Materials Company

U.S. A. Registration No. 51821

ATTENTION FARMERS

For Best Results Follow Directions

BEFORE FUMIGATION

FIRST Make the bin gas-tight. Close all cracks and openings in floor and walls of bin. Caulk or tighten, by nailing strips of wood, over all holes or cracks. Paper, if needed, with asphalt type building paper on either outside or inside of bin. Grain fumigants work best when bin is gas-tight. **SECOND** Remove all moldy or crusted layers from the surface of stored grain. Level grain surface and keep surface at least 6 in. below wall top.

HOW TO APPLY

Apply fumigant directly to the grain surface in a very wet, coarse, fan-shaped stream. A fast, uniform distribution of fumigant over the entire grain surface gives best results. If possible, apply fumigant from outside of bin to avoid toxic fumes. Seal bin after treatment and keep record. Inspect monthly for signs of re-infestation and re-treat as needed. Do not re-enter treated bin without an airline respirator or a full face gas mask with a black canister for protection against vapors.

AVERAGE DOSAGE TABLE

Gallons Per 1,000 Bushels Grain		
Type Bin	Wheat, Rye, Oats	Corn
Steel and Concrete	2	5
Wooden	4	7

Do not breathe vapors. Avoid contact with skin, eyes, nose, mouth, and clothing. Wash thoroughly with soap and water after use. Store in a cool, dry place. Keep away from children and animals.

Keep away from children and animals. Do not breathe vapors. Avoid contact with skin, eyes, nose, mouth, and clothing. Wash thoroughly with soap and water after use. Store in a cool, dry place. Keep away from children and animals.

