

WARNING: May Produce Delayed Chemical Burns. Do not breathe dust or spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after handling. Keep children and pets off treated areas until dry.

USDA Reg. No. 7453-1

ANTIDOTE: If swallowed. Give milk or white of egg beaten with water, then a tablespoon of salt in a glass of warm water, and repeat until vomit fluid is clear. Repeat milk or white of egg beaten with water. Call a Physician immediately.

If on skin. Wash immediately with soap and warm water.

If in eyes. Flush with water for at least 15 minutes, and get medical attention.

CONTAINER DISPOSAL: Decontaminate by repeated rinsing with water. Crush or perforate and bury when empty. NEVER REUSE.

NON-WARRANTY: The fitness of this chemical is not warranted by the producer for any particular use, even though such is known, and the buyer assumes all responsibility (including injury or damage) resulting from its use as such, or in combination with other materials.

DIRECTIONS FOR USE:

TURF GRASS AND LAWNS: For control of bluegrass blight (*Helminthosporium*), cooper spot, dollar spot, fading-out, melting-out, pink patch, zonate eyespot: apply 5 fl. oz. Mersolite 810 5000 sq. ft. in sufficient water to cover, or, 1 fl. oz. 3-10 gal. water 1000 sq. ft.

Begin in early spring or before diseases are expected to appear. Repeat at 7 to 10 day intervals. Do not water or mow for 24 hours after treatment. Do not use on Merion bluegrass. Do not graze treated areas or feed treated clippings to livestock. Do not allow children or pets on freshly treated areas.

MERSOLITE-810

(PHENYL MERCURIC ACETATE)

ACTIVE INGREDIENT:

Phenylmercuric Acetate 10%
(Equivalent to 6% metallic mercury)

INERT INGREDIENT 90%

TOTAL 100%

NET _____

DANGER

POISON  **POISON**

Keep Out of the Reach of Children

See Left Panel For Antidote and Additional Warning Statements

WOOD RIDGE CHEMICAL CORPORATION

PARK PLACE EAST

WOOD-RIDGE, NEW JERSEY 07075

Seed	Disease	Dosage oz. act./bu.
Barley	Covered smut, black loose smut, stripe, seed decay, seedling blights.	.078
Flax	Seed decay, damping-off, root rot	234
Oats	Smut, leaf stripe, seed decay, seedling blights	.078
Rye	Stem smut, covered smut	.078
Sorghum & Millet	Loose and covered kernel, smut, Damping-off, seed decay	.078
Wheat	Bunt, kernel smudge, seed decay, seedling blights.	.078

Do not use treated seed for food, feed, or oil purposes.

Bulbs, seeds and roots - damping-off, seed borne fungus diseases. Dilute 1 oz. Mersolite 810 to 10 oz. with water. Dilute 2 fl. oz. of this 1% solution in 3 gallons water. Soak ground to at least 3 inches below bulbs, seeds, or roots. Follow in 7-10 days with 1 oz. per 3 gallons as foliage and soil spray. Do not apply to food or feed crops.

ACCEPTED

JUN 7 1968

UNDER THE
FUNGICIDE ACT
FOR CONTROL OF
ED UNDER N 7453-1
TO ATTACHED

5385 20