

WARNING: May Produce Delayed Chemical Burns. Do not breathe dust or spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after handling. Keep children and pets off treated areas until dry.

USDA Reg. No. 7453-2

ANTIDOTE: If swallowed. Give milk or white of egg beaten with water, then a tablespoon of salt in a glass of warm water, and repeat until vomit fluid is clear. Repeat milk or white of egg beaten with water. Call a Physician immediately.

If on skin. Wash immediately with soap and warm water.

If in eyes. Flush with water for at least 15 minutes, and get medical attention.

CONTAINER DISPOSAL: Decontaminate by repeated rinsing with water. Crush or perforate and bury when empty. NEVER REUSE.

NON-WARRANTY: The fitness of this chemical is not warranted by the producer for any particular use, even though such is known, and the buyer assumes all responsibility (including injury or damage) resulting from its use as such, or in combination with other materials.

MERSOLITE®-830

(PHENYL MERCURIC ACETATE)

ACTIVE INGREDIENTS:

Phenyl Mercuric Acetate ----- 30%
(Equivalent to 18% metallic mercury)

INERT INGREDIENT ----- 70%

TOTAL 100.0%

NET _____

DANGER



Keep Out of the Reach of Children

See Left Panel For Additional Warning Statements

WOOD RIDGE CHEMICAL CORPORATION

PARK PLACE EAST

WOOD-RIDGE, NEW JERSEY 07075

write "seed" with these

DIRECTIONS FOR USE:

SEED TREATMENTS

Seed	Disease	Dosage oz. act./bu.
Barley	Covered smut, black loose smut, stripe, seed decay, seedling blights.	.078
Flax	Seed decay, damping-off, root rot	.234
Oats	Smut, leaf stripe, seed decay, seedling blights	.078
Rye	Stem smut, covered smut	.078
Sorghum & Millet	Loose and covered kernel smut, damping-off, seed decay	.078
Wheat	Bunt, kernel smudges, seed decay, seedling blights.	.078

Do not use treated seed for food, feed, or oil purposes.

Bulbs, seeds and roots - damping-off seed borne fungus diseases. Dilute 1 oz. Mersolite 830 to 30 oz. with water. Dilute 2 fl. oz. of this 1% solution in 3 gallons water. Soak ground to at least 3 inches below bulb, seed, or roots. Follow in 7-10 days with 1 oz. per 3 gallons as foliage and soil spray. Do not apply to food or food crops.

ACCEPTED
JUN 7 1968
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 7453-2 SUBJECT
TO ATTACHED COMMENTS.

5383-21