

NOTICE: This mixture is made especially for the control of both large Brown Patch and Dollar Spot on golf greens and fine lawns.

DIRECTIONS: The United States Golf Association Green Section recommends the following rates of application mixed with compost, sand or in water for mercurous mercuric chloride mixtures:

Early Spring	3 oz.	/1000 sq. ft.
Summer	2 oz.	" " "
Hot Humid Weather	1 oz.	" " "
Late Fall	4 oz.	" " "

Determine as accurately as possible the number of square feet contained in each green.

WOOD RIDGE MIXTURE 21 may be applied wet or dry with equal results. The method of use should be governed by equipment you have on hand.

URY: Thoroughly mix the required amount with sand or compost. For best results use one pail of screened sand or compost for each 4 oz. of fungicide per 1000 sq. ft. Apply one-half of the batch in one direction; cross with the other half at right angles.

DANGER: KEEP OUT OF REACH OF CHILDREN

Do not breathe dust. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling.

WOOD RIDGE MIXTURE "21"

USDA Reg. No. 7453-18

ACTIVE INGREDIENTS:

Mercurous Chloride (Calomel)	66.7%
Mercuric Chloride (Corrosive Sublimate)	33.3%
Total	100.0%

Mercury Content 81.3%

NET


WOOD RIDGE
CHEMICAL CORPORATION

WOOD-RIDGE, NEW JERSEY

WET: Agitate solution frequently so that the material will not settle. By this method 3 oz. of material may be used with as low as 10 gallons of water per 1000 sq. ft. Always rinse power sprayer with clear water after using to prevent corrosion of the metal parts. If using a barrel cart, increase the water mixture to 50 gallons for 3 oz. per 1000 sq. ft. Stir with wooden paddle during application. When hand sprinklers are used, the preparation should be mixed in wooden, glass or earthenware containers—do not use metal. Metal sprinkling cans may be used for the actual application. Use same ratio as for the barrel cart.

Either method of application should be followed with a good watering in. Do not water too heavily, as it is liable to puddle. Excessive watering will wash the mixture from the high spots and allow a concentrated solution on the low areas, which may cause a discoloration in the turf. In dry applications, the turf may be dragged with a steel mat or poled with bamboo poles in order to increase the evenness of distribution.

ANTIDOTE: Give milk or white of eggs beaten with water, then a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Repeat milk or white of eggs beaten with water. **CALL A PHYSICIAN.**



538 1-2