

SANITERG

USE DIRECTIONS:

DISINFECTION

Use at 1 ounce per gallon of water. This product will be effective against Pseudomonas aeruginosa PRD-10 at this level.

At 1 ounce per gallon of water, this product is fungicidal against pathogenic fungi.

For best results, use SANITERG as directed above with a cloth or mop and apply to walls, floors and other hard surfaces.

For heavy soil or organic matter, a pre-cleaning step is recommended.

SANITIZATION

If used for sanitization of previously cleaned food equipment or food contact items, limit the level of active quaternaries to 200 ppm (1.4 ounces per 5 gallons of water). No Rinse Required at this level.

This product when used on environmental, inanimate, hard surfaces at 1 ounce per gallon of water, is effective against Influenza A2, Herpes Simplex, Adenovirus type 2, and Vaccinia viruses.

Rinse empty container thoroughly with water and discard.

PHENOL COEFFICIENTS:

Guaranteed Minimum

S. aureus

S. tyhposa

sa 48 1

78

CONCENTRATED

- DETERGENT
- SANITIZER
- FUNGICIDE
- DISINFECTANT

がりょ DEODORIZER

ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈)	
dimethyl benzyl ammonium chlorides n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl	4.5%
ethylbenzyl ammonium chlorides	4.5%
Tetrasodium ethylenediamine tetraacetate	2.0%
Sodium Carbonate	4.0%
INERT INGREDIENTS:	85.0%
	100.0%

DANGER

KEEP OUT OF REACH OF CHILDREN. SEE RIGHT SIDE PANEL FOR ADDITIONAL PRECAUTIONARY AND FIRST AID STATE-MENTS.

NET CONTENTS

EPA Reg. No. 5404-6 EPA Est. No. 5404-N.M1 SANITERG is a concentrated detergent, disinfectant, sanitizer, fungicide and deodorizer. It is specifically designed for Institutional use, such as, Hospitals, Nursing Homes, and Schools, where housekeeping will be of prime importance in controlling the hazard of cross-infection from environmental surfaces.

SANITERG is for use on floors, walls and other hard surfaces in Hospitals, Nursing Homes, Hotels, Public Areas, Industrial Areas and in Institutions. It is excellent for Operating Rooms, Rest Rooms, Nurseries and Food Services Areas.

When used as directed, SANITERG is effective against Staphylococcus aureus, Salmonella chloreaesuis and Pseudomonas aeruginosa. SANITERG is also effective against pathogenic fungi, Trichophyton interdigitale.

if used for sanitization of previously cleaned food processing equipment or food contact items, limit the active quaternaries to 200 ppm. At this level, NO POTABLE RINSE is required.

DANGER

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food. Protect eyes and skin when handling concentrate.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN

Probable muci sal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.





P. O. BOX 3309 • ALBUQUERQUE, NEW MEXICO 87110 • (505) 881-6550 P. O. BOX 641 • FARMINGTON, NEW MEXICO 87401 • (505) 325-4132

"Serving the Southwest for Over a Quarter Century"