

SAN FRANCISCO

FORT WORTH

CHICAGO

NEWARK, N.J.

THERMAL FOGGING INSECTICIDE TYPE M

ACCEPTED

JAN 28 1974

UNDER THE FELERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NOT 1/2/2/2 SUBJECT TO ATTACHED COMMENTS.

EPA REG. NO. 5440-106

ACTIVE INGREDIENTS 100.00%

PETROLEUM HYDROCARBONS 96.50%

N-OCTYL BICYCLOHEPTANE
DICARBOXIMIDE 2.50%
*TECHNICAL PIPERONYL BUTOXIDE 0.50%
PYRETHRINS 0.50%
PYRETHRINS 0.50%
*Equivalent to 4% of (Butylcarbityl)(6-Propylpiperonyl)
Ether and 0.1% Related Compounds

JAN 22 1971 E

A thermal fogging compound for use in thermal fogging dispensers, air operated foggers or permanent automizers.

DIRECTIONS FOR USE

CONTACT FOGGING: Exposed stages of cockroaches, silverfish, weevil, spiders, ants, crickets, confused flour beetles and centipedes can be killed only on direct contact. Fog thoroughly into cracks, crevices and other places where insects may hide. Follow with thorough fogging of the area.

SPACE FOGGING: with Dyna-Fog machines, or equivalent: For gnats, flies, meal moths, mosquitos and above listed insects, fog at the rate of ½ ounce per 1,000 cu. ft.

FOR ACCESSIBLE CRAWLING INSECTS (as named above): Fog at the rate of 1 ounce per 1,000 cu. ft. May also be applied with a hand sprayer to harborages, cracks and crevices with a fine mist.

CAUTION: Keep doors and windows closed, turn off air conditioning and ventilating systems, in order to obtain maximum results. Premises should remain closed for at least ten minutes or until fog has disappeared. Repeat all applications as necessary. Do not remain in treated areas. Ventilate the area after treatment is completed. This product is compounded especially for use in food processing plants, dairies, confectionery companies, department stores, schools, institutions and offices. The carrier in this formula will not impart odor or taste when used in food processing plants according to directions.

CAUTION: Avoid exposure to fog. Do not apply to animals. Do not spray or apply on food, food products, utensils, or food handling surfaces. Remove pets, birds and cover fish aquariums before spraying. Use this product as a space spray in food processing plants only when plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy it by perforating or crushing, or discarding in a safe place away from water supplies.

CAUTION:

KEEP OUT OF REACH OF CHILDREN HARMFUL IF SWALLOWED

68A

PROTECTING THE FOODS THAT FEED THE NATION