


Front Panel

ALCO RONNEL 2 SPRAY

Contains Korlan*

CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.



Controls

FLIES, HORNFLIES, LICE, TICKS, FLEAS, SCREW-WORMS, WOOL MAGGOTS, KEDS, ROACHES, SPIDERS, BEDBUGS, ANTS, MOSQUITOES, SILVERFISH.

USE AROUND FARM, RANCH, HOME, PROCESSING AND PACKING PLANTS, FOOD HANDLING ESTABLISHMENTS, KENNELS, AND MANY OTHER AREAS.

ACTIVE INGREDIENTS	% BY WEIGHT.
Ronnel (0,0-dimethyl 0-(2,4,5-trichlorophenyl) phosphorothioate**)	23.25%
Aromatic Petroleum Derivative Solvents	68.75%
INERT INGREDIENTS	8.00%

Contains 2.0 pounds ronnel per gallon

* Korlan - Registered trademark of the Dow Chemical Co.
U. S. Patent #2,599,516

**From technical grade; not suitable for internal use in animals.

Side Panel

ALCO RONNEL 2 SPRAY KILLS INSECTS RESISTANT TO MANY OTHER INSECTICIDES. (i.e. houseflies, fleas, bedbugs, roaches, screw-worms). It has a long lasting effect.

DIRECTIONS

Mixes readily with water or oil. To prepare a finished spray follow chart below.

TYPE AND USE	LIVESTOCK REMEDY, NOT FOR HUMAN USE				REMARKS
	% Ronnel	Amt. Ronnel 2 Spray to use for			
		1 gal.	3 gal.	25 gal.	
LIVESTOCK INSECTS - Retreat as necessary but not oftener than every 2 weeks					
<u>Cattle and Calves: Hornflies, Lice, and Screw-worms.</u>	1/2%	5 Tbs.	1/2 pt.	2 qt.	
<u>Lone Star and Winter Ticks.</u>	3/4%	7 1/2 Tbs.	3/4 pt.	3 qt.	
<u>Sheep, Goats, Lambs, and Kids: Screw-worms, wool maggots, hornflies</u>	1/2%	5 Tbs.	1/2 pt.	2 qt.	Dilute with water only. Spray entire animal to wet thoroughly.
<u>Lice and sheep keds.</u>	1/4%	2 1/2 Tbs.	1/4 pt.	1 qt.	
<u>Hogs: Lice</u>	1/4%	2 1/2 Tbs.	1/4 pt.	1 qt.	

TYPE AND USE	%	Amt. Ronnel 2 Spray to use for			REMARKS
		1 gal.	3 gal.	25 gal.	
Cattle Backrubber - For Hornflies. Place where animals travel regularly or congregate.					Mix 1 gallon with 27 gallons diesel or similar oil. Saturate fibre. Recharge as necessary. Do not mix with used motor oil.
RESIDUAL FLY SPRAY - a 1/2% spray will give about 2 weeks control. A 1% spray will give 4-6 weeks control.					Spray walls, ceilings, partitions, to obtain complete coverage with no runoff. Use 1 gal. of diluted spray per 500-1000 sq. ft.
Farm Buildings - Including poultry houses, livestock shelters, dairy and hog barns.	1/2% or 1%	5 Tbs. or 1/3 pt.	1/2 pt. or 1 pt.	2 qt. or 1 gal.	Add 3 ounces of sugar per gal. for added effectiveness.
Food Processing Plants	1/2% or 1%	5 Tbs. or 1/3 pt.	1/2 pt. or 1 pt.	2 qt. or 1 gal.	Spray when not in operation. Cover produce and equipment. Spraying outside area will improve control.
Dwellings - Outside areas, including porches, patios, garages, etc.	1/2% or 1%	5 Tbs. or 1/3 pt.	1/2 pt. or 1 pt.	2 qt. or 1 gal.	Spray to point of runoff.
Restaurants, Supermarkets, Etc. - Outside areas.	1/2% or 1%	5 Tbs. or 1/3 pt.	1/2 pt. or 1 pt.	2 qt. or 1 gal.	Spray outside walls, receiving docks, doorways and refuse areas.
Meat Packing Plants	1%	1/3 pt.	1 pt.	1 gal.	Spray outer and inedible product areas.
Poultry Processing Plants	1%	1/3 pt.	1 pt.	1 gal.	(See Meat Packing Plants)
Refuse Areas & Containers	1%	1/3 pt.	1 pt.	1 gal.	Spray to soak all exposed surfaces.
Caged Layers - on wire.	2%	2/3 pt.	1 qt.	2 gal.	Spray poultry droppings beneath cages.
FLY BAIT - Where flies congregate such as floors, windows, doorways, etc.	2%	2/3 pt.	1 qt.	2 gal.	Add 3 oz. of sugar per gallon diluted spray. Spray or sprinkle at 1 to 2 day intervals until control is obtained.
COCKROACH CONTROL - Use 1% spray for German roaches, 2% for others.					Apply as a spot spray to all areas frequented by cockroaches.

ACCEPTED
6/27/68
5481-33

TYPE AND USE	% Konnel	1 gal.	3 gal.	25 gal.	REMARKS
Buildings - Including dwellings, institutions, and food handling establishments.	1% or 2%	1/3 pt. or 2/3 pt.	1 pt. or 1 qt.	1 gal. or 2 gal.	Apply as a spot spray to all areas frequented by cockroaches.
Restaurants, Supermarkets, etc. - Oil mist spray inside buildings for German cockroaches.	1% or 2%	1/3 pt. or 2/3 pt.	1 pt. or 1 qt.	1 gal. or 2 gal.	Dilute with oil and apply by remote control when buildings are not occupied. Cover exposed food and equipment. Use 1 gal. diluted spray per 50,000 cubic feet.
Food Processing Plants	1% or 2%	1/3 pt. or 2/3 pt.	1 pt. or 1 qt.	1 gal. or 2 gal.	Spray when not in operation Cover produce and equipment.
Meat Packing Plants	2%	2/3 pt.	1 qt.	2 gal.	Spray outer and inedible product areas.
Poultry Processing Plants	2%	2/3 pt.	1 qt.	2 gal.	(See Meat Packing Plants)
MISCELLANEOUS INSECTS					Apply 1 gal. of dilute spray per 1000 sq. ft.
Fleas and spiders in buildings, laundry trucks dog kennels, etc.	1%	1/3 pt.	1 pt.	1 gal.	Spray floors, walls, cracks, crevices, sleeping quarters of pets, etc. where insects breed and harbor.
Bedbugs in buildings and sleeping quarters, on beds and bedding.	1%	1/3 pt.	1 pt.	1 gal.	Spray mattress and bedding lightly, including seams and folds, cracks and crevices in bed frames, walls, baseboards, and floors.
Fleas in lawns and kennel runs.	1%	1/3 pt.	1 pt.	1 gal.	Spray infested areas. Keep children off treated areas until spray has dried.
Ants, Mosquitoes and Silverfish	1%	1/3 pt.	1 pt.	1 gal.	Spray areas where insects frequent.

ACCEPTED
6/27/68
2481-33

WARNING: Do not treat swine within 6 weeks of slaughter. Do not apply to swine more frequently than once every 2 weeks. Do not treat cattle within 8 weeks of slaughter. Do not treat sheep and goats within 12 weeks of slaughter. Do not apply more frequently than once every 2 weeks. Do not treat lactating dairy animals, or dry dairy animals within 21 days of the lactating period. Do not use in rooms where milk is processed or stored. Do not allow spray to come in contact with food, feed, or water. Remove lactating dairy animals from buildings during application of residual fly spray. DO NOT USE ON CATS.

CAUTION: INJURIOUS TO THE EYES -- IRRITATING TO THE SKIN -- HARMFUL IF SWALLOWED. Do not get in the eyes. Avoid prolonged or repeated contact with the skin. Avoid prolonged breathing of spray mists. Avoid contamination of food, feedstuffs, and water supplies. Keep material out of reach of children and animals. Keep away from heat and open flame. Keep container closed. Wash thoroughly with soap and water after handling.

FISH AND WILDLIFE CAUTION: To protect fish and wildlife, do not contaminate streams, ponds or lakes.

COMBUSTIBLE LIQUID

DESTROY CONTAINERS.

NEVER RE-USE.

NOTE TO PHYSICIAN: This material is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

U.S.D.A. Reg. #5481 - 33

ACL 113-963

Net Contents _____

Manufactured by

ALCO CHEMICAL COMPANY

Artesia, California

