

COMFILE \AMVAC\5481-58.LAB HD#13  
EPA Reg. No. 5481-58-AA  
Last EPA Acceptance 10/18/84



Page 1 of 4  
These copy changes  
made to accomodate the  
Union Carbide SEVIN®  
Trademark requirements.

**SEVIN®**  
**Brand Carbaryl Insecticide**  
**5% DUST**

**ACCEPTED**  
**AUG 12 1986**  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 5481-58

FOR CONTROL OF CERTAIN INSECT PESTS OF VEGETABLES  
FRUIT, ORNAMENTAL PLANTS, TURF, DOGS AND ADULT CATS

**ACTIVE INGREDIENT:**  
Carbaryl:1-Naphthyl N-methylcarbamate . . . . . 5.0%  
**INERT INGREDIENTS:** . . . . . 95.0%  
Total . 100.0%

SEVIN® is a registered Trademark of Union Carbide Corporation for Carbaryl Insecticides.

KEEP OUT OF REACH OF CHILDREN

**C A U T I O N**

"PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente"

STATEMENTS OF PRACTICAL TREATMENTS

**IF SWALLOWED:** Call a physician of Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

**IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists.

**IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation persists.

**IF INHALED:** Remove victim to fresh air. Apply Artificial Respiration if indicated. Get medical attention.


SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 5481-58-AA

EPA Est. No. 5481-CA-1

Net Weight  1  Pound ( 50 lb.)

**AMVAC CHEMICAL CORPORATION**  
**4100 EAST WASHINGTON BLVD.**  
**LOS ANGELES CA 90023**



**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

**ENVIRONMENTAL HAZARDS**

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the proposed treatment areas.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other Carbaryl labels.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original container in a cool, dry, locked place out of reach of children. **DISPOSAL:** Do not re-use empty container. Rinse thoroughly before discarding in trash.

**READ ENTIRE LABEL BEFORE USING.....USE ONLY AS DIRECTED**

ALCO SEVIN® Brand Carbaryl Insecticide 5% DUST can be applied by sifting from shaker top can, by means of a dust gun or through a squeeze duster. A dust gun will give a more uniform coverage of the plants to be treated.

**DIRECTIONS FOR USE (Cont.)**

**FOR BEST RESULTS MAKE APPLICATION IN THE MORNING OR EVENING WHEN IT IS COOL AND THE AIR IS STILL. BE SURE TO TREAT BOTH UPPER AND LOWER SURFACES OF LEAVES. APPLY TO GIVE THOROUGH COVERAGE BUT AVOID APPLYING IN EXCESS.**

Applications should be started when insects first appear. Repeat applications as necessary. This would normally be at 7 to 10 day intervals, but in rainy weather applications may have to be made more often. Listed below are the crops with the pests controlled by this product followed by the time interval to stop applications prior to harvest. As a general rule apply enough dust to give thorough coverage to all plant parts or 11 to 15 ounces per 1,000 Sq. Ft.



DIRECTIONS FOR USE (Cont.)

PEST CONTROL FOR VEGETABLES & FRUIT

| CROP AND PEST CONTROLLED  | INTERVAL TO STOP USE BEFORE HARVEST                                 |
|---|---|
| <b>BEANS:</b> Mexican Bean Beetle, Leafhoppers, Flea Beetles, Corn Earworms, Lygus Bugs, Stink Bugs, Lima Bean Pod Borer.   | Can be applied day of harvest.                                      |
| <b>CARROTS:</b> Leafhoppers, Flea Beetles.  | Can be applied day of harvest.                                      |
| <b>CORN:</b> Corn Earworm, Flea Beetles, Leafhoppers, Fall Armyworms.<br>For Corn Earworm control treat corn silks every 3 days until silks dry.  | Can be applied day of harvest.                                      |
| <b>POTATOES:</b> (White); Flea Beetles, Leafhoppers.  | Can be applied day of harvest.                                      |
| <b>TOMATOES:</b> Tomato Fruit Worm, Tomato Hornworm, Flea Beetles, Stink Bug, Leafhoppers.  | Can be applied day of harvest.                                      |
| <b>EGGPLANT:</b> Corn Earworm, Flea Beetles, Hornworms, Cucumber Beetles.   | Can be applied day of harvest.                                      |
| <b>PEPPERS:</b> Corn Earworm, Leafhoppers, Flea Beetles.  | Can be applied day of harvest.                                      |
| <b>CUCUMBERS, MELONS, SQUASH, PUMPKINS:</b> Squash Bugs, Leafhoppers, Cucumber Beetles, Flea Beetles.<br>Do not apply to Melons or Cucumbers when temperatures exceed 80° F or when vines are wet.            | Can be applied day of harvest.                                      |
| <b>HEAD LETTUCE:</b> Armyworms, Small Cabbage Looper (treat every 5 to 7 days for control), Leafhoppers, Salt Marsh Caterpillars.   | 3 Days before harvest.  |
| <b>LEAF LETTUCE:</b> Same insects as for Head Lettuce.  | 14 days before harvest.   |
| <b>TABLE BEETS:</b> Armyworms, Flea Beetles, Leafhoppers.   | 3 days. Do not apply with 14 days if tops are used as food or feed. |
| <b>BRUSSELS SPROUTS, BROCCOLI, CABBAGE, CAULIFLOWER:</b> Small Cabbage Looper, (treat every 5 to 7 days for control), Armyworms, Flea Beetles.  | 3 days before harvest.  |
| <b>ASPARAGUS:</b> Asparagus Beetles.  | 1 day before harvest. Do not repeat within 3 days.                  |
| <b>STRAWBERRIES:</b> Leaf Rollers, Lygus Bugs, Salt Marsh Caterpillar.  | 1 day before harvest.   |
| <b>CITRUS:</b> Black Scale, California Red Scale, Soft Brown Scale, Orange Tortrix, California Orange Dog, Fruit Tree Leaf Rollers.<br>For Scale insects best results are obtained when crawlers are present. | 5 days before harvest.  |

## DIRECTIONS FOR USE (Cont.)

**PEST CONTROL FOR ORNAMENTALS****ORNAMENTAL FLOWERS, SHRUBS AND TREES**

Apply this product to thoroughly cover the plant foliage for control of the following pests. Repeat as needed for continued control.

|                 |                         |                     |
|-----------------|-------------------------|---------------------|
| APPLE APHID     | MIMOSA WEBWORM          | BLISTER BEETLES     |
| BOXELDER BUG    | OAK LEAF MINER          | SOFT BROWN SCALE    |
| ELM LEAF BEETLE | ORANGE TORTRIX          | EUROPEAN ELM SCALE  |
| FLEA BEETLE     | PERIODICAL CICADA       | OAK KERMES SCALE    |
| JUNE BEETLE     | PSYLLIDS                | OYSTERSHELL SCALE   |
| LEAFHOPPERS     | ROSE APHIDS             | BLACK SCALE         |
| LEAF ROLLERS    | ROSE SLUG               | COTTONY MAPLE SCALE |
| MEALY BUGS      | THRIPS (Exposed Stages) |                     |

For Scale Insect control best results are obtained when crawlers are present DO NOT APPLY SEVIN® Brand Carbaryl Insecticide to BOSTON IVY AS INJURY WILL RESULT..

**PEST CONTROL FOR TURF**

Apply one pound of this product to cover 250 square feet of turf area to control the following pests:

**EARWIGS - MILLIPEDES - SOD WEBWORMS - LEAFHOPPERS - ANTS**

Turf should be lightly sprinkled after application to wash dust off foliage. Repeat as necessary. Keep children and pets off treated areas until grass has been sprinkled and is dry.

**DOGS & ADULT CATS**

For control of Fleas, and Brown Dog Ticks - Rub into skin and apply in sleeping quarters. Avoid getting into eyes.

**DO NOT TREAT KITTENS UNDER FOUR WEEKS OLD**

**DO NOT USE ON PREGNANT DOGS.**

**DO NOT USE THIS PRODUCT ON DIARY ANIMALS, MEAT ANIMALS OR POULTRY.**

NOTICE: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label, when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the manufacturer and/or seller and the buyer assumes the risk of any such use.

