

ALCO

DEVIN\*

50W

For Control of Certain Insect Pest of Fruit, Vegetables and Ornamentals.

6 Oz. Makes up to 20 Gallons of Spray.

ACTIVE INGREDIENT

% BY WEIGHT

Carbaryl (1-Naphthyl N methyl carbamate)

50%

INERT INGREDIENTS

50%  
100%

\*Registered Trade Mark Union Carbide Corp. for 1-Naphthyl N Methyl Carbamate.  
U. S. Patent #2,903,478

CUTION: Keep Out of Reach of Children.  
See Back Panel for Additional Cautions.

Net Weight \_\_\_\_\_

Side Panel

USE ONLY AS DIRECTED

READ ENTIRE LABEL BEFORE USING.

Alco Devin\* 50W is a wettable powder formulation to be applied as a spray. Keep spray mixture thoroughly agitated. This product can be safely mixed with most fungicides and miticides but do not mix with Bordeaux sprays or other alkaline materials.

DIRECTIONS FOR USE

Ornamentals: For the control of Boxelder bug, Elm leaf beetles, flea beetles, June beetles, leafhoppers, leaf rollers, apple aphid, mealybugs, mimosa webworm, oak leaf miner, orange tortrix, Periodical cicada, Psyllids, Rose aphid, Rose slug, Thrips, (exposed), blister beetles and the following scale insects: Soft Brown, European Elm, Oak Kermes, Oystershell, Cottony Maple, Black. Mix 2 Tablespoonful per gallon of water. Do not apply Alco Devin\* 50W to Boston Ivy as injury will result.

Peaches and Apricots: Peach twig borer, codling moth, orange tortrix, scale, oriental fruit moth, cat facing bugs. Mix 1 Tablespoonful per gallon of water. May be applied within one day of harvest on peaches. 3 days on apricots.

Apples and Pears: Pear psylla, codling moth, fruit tree leaf roller, apple aphids, scale. Mix 2 Tablespoonsful per gallon of water. May be applied 1 day before harvest.

Grapes: Grape berry moth, leafhoppers, June beetles, Grape leaf skeletonizer. Mix 1 Tablespoon per gallon of water. May be applied day of harvest.

Strawberries: Leaf rollers, aphids, salt marsh caterpillar. Mix 2 Tablespoonsful per gallon of water. May be applied one day before harvest.

Beans: Mexican bean beetle, leafhoppers, flea beetles, corn earworms, lygus bugs, lima bean pod bores, stink bugs. Mix 2 Tablespoonsful per gallon of water. Spray to wet all foliage. Repeat application in 7 - 10 days. Can be applied day of harvest. Do not feed vines to dairy or meat animals.

Corn: Corn earworms, flea beetles, leafhoppers, fall armyworm, grasshoppers. For corn earworm control treat silks at 2 to 3 day intervals until silks are dry. Mix 2 Tablespoonsful per gallon of water. Do not feed treated fodder to animals. May be applied day of harvest.

Tomatoes, Peppers, Eggplant: Tomato Fruitworm, Tomato Hornworm, Flea beetles, stink bugs, leafhoppers, corn earworm, cucumber beetle. Mix 2 Tablespoonsful per gallon of water. May be applied day of harvest.

Cucumbers, Melons, Squash, Pumpkins: Squash bug, leafhoppers, flea beetles, cucumber beetles. Mix 2 Tablespoonsful per gallon of water. (Do not apply when temperatures exceed 80° F. or when vines are wet. Do not exceed recommended dosage as plant injury will occur. May be applied day of harvest.

Potatoes (white): Flea beetles, leafhoppers. Apply 2 Tablespoonsful per gallon of water. May be applied day of harvest.

CAUTION: Harmful if inhaled or swallowed. Avoid prolonged or repeated breathing of dust. Avoid unnecessary skin contact. May cause irritation of eyes, nose throat. Wash thoroughly after use. Store out of reach of children or animals. Toxic to honey-bees. Do not use during periods of bee activity. Avoid contamination of Food, Feed, Feeding troughs, water, watering receptacles. Keep out of reach of children and animals. Remove any residues at harvest by peeling, brushing or washing.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Manufactured for  
ALCO CHEMICAL CO.  
Artesia, California