

ALCO

(1,000,00)

DUR-BAN 12

Emulsifiable Insectición

For Use by Pest Control Operators Only to Control Resistant and Non-Resistant Cockroaches.

WARNING: KEEP OUT OF REACH OF CHILDREN.

See side panel for additional warnings and cautions.

ACTIVE INGREDIENTS	BY WI.
0,0-diethyl 8-3,5,6-trichloro-2-pyridyl phosphorothioate*	13.88
Ar natic petroleum derivative solvent (xylene)	44.39
INERT INGREDIENTS	41.73 100.00

* DURSBANR - Trademark of THE DOW CHEMICAL COMPANY

Contains one sound 0.0-liethyl 0-3.5.6-triblor -2-pyridyl phosphorottioate per sallen.

READ ENTIRE LABEL BEFORE USING, USE ONLY AS DIRECTED.

MIXING AND USE DIRECTIONS:

amount of DURSBAN IE in water or in a base oil such as decodorized kerosche. Mill according to the following table to obtain the desired concentration of 0.25% or .56% DURSBAN insecticide. Use the higher concentration (4.56%) to quickly reduce heavy cockroach infestations and the lower concentration (4.25%) on light infestations.

FOR WATER DILUTION	Amount of 1	DURSBAN IF to use
Amount of Spray Mixture Desired	for 0.25/	for 0.507
1 gallon	2 2/2 00.	5 1, 5 02.
3 gallons	ชี Gunces	16 ounces
5 gallons	13.1/3 32.	26 2/3 62.

1/1/10

1411-68

FOR OIL DILUTION			
Amount of Spray	Amount of DURSBAN LE to Use		
Mixture Desired	F · U.25%	For 0.50%	
1 gallon	2 1 8 oz.*	4 1. oz.	
3 gallons	6 3/8 oz.	12 3/4 oz.	
5 gallons	10 5/8 oz.	21 % 02.	

*1 ... = 2 Tablespoons = 6 Teaspoons $1.5z. = 1\frac{1}{5}$ Teaspoons

Use as a spot treatment by applying as a coarse spray or with a paint brush, to creas where cockroaches are normally found. These areas include dark corners of rooms and closets, at the base of walls, in cracks in the wall, behind baseboards, behind and beneath sinks, refrigerators, stoves and cabinets and around draingipes. Apply in a manner to avoid spray or mist contacting food, feedstuff and water applies. Dishes and food handling utensils should be thoroughly waster to soap and water if they have become contaminated by the application of talk predict. Repeat treatment as needed.

NOTE: A period of 4 to 7 days after treatment is normally required for maximum

NOTE: A period of 4 to 7 days after treatment is normally required for maximum affect on cackroaches. DO NOT USE IN EDIBLE PRODUCTS AREAS OF FOOD PROCESSING PLANCE.

Principle crish empty containers and bury them in locations isolated from cannot.

WARNING: KEEP OUT OF REACH OF CHILDREN. May be fatal if swallowed. May be absorbed through the skin. May be irritating to eyes and skin. Do not get in eyes or on skin. Do not wear contaminated clothing. Wash thoroughly after handling. Avoid breathing spray particles. Handle concentrates in ventilated area. Keep away from food, feedstuffs and water supplies. Do not let children or pets contact treated area until thoroughly dry. Flush contaminated eyes and skin with meent of water and for eyes get medical attention.

Note to Physician: Dursban insecticide is a cholinesterase inhibitor. Freat symptomatically. Atropine is antidotal.

Keep container closed and away from heat or pun flame.

Seller warrants that the product conforms to its enemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty or any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE express or implied, extends to the use of this product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

C.s. Patent No. 3,244,586 ACL 269-187

1.5.11.6. Pos. No. 3481-65

Not Contents ______