

ACCEPTED  
11-12-70  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO. 5481-71



# L-T Stock Spray

Contains Lindane and Toxaphene for use on  
Cattle - Hogs - Sheep - Goats

For control of Ticks - Hornflies - Lice - Keds -  
Stable Flies - Mosquitoes

ACTIVE INGREDIENTS	% BY WT.
Toxaphene*	45.0%
Gamma Isomer of Benzene Hexachloride from Lindane	7.0%
Petroleum Distillate	37.0%
Aromatic Petroleum Derivative Solvents	10.0%
<b>INERT INGREDIENTS</b>	<b>6.0%</b>
	<b>100.0%</b>

\*Technical chlorinated camphene (chlorine content, 67% - 69%).

**WARNING: KEEP OUT OF REACH OF CHILDREN**  
**SEE SIDE PANELS FOR ADDITIONAL WARNINGS**

ACL-61563 R-470 NET CONTENTS \_\_\_\_\_  
Manufactured By **ALCO CHEMICAL CO.** Artesia, California

**READ ENTIRE LABEL BEFORE USING.  
USE ONLY AS DIRECTED.**

ALCO L-T STOCK SPRAY is formulated for the control of ticks, hornflies, lice, keds, stableflies and mosquitoes on livestock and also is an effective premise spray for the control of mosquitoes, fleas, stableflies, roaches on such agricultural premises as barns, sheds, animal shelters, fences, other farm buildings. **DO NOT USE IN OR AROUND DAIRY BARN, MILK ROOMS, OR POULTRY HOUSES.**

**DIRECTIONS FOR USE:**

Mix ALCO L-T STOCK SPRAY only as directed under specific usage recommendations. Apply only as specified. Do not treat animals under 3 months of age. Very young animals can be injured by this treatment. Do not apply to dairy cattle or dairy goats. Do not treat meat animals within 30 days of slaughter.

**BEEF CATTLE: TICKS, HORNFLIES, STABLE FLIES** (Supplemental control), **LICE, MOSQUITOES** - Mix one quart in forty gallons of water (5 pints in 100 gallons of water). For control of these pests and protection against reinfestation, spray animals thoroughly, including legs, using two quarts of dilute spray per animal. Repeat as necessary. For lice, one treatment should give season-long control. Do not use on dairy animals.

**SHEEP AND GOATS: LICE AND KEDS** (Sheep Ticks) - Mix one quart in 120 gallons of water (27 fluid ounces per one hundred gallons of water). Spray animals thoroughly to wet, including legs. Better control will be had if the animals are sprayed after shearing. One application will be normally sufficient for control.

**FLEECE WORMS AND TICKS** - Mix one quart in 40 gallons of water (5 pints in 100 gallons of water). Spray animals thoroughly to wet, including legs. Repeat as necessary.

**HOGS: LICE, TICKS** - Mix one quart in 40 gallons of water (5 pints in 100 gallons of water). Apply as a thorough spray. For ticks repeat as necessary. One application is usually sufficient to control lice. **DO NOT SPRAY SOWS WITHIN 2 WEEKS OF FARROWING NOR FOR AT LEAST 3 WEEKS AFTER.**

**BACK RUBBERS - OILERS: HORNFLIES, LICE on BEEF CATTLE** - Mix one quart in 3 gallons fuel oil, diesel oil or other recommended back rubber oil. Charge backrubber tank or soak burlap oiler until thoroughly wet but not dripping. Recharge as necessary. This treatment will give a rapid reduction of hornflies and act as an aid in reducing infestations of lice on cattle. Do not use crankcase oil or other like oils as they will be harmful to the animals. Do not allow meat animals access to backrubbers or oilers within 30 days of slaughter. Do not apply to animals under 6 months of age. Do not apply to dairy animals.

**PREMISE OR AREA SPRAY, AGRICULTURAL PREMISES SUCH AS BARN, SHEDS, FENCES, ANIMAL SHELTERS, OTHER FARM BUILDINGS (EXCEPT DAIRY BARN, MILK ROOMS OR POULTRY HOUSES):** Fleas, mosquitoes, stableflies, roaches. Mix one pint in five gallons of water. Apply to interior and exterior surfaces where insects alight or crawl. For fleas spray cracks, crevices, base of walls and floors. For roaches, spray cracks, crevices, base of walls in barns and areas where seen running. Apply spray to point of runoff. In animal quarters, treat bedding areas as well as ground areas adjacent for improved flea control. Do not treat animals with this dilution. Do not apply in dwellings. Repeat treatment as necessary. Remove or protect all feed and water during spraying operations.

**WARNING:**

Use this product only at recommended rates. Treat as specified only. Do not treat thirsty, overheated or emaciated animals. Do not treat animals when weather is cold or stormy as injury may result from chilling. Do not treat pets with this product. Do not use in any manner not specified on the label. Not for use or storage in or around the home.

**DISPOSAL AND DECONTAMINATION**

Dispose of excess spray material in such a manner so as to not create a hazard, away from any body of water. Clean spills of the concentrate, using a strong caustic and detergent. Flush area with plenty of clear water. Dispose of cleaned up residue in a manner so as to not create a hazard. Wash spray container with plenty of running water. When empty, wash containers thoroughly with detergent and water. Dispose of washings in a safe manner. Puncture and crush metal containers. Break glass containers. Bury or otherwise dispose in a manner so as to not create a hazard. Do not contaminate any body of water.

**THIS PRODUCT IS TOXIC TO FISH AND WILDLIFE.** Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes.

**WARNING:** May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes or clothing. In case of contact of skin, wash immediately with soap and water. For eyes, wash with plenty of clear water. Get medical attention. Wash thoroughly with soap and water. Keep out of reach of children. Do not contaminate water, feed or foodstuffs. Keep away from heat or open flame.

U.S.D.A. Reg. No. 5481-71