

DURHAM MALATHION DUST

COMPOSITION

ACTIVE INGREDIENT:

*Malathion..... 5.0%

INERT INGREDIENTS:..... 95.0%
100.0%

*O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate.

DIRECTIONS FOR USE:

Broccoli, Brussels Sprouts, Cabbage and Cauliflower - For the control of Aphid, apply at the rate of 20 to 35 lbs. per acre. MAKE NO APPLICATION WITHIN 7 DAYS OF HARVEST EXCEPT ON BROCCOLI WHERE NO APPLICATION SHOULD BE MADE WITHIN 3 DAYS OF HARVEST.

Beans, Cucumbers, Melons and Squash - For the control of Aphid, Lygus Bug, Cucumber Beetle, Flea Beetle, Darkling Ground Beetle, Cutworm, Leafhopper and Thrips, apply at the

rate of 25 to 35 lbs. per acre. NO APPLICATION TO BE MADE WITHIN 24 HOURS OF HARVEST.

Eggplant, Grapes, Peas, Peppers and Strawberries - For the control of Aphid and Leafhoppers, apply at the rate of 20 to 25 lbs. per acre. NO APPLICATION TO BE MADE WITHIN 3 DAYS OF HARVEST.

CAUTION: Avoid prolonged breathing of dust. Avoid prolonged or repeated contact with skin, wash thoroughly after using. Avoid contamination of feed and foodstuffs. KEEP OUT OF REACH OF CHILDREN.

2-60

ACCEPTED
FEB 3 - 1966
UNDER THE FEDERAL
FOR THE
ED UNDER
TO ADAP

422-217

5481-81

5



DURHAM CHEMICAL COMPANY
4124 E. PACIFIC WAY
LOS ANGELES 23, CALIFORNIA

NONWARRANTY
of any kind
use of this
use or har
directions