

SEP 24 1987

Mr. H.R. McLane
AMVAC Chemical Corporation
4100 East Washington Boulevard
Los Angeles, CA 90023

Dear Mr. McLane:

Subject: DURHAM-END-OF-TRAIL Snail
EPA Registration No. 5481-95
Your Application Dated May 14, 1987

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Dennis H. Edwards, Jr.
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure

15814:I:Edwards:E-03:KENCO:9/18/87:9/29/87:CB:LF:EK:CB



DURHAM
END OF TRAIL

ACCEPTED
with COMMENTS
in EPA Letter Dated

SNAIL - SLUG AND INSECT GRANULES

SEP 24 1987

A SOIL SURFACE TREATMENT
FOR CONTROL OF

SNAILS, SLUGS, ANTS, EARWIGS, FALL ARMYWORMS,
SOD WEBWORMS, (LAWN MOTHS), AND SOWBUGS

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
amended, for the pesticide
registered under EPA Reg. No.

5481-95

IN

CITRUS GROVES - APPLE & PEACH ORCHARDS - SMALL FRUITS
VEGETABLE GARDENS - ORNAMENTAL GARDENS - GROUND COVERS
- SEED GRASSES -
DICHONDRA & TURFGRASS LAWNS - IN ORNAMENTAL GREENHOUSES

EXCEPT FOR TURF DO NOT APPLY DIRECTLY ONTO PLANTS

For Snails and Slugs

One Pound Treats up to 1,500 Sq.Ft.

For Insect Control Use at the Rate of 1 Pound to Cover 200 Sq.Ft.

*GRANULAR FORMULATION IN CONVENIENT SHAKER APPLICATOR
For Faster More Effective Kill - Blends into Lawns &
Ornamental Garden Areas.

ACTIVE INGREDIENTS:

Metaldehyde	4.0%
Carbaryl (1 naphthyl N-methylcarbamate)	3.0%
INERT INGREDIENTS:	93.0%
Total	100.0%

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

THIS PESTICIDE MAY BE FATAL TO CHILDREN AND DOGS OR OTHER PETS IF EATEN.
KEEP CHILDREN AND PETS OUT OF TREATED AREA.

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.
IF ON SKIN: Remove contaminated clothing and wash effected area with soap and warm water.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 5481-95-AA

EPA EST. NO. 5481-CA-1

NET WEIGHT 50 POUND (*1; 2; 5 LB. Containers)

AMVAC CHEMICAL CORPORATION
4100 EAST WASHINGTON BLVD.
LOS ANGELES CA 90023

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed inhaled or absorbed through the skin. Do not breathe dust, get in eyes, on skin or on clothing. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and warm water before re-use. Keep children and pets out of treated area. Bait should be spread in such a way that it cannot be consumed by children, pets or farm animals.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Birds feeding on treated areas may be killed. Do not apply where run-off is likely to occur. Apply this product only as specified on this label. This pesticide is highly toxic to bees. Protective information may be obtained through your Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconsistent with its labeling. Do not allow bait to contact the edible portions of any food crop.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store product in original labeled container in a cool, dry, locked place out of reach of children. Keep container tightly closed when not in use. Do not store in direct sunlight.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State & local authorities, by burning. If burned stay out of smoke.
HOUSEHOLD: Do not re-use empty container. Wrap and put in trash.

This package is sold by weight when packed, but due to its granular nature the material is inclined to settle.

USE ONLY AS DIRECTED - READ ENTIRE LABEL BEFORE USING

When used as directed DURHAM END-OF-TRAIL SNAIL, SLUG AND INSECT GRANULES will give good control of SNAILS, SLUGS, ANTS, EARWIGS, FALL ARMYWORMS, SOD WEBWORMS (LAWN MOTH), AND SOWBUGS.

Thoroughly water area prior to application. This product can be applied any time during the day but late afternoon or evening is preferred.

*Simply shake on the proper amount of material from the convenient shaker top can. May also be applied from a fertilizer broadcasting unit. Do not water treated area for 48 hours after application.

NOTICE: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label, when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the manufacturer and/or seller and the buyer assumes the risk of any such use.

EPA Reg. No. 5481-95 DURHAM END OF TRAIL
WHERE TO USE AS A SOIL TREATMENT
TREE FRUIT ORCHARDS & GROVES

Includes: Apples, Apricots, Avocado, Cherries, Peaches, and Pears.
Citrus such as - Oranges, Lemons, Limes, Grapefruit, and their Hybrids.

AROUND SMALL FRUIT

Includes: Blackberries, Boysenberries, Loganberries, Raspberries, Grapes, and Strawberries.

APPLY TO SOIL IN VEGETABLE GARDENS

Includes: Artichokes, Asparagus, Beans, Beets, Blackeyed Peas, Broccoli, Brussels Sprouts, Cabbage, Cantaloupe, Carrots, Cauliflower, Celery, Collards, Corn, Cow Peas, Cucumbers, Kale, Kohlrabi, Leeks, Garlic, Melons, Mustard Greens, Okra, Onions, Parsnips, Peas, Peppers, Pimentos, Potatoes, Pumpkins, Radishes, Rhubarb, Rutabagas, Salsify, Shallots, Spinach, Squash, Sweet Potatoes, Swiss Chard, Tomatoes, Turnips, Watermelons,

To control SNAILS and SLUGS, water or irrigate the area to be treated thoroughly. Withhold watering for 48 hours after treatment. Apply this material over the infested area and around the edges of border plants. Also apply to the bases of fences and walls and in shady, protected areas. APPLY AT THE RATE OF ONE POUND TO 1500 SQUARE FEET. Repeat treatment every 3 to 4 weeks during "season" or as needed for control. Do not allow bait to contact the edible portions of any food crop.

To control ANTS, EARWIGS, FALL ARMYWORMS, SOD WEBWORMS(LAWN MOTH) & SOWBUGS APPLY AT THE RATE OF ONE POUND TO 200 SQUARE FEET. Follow watering practices as outlined above. Repeat treatment every 3 to 4 weeks during "season" or as needed for control. Do not allow bait to contact the edible portions of any food crop.

AROUND ORNAMENTALS
OUTDOORS AND IN GREENHOUSES; IN FLOWER GARDENS,
BORDER PLANTINGS, AND AROUND GROUND COVERS.

Azaleas, Begonias, Camellias, Chrysanthemums, Daffodils, Dahlias, Daisies, Geraniums, Honeysuckle, Hydrangeas, Ice Plants, Impatiens, Iris, Ivy, Marigolds, Pansies, Petunias, Rhododendrons, Roses, Snapdragons, Spathaphyllum, Sweet Peas, Tulips, Do not get granules on Boston Ivy.

To control SNAILS and SLUGS, thoroughly water area to be treated. Withhold watering for 48 hours after treatment. Apply this material over the infested area and around the edges of border and ground covers plants. Apply also to the bases of fences and walls and in shady, protected areas. APPLY AT THE RATE OF ONE POUND TO 1500 SQUARE FEET. Repeat treatment every 3 to 4 weeks during "season" or as needed for control.

To control ANTS, EARWIGS, FALL ARMYWORMS, SOD WEBWORMS(LAWN MOTH) & SOWBUGS APPLY AT THE RATE OF ONE POUND TO 200 SQUARE FEET. Follow watering practices as outlined above. Repeat treatment every 3 to 4 weeks during "season" or as needed for control.

FOR DICHONDRA AND TURF LAWNS

To control SNAILS and SLUGS, water area to be treated thoroughly before application. Apply DURHAM END OF TRAIL SNAIL, SLUG AND INSECT GRANULES at the rate of ONE POUND TO 1500 SQUARE FEET.

To control ANTS, EARWIGS, FALL ARMYWORMS, SOD WEBWORMS(LAWN MOTH) & SOWBUGS APPLY AT THE RATE OF ONE POUND TO 200 SQUARE FEET. Follow watering practices as outlined above. Repeat treatment every 3 to 4 weeks during "season" or as needed for control.

FOR SEED GRASSES

Including: Alfalfa, Hay, Wheat, Oats, Barley. For Snails use 29-30 lbs./acre. For Snails, Slugs & Insects use at same rate as for Dichondra & Turf Lawns.