

# DURHAM SEVIN METALDEHYDE GRANULES 5-2.5

ACCEPTED

MAY 30 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 5481-100

## ACTIVE INGREDIENTS:

Carbaryl\* (1-Naphthyl N-Methylcarbamate) ..... 5.0%  
Metaldehyde ..... 2.5%

## INERT INGREDIENTS: .....

92.5%

TOTAL 100.0%

\*Trademark of Union Carbide Corp. for the active ingredient, 1-Naphthyl N-Methylcarbamate.

## WARNING

May be fatal if swallowed.

Do not take internally. Harmful dust. Avoid breathing dust and vapor. Avoid contact with skin, eyes, and clothing. Do not contaminate or store near feed or foodstuffs.

Wash after handling. Baits should be spread in such a way that children, pets, or farm animals cannot reach them.

This product is toxic to fish.

Birds feeding in treated areas may be killed.

Do not store in direct sunlight.

## USE PRECAUTIONS

Use only on crops and for purposes recommended on the label. Toxic to honey bees. Do not apply during bloom when honey bees or other pollinators are active. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe place until one week after application. Do not allow to drift to adjoining crops. The grower is responsible for residues on his crops, as well as for damages caused by drift from his property to that of others. In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops.

## CONTAINER DISPOSAL

Do not reuse empty containers. Destroy it by burying with waste.

## ENVIRONMENTAL HAZARD

This product is toxic to fish, shrimp, crab, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NET WEIGHT: 50 POUNDS

## RECOMMENDATIONS

Durham Sevin Metaldehyde Granules 5-2.5 is recommended for the control of snails, slugs, ants, earwigs, sowbugs, fall armyants, sod webworms, (lawn moths), and cutworms on crops listed.

Water area to be treated well, prior to application. Withhold rewatering treated area for 48 hours. This product can be applied anytime during the day, but late afternoon or evening applications are preferred.

## DIRECTIONS FOR USE

**CITRUS** (Oranges, Lemons and Grapefruit): For the control of slugs and snails apply at the rate of 30 to 50 pounds per acre through any cyclone type hand seeder or mechanical ground applicators that gives even distribution of granules. Broadcast Durham Sevin Metaldehyde Granules 5-2.5 liberally under trees around irrigation pipes, farm buildings and any other place slugs and snails may congregate.

**CAUTION:** Do not apply to tree foliage or edible parts. Durham Sevin Metaldehyde Granules 5-2.5 should never be applied to a dry soil. It is suggested that the application be made following an irrigation.

**ORNAMENTAL GARDENS, PLANTED BORDERS, AND GROUND COVERS:** For the control of snails, slugs, cutworms, armyworms, sowbugs, dark ground beetles, and crickets apply at the rate of 30 to 50 pounds per acre (one pound will cover 1000 to 1500 square feet) through any cyclone type hand or mechanical seeder that gives even distribution of the granules. Water area to be treated well, prior to application. This product can be used any time during the day, but late afternoon or evening application is preferred. Do not water treated area for 48 hours after application.

**DICHONDRA AND TURF LAWNS:** For the control of snails and slugs apply at the rate of 30 to 50 pounds per acre (one pound will cover 1000 to 1500 square feet) through a cyclone type hand or mechanical seeder that gives even distribution of the granules. For the control of ants, earwigs, fall armyworms, sod webworms (lawn moths) and sow bugs apply 4 pounds of this material per 1000 square feet in the same manner as described above. Water area to be treated well, prior to application. This product can be used any time during the day, but late afternoon or evening application is preferred. Do not water treated area for 48 hours after application.

EPA Reg. No. 5481-100-AA

- EPA Est. 5481-CA-1

0475

5481-100  
05/30/1975  
1/1

AMVAC CHEMICAL CORP.  
LOS ANGELES, CALIFORNIA 90023

**WARNING!**  
**Keep Out of Reach of Children**  
This pesticide may be fatal to children, and dogs or other pets if eaten. Keep children and pets out of treated area.

**NOTICE:** Amvac Chemical Corporation makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Amvac's label.