


Reg no: 5481-100

PM-12

181



DURHAM CARBARYL METALDEHYDE GRANULES 5 - 2.5

FOR USE AS A SOIL SURFACE TREATMENT
IN CITRUS - ORNAMENTAL GARDENS
DICHONDRA & TURF LAWNS

For control of Snails, Slugs, Ants, Earwigs,
Sowbugs, Fall Armyworms, Sod Webworms
(Lawn Moth), and Cutworms

ACTIVE INGREDIENTS:

Metaldehyde 2,4,6,8-tetramethyl-1,3,5,7-tetraoxocyclo-octane) — — — — —	2.5%
Carbaryl (1 naphthyl N-methylcarbamate) — — — — —	5.0%
INERT INGREDIENTS: — — — — —	92.5%
Total	100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

THE PESTICIDE MAY BE FATAL TO DOGS OR OTHER PETS IF EATEN
KEEP PETS OUT OF TREATED AREA

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or comatose person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists. **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and warm water.

SEE BACK PANEL FOR NOTE TO PHYSICIAN AND ADDITIONAL PRECAUTIONARY STATEMENTS
EPA REG. NO. 5481-100 EPA EST. NO. 5431 CA-1
NET WEIGHT: 50 POUND (51 lb. Shaker Can; 2 & 5 lb. Containers)

AMVAC CHEMICAL CORPORATION
4100 EAST WASHINGTON BLVD.
LOS ANGELES, CALIFORNIA 9002

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: May be harmful if swallowed inhaled or absorbed through the skin. Do not breathe dust, get in eyes, on skin or on clothing. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and warm water before re-use. Keep pets out of treated area. Bait should be spread in such a way that it cannot be consumed by pets or farm animals.
NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, Opiates or cholinesterase inhibiting drugs.

ENVIRONMENTAL HAZARDS
This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes). Do not contaminate water when disposing of equipment washwaters. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in manner inconsistent with its labeling. Do not allow bait to contact the edible portions of any food crop.

STORAGE AND DISPOSAL
Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store product in original labeled container in a cool, dry, locked place out of reach of children. Keep container tightly closed when not in use. Do not store in direct sunlight.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State & local authorities, by burning. If burned stay out of smoke.
HOUSEHOLD: Do not re-use empty container. Wrap in several layers of newspaper and put in trash.



RECOMMENDATIONS

Durham Carbaryl Metaldehyde Granules 5-2.5 is recommended for the control of Snails, Slugs, Ants, Earwigs, Sowbugs, Fall Armyworms, Sod Webworms (Lawn Moth) and Cutworms at crop sites as listed. Water area to be treated well, prior to application. Withhold rewatering treated area for 48 hours. This product can be applied anytime during the day, but late afternoon or evening applications are preferred.

CITRUS

(Oranges, Lemons, Grapefruit): For the control of Slugs and Snails apply at the rate of 30 to 50 pounds per acre through any cyclone type hand spreader or mechanical ground applicators that give even distribution of granules. Broadcast Durham Carbaryl Metaldehyde Granules 5-2.5 liberally under trees around irrigation pipes, farm buildings and any other place slugs and snails may congregate. **IMPORTANT:** Do not apply to tree foliage or edible parts. This product should never be applied to a dry soil. It is suggested that the application be made following an irrigation.

ORNAMENTAL GARDENS, PLANTED BORDERS, AND GROUND COVERS

For the control of Snails, Slugs, Cutworms, Armyworms, Sowbugs, Darkling Ground Beetles, and Crickets apply at the rate of 30 to 50 lbs. per acre (one pound will cover 1000 to 1500 Sq.Ft.), through any cyclone type hand or mechanical seeder that gives even distribution of the granules. Water area to be treated well, prior to application. This product can be used any time during the day, but late afternoon or evening application is preferred. Do not water treated area for 48 hours after application.

DICHONDRA AND TURF LAWNS

For the control of Snails and Slugs apply at the rate of 30 to 50 lbs. per acre (one pound will cover 1000 to 1500 Sq. Ft.) through a cyclone type hand or mechanical seeder that gives even distribution of the granules. For the control of Ants, Earwigs, Fall Armyworms, Sod Webworms (Lawn Moth), and Sow bugs apply 4 pounds of this material per 1,000 Sq. Ft. in the same manner as described above. Water area to be treated well, prior to application. This product can be used any time during the day, but late afternoon or evening is preferred. Do not water treated area for 48 hrs. after application.

NOTICE: This product conforms to its chemical description and is reasonably effective for the purpose stated on the label, when used in accordance with the directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the manufacturer and/or seller and the buyer assumes the risk of any such use.

07/01/89