

DURHAM SEVIN^{*} DUST 10

ACTIVE INGREDIENTS:
CARBARYL (1-NAPHTHYL N-METHYLCARBAMATE) 10.0%
INERT INGREDIENTS: 90.0%
TOTAL 100.0%

*TRADE-MARK OF UNION CARBIDE FOR THE ACTIVE INGREDIENT, 1-NAPHTHYL N-METHYLCARBAMATE.

CAUTION:

HARMFUL IF SWALLOWED OR INHALED.
DO NOT BREATHE DUST. DO NOT TAKE INTERNALLY.
SKIN CONTACT MAY BE HARMFUL. AVOID CONTACT.
WASH HANDS BEFORE EATING. TAKE SHOWER OR BATH AFTER WORK.
WEAR REGULAR LONG SLEEVED WORK CLOTHING. CHANGE TO CLEAN CLOTHING DAILY.

NOTE FOR PHYSICIANS:

CARBARYL IS A MODERATE, REVERSIBLE, CHOLINESTERASE INHIBITOR. ATROPINE IS ANTIDOTAL.
AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLE.

USE PRECAUTIONS:

USE ONLY FOR CLAIMS LISTED AND IN ACCORDANCE WITH ALL DIRECTIONS.

TOXIC TO HONEY BEES. DO NOT APPLY DURING BLOOM WHEN HONEY BEES OR OTHER POLLINATORS ARE ACTIVE. WHEN NECESSARY TO USE DURING SUCH PERIODS, WARN BEEKEEPERS WELL IN ADVANCE TO LOCATE HIVES AT A SAFE DISTANCE UNTIL ONE WEEK AFTER APPLICATION.

AVOID DRIFT TO ADJOINING FORAGE CROPS.

IN ORDER THAT PESTICIDAL RESIDUES ON FOOD AND FORAGE CROPS WILL NOT EXCEED TOLERANCES ESTABLISHED BY THE FEDERAL FOOD AND DRUG ADMINISTRATION, USE ONLY AT RECOMMENDED RATES AND INTERVALS, AND DO NOT APPLY CLOSER TO HARVEST THAN SPECIFIED. DO NOT APPLY OR ALLOW TO DRIFT TO AREAS OCCUPIED BY UNPROTECTED HUMANS OR BENEFICIAL ANIMALS OR ONTO ADJOINING FOOD, FIBER OR PASTURE CROPS. THE GROWER IS RESPONSIBLE FOR RESIDUES ON HIS CROPS, AS WELL AS FOR DAMAGES CAUSED BY DRIFT FROM HIS PROPERTY TO THAT OF OTHERS.

CONTAINER DISPOSAL:

DO NOT REUSE EMPTY CONTAINER. DESTROY BY BURYING IN A SAFE PLACE.

DIRECTIONS FOR USE:

APPLY THE SPECIFIED AMOUNTS OF THIS PRODUCT UNIFORMLY TO THE FOLIAGE WHEN AIR IS CALM, USING GROUND EQUIPMENT OR AIRCRAFT. APPLY AT THE FIRST SIGN OF INFESTATION AND INFECTION AND REPEAT AT 7 TO 10 DAY INTERVALS AS NEEDED TO MAINTAIN CONTROL.

U.S.D.A. REG. NUMBER 1422-454

NET WEIGHT 50 POUNDS

ACCEPTED
9-15-71
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 5481-108

ACCEPTED WITH COMMENTS

DURHAM CHEMICAL COMPANY

4124 E. PACIFIC WAY

LOS ANGELES, CALIFORNIA 90023

CAUTION!
KEEP OUT OF REACH OF CHILDREN

ACCEPTED

9-15-71

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR THE POISON REGISTERED
ED UNDER NO. 5481-108

RATES ARE GIVEN FOR AVERAGE CONDITIONS AND EXPRESSED AS POUNDS OF PRODUCT PER ACRE APPLICATION. FOR UNUSUAL CONDITIONS MAXIMUM ALLOWABLE RATES PER ACRE ARE GIVEN IN PARENTHESES (), AFTER EACH CLAIM.

RECOMMENDATIONS:

(READ ALL PRECAUTIONS AND DIRECTIONS FOR USE BEFORE USING.)

BEANS -- USE 10 LB. FOR MEXICAN BEAN BEETLE; USE 10 LB. FOR BEAN LEAF BEETLE, CUCUMBER BEETLES, FLEA BEETLES, JAPANESE BEETLE, LEAFHOPPERS, VELVETREAN CATERPILLAR AND WESTERN BEAN CUTWORM. USE 10-15 LB. FOR ARMYWORMS AND TARNISHED PLANT BUGS; 20 LB. FOR CORN EARWORM, LIMA BEAN POD BORER, LYGUS AND STINK BUGS. MAY BE APPLIED UP TO AND INCLUDING THE DAY OF HARVEST. (21 LB.)

COTTON -- USE 10 LB. FOR EARLY SEASON CONTROL OF: COTTON FLEAHOPPER, COTTON LEAFWORM, FLEA BEETLES, STRIPED BLISTER BEETLE, THRIPS. USE 10-20 LB. FOR MID AND LATE-SEASON CONTROL OF: BOLL WEEVIL, BOLLWORM FALL ARMYWORM, COTTON LEAF PERFORATOR, LEAFROLLERS, LEAFHOPPERS, LYGUS BUGS (LIGHT TO MODERATE INFESTATIONS IN WESTERN IRRIGATED COTTON). TARNISHED PLANT BUGS, APHIDS (POPULATIONS WILL BE SUPPRESSED BY REPEATED APPLICATIONS OF THIS INSECTICIDE). USE 15-25 LB. FOR PINK BOLLWORM. (APPLY ON A 5 - 7 DAY SCHEDULE DEPENDING ON THE SIZE OF THE COTTON AND DEGREE OF INFESTATION.) USE 20 LB. FOR SALT-MARSH CATERPILLAR, STINK BUGS. USE 10-15 LB. FOR GRASSHOPPERS. THE HIGHER DOSAGE SHOULD BE USED WHEN GRASSHOPPERS ARE MATURE OR MATERIAL IS APPLIED ON PARTLY DEFOLIATED PLANTS. ALLOW 7 DAYS BETWEEN LAST APPLICATION AND GRAZING TREATED FIELDS OR HARVESTING CROP RESIDUE FOR USE AS FEED OR BEDDING FOR DAIRY OR MEAT ANIMALS.

EGGPLANT, PEPPER, POTATO AND TOMATO:

USE 10 LB. FOR COLORADO POTATO BEETLE, FLEA BEETLES, AND LEAFHOPPERS. USE 10-20 LB. FOR EUROPEAN CORN BORER, FALL ARMYWORM, LACE BUGS, (EGGPLANT AND PEPPER), TOMATO FRUITWORM, TOMATO HORNWORM, TARNISHED PLANT BUGS AND STINK BUGS. MAY BE APPLIED UP TO AND INCLUDING DAY OF HARVEST. (POTATO --- 20 LB.; OTHER CROPS --- 40 LB.)

GRAPES: -- USE 10-20 LB. FOR LEAF FOLDER, GRAPE LEAFHOPPER, AND GRAPE LEAF SKELETONIZER. FOR GRAPE LEAFROLLER, APPLY 10-15 LB. JUST BEFORE FIRST BROOD LARVAE EMERGE FROM THE FIRST ROLLS. APPLY 10-20 LB. FOR THE CONTROL OF 2ND AND 3RD BROODS. APPLY AS NEEDED FOR LEAFHOPPERS. USE 20 LB. FOR GRAPE BERRY MOTH, JAPANESE BEETLES, JUNE BEETLES, AND RED-BANDED LEAFROLLER. MAY BE APPLIED UP TO AND INCLUDING THE DAY OF HARVEST (90 LB.)

PEACHES: ORIENTAL FRUIT MOTH, AND PEACH TWIG BORER; APPLY 50 LB. PER ACRE IN MID-SUMMER AND LATE WINTER. MAKE 4 APPLICATIONS AT 10-60 DAY INTERVALS. MAY BE APPLIED UP TO ONE DAY OF HARVEST. (80 LB.)

STRAWBERRIES: USE 10-20 LB. FOR MEADOW SPITTLEBUG, STRAWBERRY LEAF ROLLER AND STRAWBERRY WEEVIL. MAY BE APPLIED UP TO ONE DAY OF HARVEST. (20 LB.)

LJB

9-2-71

128

ACCEPTED WITH COMMENTS

NOTICE: Durham Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Durham's label.