

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER - POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. RAPIDLY ABSORBED THROUGH THE SKIN. REPEATED INHALATION OR SKIN CONTACT MAY, WITHOUT SYMPTOMS, PROGRESSIVELY INCREASE SUSCEPTIBILITY TO MEVINPHOS INSECTICIDE POISONING** Do not swallow or get in eyes, on skin or on clothing. Do not breathe vapors. Do not contaminate food or feed products. Keep away from heat or open flame.

**PRECAUTIONS IN USING**

Wear clean natural rubber gloves and waterproof protective clothing, and goggles or face shield. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type jointly approved by the Mine Safety and Health Administration and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part 11 approved for Mevinphos protection.

Wear clean clothes daily. Wash thoroughly with soap and water after handling and before eating or smoking. Pilot should not assist in mixing and loading operation. Do not contaminate or store near food or feedstuff. Note Reentry Statements section.

**IMPORTANT:** Not for use or storage in or around the home. Keep animals and unprotected persons out of operational areas and while there is a danger of drift. In case of spillage, decontaminate areas and equipment by washing with a dilute solution of alkali less than 5% and detergent and rinse with water.

**NOTE TO PHYSICIAN - POISONING SYMPTOMS**

Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

**TREATMENT:** Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams, until complete atropinization has occurred.

Pralidoxime chloride (2-PAM chloride) may also be used as an effective antidote in addition to and while maintaining full atropinization. In adults, an initial dose of 1 gram of 2-PAM should be injected, preferably as an infusion in 250cc of saline over a 15 to 20 minute period. If this is not practical, 2-PAM may be administered slowly by intravenous injection as a 5 percent solution in water over not less than two minutes. After about an hour, a second dose of 1 gram of 2-PAM will be indicated if muscle weakness has not been relieved. For infants and children the dose of 2-PAM is 0.25 grams.

Morphine is an improper treatment.

Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been attained as determined by blood test.

For 24 hours emergency medical assistance call (213) 264-3910.

**ENVIRONMENTAL HAZARDS**

This product is toxic to birds, fish, shrimp, crab, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on this label. Not for use or storage in or around the home.

Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Unprotected humans or animals should be kept out of treated fields on the day of treatment. See Reentry Statement. Do not apply when weather conditions favor drift from areas treated.

**PHYSICAL & CHEMICAL HAZARDS**

**FLAMMABLE** Keep away from heat or open flame. Liquid phases separate at lower temperatures. Warm or store at higher temperatures and mix thoroughly to recombine liquid phases and assure uniformity before use.

**RESTRICTED USE PESTICIDE**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

**DURHAM DURAPHOS EM 4**  
**ORGANOPHOSPHORUS INSECTICIDE**

Contains 4 lbs. MEVINPHOS per Gallon

ACTIVE INGREDIENTS:

2-Carbomethoxy-1-methylpyridinemethyl phosphate, alpha isomer. . . . . 28.2%

Related Compounds. . . . . 18.8%

INERT INGREDIENTS: \* . . . . . 53.0%

Total 100.0%

Equivalent to 50%/w MEVINPHOS Insecticide

**KEEP OUT OF REACH OF CHILDREN**

**DANGER - POISON**  
**PELIGRO**



"PRECAUCION AL USARLO. Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente"

**STATEMENTS OF PRACTICAL TREATMENT**

**IF SWALLOWED** - Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Never induce vomiting or give anything by mouth to an unconscious or convulsing person.

**IF INHALED** - Remove victim to fresh air immediately. Apply artificial respiration if indicated. Get medical attention.

**IF IN EYES** - Flush eyes with plenty of water for at least 10 minutes. Get medical attention immediately.

**IF ON SKIN** - Remove contaminated clothing immediately and wash effected area with soap and water. Call a physician immediately. Wash clothing before re-use.

**IF POISONING SYMPTOMS** appear (See Poisoning Symptoms under NOTE TO PHYSICIAN) keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest hospital.

SEE ANTIDOTE AND OTHER PRECAUTIONS ON SIDE PANEL

EPA Reg. No. 5481-113-AA

EPA Est. No. 5481-CA-1

**NET CONTENTS 30 GALLONS (2½ Gallons)**

MADE IN U.S.A.

**AMVAC CHEMICAL CORPORATION**

Do  
at  
tre  
re  
yo  
W  
in  
of  
the  
wa  
Or  
un  
D/  
ap  
ST.

Do  
Of  
dr  
fre  
(80  
of  
Pl  
dis  
CA  
PE  
cid  
of  
Co  
gu  
rec  
ap  
rec  
if c

RE

Applic  
approv  
Pour  
Keep  
tends  
wettab  
plants  
table  
SUGGE  
provide  
for cert

# RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

# DURHAM DURAPHOS EM 4

## ORGANOPHOSPHORUS INSECTICIDE

Contains 4 lbs. MEVINPHOS per Gallon

### ACTIVE INGREDIENTS:

2-Carbomethoxy-1-methylvinyl dimethyl phosphate, alpha isomer ..... 28.2%

Related Compounds ..... 18.8%

INERT INGREDIENTS: \* ..... 53.0%

Total 100.0%

Equivalent to 50%/w MEVINPHOS Insecticide

KEEP OUT OF REACH OF CHILDREN

**DANGER - POISON**  
**PELIGRO**



"PRECAUCION AL USARLO Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente"

### STATEMENTS OF PRACTICAL TREATMENT

**IF SWALLOWED** — Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Never induce vomiting or give anything by mouth to an unconscious or convulsing person.

**IF INHALED** — Remove victim to fresh air immediately. Apply artificial respiration if indicated. Get medical attention.

**IF IN EYES** — Flush eyes with plenty of water for at least 10 minutes. Get medical attention immediately.

**IF ON SKIN** — Remove contaminated clothing immediately and wash affected area with soap and water. Call a physician immediately. Wash clothing before re-use.

**IF POISONING SYMPTOMS** appear (See Poisoning Symptoms under NOTE TO PHYSICIAN) keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest hospital.

SEE ANTIDOTE AND OTHER PRECAUTIONS ON SIDE PANEL

EPA Reg. No. 5481-113-AA

EPA Est. No. 5481-CA-1

NET CONTENTS 30 GALLONS (2½ Gallons)

MADE IN U.S.A.

# AMVAC CHEMICAL CORPORATION

4100 E. WASHINGTON BLVD., LOS ANGELES, CALIF. 90023

# BEST AVAILABLE COPY

## DIRECTIONS FOR USE

293

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**FOR END USE ONLY. DO NOT REPACKAGE OR REFORMULATE WITHOUT MANUFACTURER'S WRITTEN APPROVAL**

### RE-ENTRY AND WORKER SAFETY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in treated areas or in an area to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure.) When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

**DANGER:** Area treated with MEVINPHOS on (Date)..... Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure see STATEMENTS OF PRACTICAL TREATMENT found on the MEVINPHOS product label.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Open dumping is prohibited. **STORAGE:** Do not store in or around the home. Store in a secure, dry, and temperate area. Store in original container only. Product is flammable. Keep away from heat, sparks, or open flame. If spilled, avoid exposure. If significant spill, call CHEMTREC, (800) 424-9300 or (213) 264-3910. Spill can be contained by covering with a sufficient amount of absorbent material, such as clay, diatomaceous earth, sand, sawdust, or sweeping compound. Place in drum, label contents, and dispose of according to the instructions below in pesticide disposal.

**CAUTION:** Wear a pesticide respirator and avoid forming dust clouds during clean-up. **PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your nearest State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance. **CONTAINER DISPOSAL:** (Metal) Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary land fill, or by other procedures approved by State and local authorities. (Plastic) Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### READ THE DIRECTIONS CAREFULLY AND FOLLOW THEM AT ALL TIMES USE AND APPLICATION DIRECTIONS

Application can be made by aircraft or ground power equipment by trained personnel only using approved protective equipment. Do not apply with hand equipment. Pour specified amount of this product into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine with wettable powders unless previous use of the mixture has proven physically compatible and safe to plants. Always thoroughly emulsify this product with at least half of total water before adding wettable powder.

**SUGGESTED WATER RATES FOR AIRCRAFT AND GROUND APPLICATION** — The actual rate required to provide thorough, uniform coverage varies with plant growth at time of application. Except as specified for certain uses, the following rates are therefore intended to cover a broad range of conditions:

Crop	Gal. Water/Acre	
	Aircraft	Ground
Vegetable and Field Crops .....	5-20	20-125
Grapes .....	7-25	20-200
Orchard Crops .....	10-20	40-800

### FRUIT AND

**APPLES, PEACHES** and mites use control of grasshopper use ½ t roller use ½ t treat within 1 t per acre on peaches and peaches be permitted to activity require for 4 days fol or more organ entry interval est applicable terval.

**CHERRIES (SOU)** use ¼ to ¾ t in 2 days of h

**CITRUS: ORANGE** aphids use 1 water. For con and omnivorous water. For con cutworm, varie pillar use 2 q least 7 days 1 day of harv enter treated. substantial co lowing treatm ganophosphate val shall be e able interval t

**GRAPES:** For c Do not treat leaf folder, le lygus bugs us do not treat not be permit any activity r foliage for 4 of 2 or mor the re-entry in longest applic interval.

**STRAWBERRIES** use ¼ pt. per pers. strawbe lygus bugs us treat within 1 and mites, us treat within 1 per acre

**WALNUTS:** For 100 gals. wat orange tortrix moth use ½ in 1 day of h

### FIELD C

**ALFALFA, CLO** caterpillar us grasshoppers, alfalfa weevil acre. Do not

**MUSTARD BR** use ¼ to 1 looper, impor terous leafm mites use ½ 3 days of h

a manner inconsistent with its labeling.

PACKAGE OR REFORMULATE

WRITTEN APPROVAL

SAFETY STATEMENT

Directly or through drift expose workers or others not protected by unprotective persons. Do not enter treated areas until the spray has dried. Because certain states may have laws which prohibit the use of crops treated with this product, consult your state agricultural extension agent.

Do not enter without proper protective clothing. In case of accidental exposure see MEVINPHOS product label.

DISPOSAL

Proper disposal. Do not reuse empty container. Do not use in or around the home. Store in a secure, fire-resistant container. Product is flammable. Keep away from heat and fire. If significant spill, call CHEMTREC. Clean up by covering with a sufficient amount of soil, sand, sawdust, or sweeping compound. Follow the instructions below in pesticide disposal.

FOLLOW THEM AT ALL TIMES

DIRECTIONS

Apply by trained personnel only using appropriate equipment. Add balance of water to fill tank. Do not mix with other pesticides unless proven physically compatible and safe to use. Do not combine with other pesticides unless proven physically compatible and safe to use.

APPLICATION - The actual rate required to control pests varies with time of application. Except as specified below, apply to cover a broad range of conditions:

Table with 2 columns: Gal. Water/Acre and Application Rate. Rows for Aircraft and Ground, with sub-rows for 5-20, 7-25, and 10-20 gal. water per acre.

FRUIT AND NUT CROPS

APPLES, PEACHES, PEARs, PLUMs: For control of aphids and mites use 1/4 to 1/2 pt. per 100 gals. water. For control of grasshoppers, lygus bugs and red-banded leafhopper use 1/2 to 3/4 pt. per 100 gals. of water. Do not treat within 1 day of harvest or apply more than 5 pts. per acre on peaches and plums. 6 1/2 pts. per acre on apples and pears. PEACHES ONLY: Workers should not be permitted to enter treated areas to engage in any activity requiring substantial contact with treated foliage for 4 days following treatment.

CHERRIES (SOFT ONLY): For control of aphids and mites use 1/2 to 3/4 pt. per 100 gals. water. Do not treat within 2 days of harvest or apply more than 5 pts. per acre.

CITRUS, ORANGES, LEMONS, GRAPEFRUIT: For control of aphids use 1 to 2 pt. per acre in 200 or more gals. water. For control of fruit tree leafroller, orange tortrix and omnivorous leafroller use 1 qt. per acre in 500 gals. water. For control of Western tussock moth larvae, citrus cutworm, variegated cutworm and pink scavenger caterpillar use 2 qt. per acre in 1200 gals. water. Allow at least 3 days between applications. Do not treat within 1 day of harvest. Workers should not be permitted to enter treated areas to engage in any activity requiring substantial contact with treated foliage for 4 days following treatment.

CUCUMBERS: For control of aphids use 1/4 to 1 pt. per acre. Do not treat within 2 days of harvest. For control of cucumber beetle, leafhoppers, mites, red-banded leafhopper and lygus bugs use 1 quart per acre. At the 2 qt. dosage do not treat within 5 days of harvest. Workers should not be permitted to enter treated areas to engage in any activity requiring substantial contact with treated foliage for 4 days following treatment.

STRAWBERRIES: For control of aphids and mites use 1/2 to 3/4 pt. per 100 gals. water. For control of grasshoppers, strawberry leafroller, salt-marsh caterpillar and lygus bugs use 1/2 to 1 pt. per 100 gals. water. Do not treat within 1 day of harvest. For hard to kill aphids and mites, use 1 qt. per 100 gals. water, but do not treat within 1 day of harvest or apply more than 1 qt. per acre.

WALNUTS: For control of aphids use 1/4 to 1/2 pt. per 100 gals. water. For control of mites, omnivorous looper, orange tortrix, fruit tree leafroller, and Western tussock moth use 1/2 pt. per 100 gals. water. Do not treat within 1 day of harvest or apply more than 5 pts. per acre.

FIELD CROPS

ALFALFA, CLOVER: For control of aphids and alfalfa caterpillar use 1/4 to 1/2 pt. per acre. For control of grasshoppers, leafhoppers, cutworms (climbing), alfalfa weevil larvae, lygus bugs use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

MUSTARD GREENS, TURNIP TOPS: For control of aphids use 1/4 to 1/2 pt. per acre. For control of cabbage looper, imported cabbage worm, false chinch bug, dipterous leafminer (adult), grasshoppers, leafhoppers and mites use 1/2 to 1 pt. per acre. Do not treat within 3 days of harvest.

SORGHUM (FOR FORAGE AND GRAIN): For control of aphids use 1/4 to 1/2 pt. per acre. For control of corn earworm and webworm use 1/2 to 1 pt. per acre. For control of fall armyworms use 1 pt. per acre. Do not treat within 3 days of harvest.

VEGETABLE CROPS

ARTICHOKES: For control of aphids use 1/4 to 1/2 pt. per acre. For control of plume moth use 1 to 2 pts. per acre. Do not treat within 2 days of harvest.

BEANS: For control of aphids use 1/4 to 1/2 pt. per acre. For control of grasshoppers, leafhoppers, mites and Mexican bean beetle use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

BEETS (INCLUDING TOPS): For control of aphids use 1/4 to 1/2 pt. per acre. For control of cabbage looper, cutworm (climbing), dipterous leafminer (adult), false chinch bug, grasshoppers, imported cabbage worm, leafhopper, mites and salt-marsh caterpillar use 1/2 to 1 pt. per acre. Do not treat within 3 days of harvest.

BROCCOLI, CABBAGE: For control of aphids use 1/4 to 1/2 pt. per acre. For control of cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, salt-marsh caterpillar, mites, cutworms (climbing), dipterous leafminer (adult) and lygus bugs use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest. For hard-to-kill aphids, use 1 qt. per acre, but do not treat within 3 days of harvest.

BRUSSELS SPROUTS, CAULIFLOWER, COLLARDS, KALE: For control of aphids use 1/4 to 1/2 pt. per acre. For control of cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, salt-marsh caterpillar, mites, cutworms (climbing), dipterous leafminer (adult) and lygus bugs use 1/2 to 1 pt. per acre. Do not treat within 3 days of harvest. For hard to kill aphids, use 1 qt. per acre, but do not treat collards and kale within 7 days of harvest; 3 days on Brussels sprouts and cauliflower.

CARROTS: For control of aphids use 1/4 to 1/2 pt. per acre. For control of leafhoppers, lygus bugs, mites, cabbage looper, dipterous leafminer (adult), cutworm (climbing) and salt-marsh caterpillar use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

CELERY: For control of aphids use 1/4 to 1/2 pt. per acre. For control of dipterous leafminer (adult), lygus bugs, salt-marsh caterpillar, leafhoppers, cabbage loopers and mites use 1/2 to 1 pt. per acre. Do not treat within 3 days of harvest. For hard to kill aphids, use 1 qt. per acre, but do not treat within 5 days of harvest.

CORN (FIELD, SWEET, AND POPCORN) FOR GRAIN ONLY: For control of aphids use 1/4 to 1 pt. per acre. Do not treat within 1 day of harvest.

CUCUMBERS: For control of aphids use 1/4 to 1/2 pt. per acre. For control of grasshoppers, leafhoppers and mites use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

EGGPLANT, PEPPERS: For control of aphids use 1/4 to 1/2 pt. per acre. For control of grasshoppers, leafhoppers and mites use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

LETTUCE: For control of aphids use 1/4 to 1/2 pt. per acre. For control of corn earworm, cutworms (climbing), dipterous leafminer (adult), cabbage looper, imported cabbage worm, grasshoppers, mites, lygus bugs, salt-marsh caterpillar, false chinch bug and thrips use 1/2 to 1 pt. per acre. Do not treat within 2 days of harvest. For hard to kill aphids, use 1 qt. per acre, but do not treat within 4 days of harvest.

MELONS (INCLUDING CANTALOUPE, HONEY-DEW MELONS, MUSKMELONS, WATERMELONS): For control of aphids use 1/4 to 1/2 pt. per acre. For control of cabbage looper, dipterous leafminer (adult), leafhoppers, lygus bugs, mites, false chinch bugs, salt-marsh caterpillars, cutworms (climbing) and grasshoppers use 1/2 to 1 pt. per acre. To control ringworms (cabbage looper, cutworms, salt-marsh caterpillars, tobacco budworms) on watermelons apply 1/2 pt. per acre. Do not treat cantaloupe, honey-dew melons, muskmelons or watermelons within 1 day of harvest.

ONIONS (INCLUDING GREEN ONIONS): For control of thrips and cutworms (climbing) use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

PEAS (INCLUDING VINES): For control of aphids use 1/4 to 1/2 pt. per acre. For control of grasshoppers, leafhoppers, mites and cutworms (climbing) use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

POTATOES: For control of aphids use 1/4 to 1/2 pt. per acre. For control of grasshoppers, leafhoppers and mites use 1/2 pt. per acre. Do not treat within 1 day of harvest.

SPINACH: For control of aphids use 1/4 to 1/2 pt. per acre. For control of cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, mites, dipterous leafminer (adults), cutworms (climbing), salt-marsh caterpillar and false chinch bug use 1/2 to 1 pt. per acre. Do not treat within 4 days of harvest. For hard-to-kill aphids use 1 qt. per acre, but do not treat within 7 days of harvest.

SUMMER SQUASH: For control of aphids use 1/4 to 1/2 pt. per acre. For control of cabbage looper, dipterous leafminer (adults), leafhoppers, lygus bugs, mites, false chinch bugs, salt-marsh caterpillar, cutworms (climbing) and grasshoppers use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

TOMATOES: For control of aphids use 1/4 to 1/2 pt. per acre. For control of grasshoppers, leafhopper and mites use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

TURNIPS: For control of aphids use 1/4 to 1/2 pt. per acre. For control of cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, mites, false chinch bug and dipterous leafminer (adults) use 1/2 to 1 pt. per acre. Do not treat within 3 days of harvest.

NOTICE: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label, when used in accordance with directions under normal conditions or use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the manufacturer and/or seller, and buyer assumes the risk of any such use.

MEVINPHOS MIXTURE LIQUID-POISON B

RO MEVINPHOS NA 2783

ACCELED

1-87

MAR 30 1987 544-113

BEST AVAILABLE COPY