

5481-130

04/12/1988

1/1

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. In case of eye contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse.

RE-ENTRY AND WORKER SAFETY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in treated areas or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written and oral warnings must include the following information. **CAUTION:** Areas treated with Fruit Fix SUPER CONCENTRATE 800 on (Date) _____. Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, refer to the Precautionary Statements found on this product label. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwater. Avoid drift of spray mist.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconsistent with its labeling. Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original container in a cool, (not in direct sunlight), dry, locked place, out of reach of children. Do not allow product to freeze. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

Fruit Fix SUPER CONCENTRATE 800 serves to control preharvest drop of apples & pears, leading to larger harvest & reducing losses from windfall and knockdown. Fruit Fix SUPER CONCENTRATE 800 is also effective in thinning the crop of olive trees when an excessively heavy fruit set is expected. Fruit Fix SUPER CONCENTRATE 800 may be used to prevent fruiting of ornamental olive trees. In addition, Fruit Fix SUPER CONCENTRATE 800 may be used for fruit thinning of ORANGES, TANGERINES (Mandarin), TANGELOS AND TANGORS.

DIRECTIONS FOR USE ON APPLES AND PEARS

Spray leaves thoroughly, wetting all fruit stems and adjacent leaves. Apply 7 to 14 days before harvest for apples and 5 to 7 days before harvest for pears. Do not spray within 5 days of harvest. This product becomes effective in 2 to 3 days after application and controls drop for 10 days to 4 weeks (see for McIntosh variety) depending on weather conditions. Do not make more than 2 applications and do not delay picking beyond optimum maturity.

GROUND SPRAY - FOR APPLES, thoroughly mix 3 fl.oz. with 250 gallons of water (1.2 fl. oz./100 gals. water) to prepare a 20 ppm spray. **FOR PEARS,** use a 10 ppm spray obtained by mixing 3 fl. oz. with 500 gals. water. Drench the trees with the appropriate spray, using up to 670 gallons per acre of this 10 ppm spray, which provides and equivalent of 25 grams of 1-Naphthaleneacetic acid per acre. Apply to fruit 5 to 7 days before harvest. Do not make more than 2 applications. Do not apply within 2 days of harvest. (For an "oil-type" spray, add 1 pint of flowable summer oil for each 100 gallons of spray solution).

Fruit Fix SUPER CONCENTRATE 800

A HORMONE TYPE PRODUCT FOR CONTROL OF
PREHARVEST DROP OF APPLES AND PEARS

ACTIVE INGREDIENT:

1-Naphthalene Acetic Acid, Ammonium Salt . . . 21.40%

INERT INGREDIENTS:

78.60%

Total 100.00%

Equivalent to 19.6% 1-Naphthaleneacetic Acid or 800 grams (1.78 lbs./gallon).

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

EPA Reg. No. 5481-130-AA

EPA Est. No. 5481-CA-1

AMVAC CHEMICAL CORPORATION

4100 EAST WASHINGTON BLVD
LOS ANGELES, CALIFORNIA 90023

NET CONTENTS 1 GALLON

AIRPLANE SPRAY - APPLES & PEARS - Mix 5-1/2 fl. oz. in 3 gallons of water and emulsify with 2 gallons of flowable summer oil by thorough agitation. Apply 5 gallons of emulsion per acre of APPLES. For PEARS, use 2-3/4 fl. oz. of this product in 5 gallons of emulsion per acre.

DIRECTIONS FOR THINNING OLIVE TREES

TIMING IS CRITICAL: Spray when young fruits are 1/8" to 3/16" in diameter, which is from 1-1/2 to 2-1/2 weeks following full bloom. Thinning is effective on all varieties except Savillano.

WITHOUT OIL - Apply a solution of 150 ppm 1-Naphthaleneacetic acid equivalent obtained by adding 9 fl. oz. and a wetting agent (e.g. "Dreit" 14 to 16 oz.) to 100 gallons of water.

WITH OIL - Apply a solution of 100 ppm 1-Naphthaleneacetic acid equivalent obtained by mixing thoroughly 6 fl. oz. with 100 gallons of water. Emulsify by agitation with 1-1/2 gallon of light or medium summer oil. Spray only the heavily loaded trees until the leaves are thoroughly wet. This will require 10 to 30 gallons per tree.

PRECAUTION - Unless you wish to remove all your fruit, do not apply during bloom.

DIRECTIONS TO PREVENT FRUITING OF ORNAMENTAL OLIVE TREES

For best result, apply when the olive trees are in full bloom but before fruit sets. When extended bloom occurs, spray between 1/2 to 3/4 bloom, with a second spraying 7 to 14 days later. Mix 6 fl.oz. Fruit Fix SUPER CONCENTRATE 800 and a wetting agent (e.g. 12 oz. liquid detergent) to 80 gallons of water. Spray trees until leaves are thoroughly wet. **PRECAUTIONS:** Applications should not be made when temperatures are over 85 degrees F as leaf curl may occur. Adjacent plants should be covered with polyethylene sheet while spraying and the cover left on until danger of dripping and spray drift has passed, to prevent possible injury to these plants.

DIRECTIONS FOR FRUIT THINNING ORANGES, TANGERINES (Mandarin), TANGELOS AND TANGORS

Spray at the rate of 100 to 800 ppm (6 fl.oz. per 100 gals. water equals 100 ppm). The degree of thinning obtained with Fruit Fix SUPER CONCENTRATE 800 is highly variable and temperature related. Severe over-thinning may result from application to trees during hot and dry conditions. Do not apply to "non-thrifty" trees. The grower must determine a suitable concentration and time of application for his cultivars and location. Concentrations in the 100 to 500 ppm range should be tried and applications made when the fruit are in an early stage of development (5 to 20 mm in diameter) and near the normal period of "June Drop" (in Florida the period of natural drop occurs in May.) Higher thinning activity should be expected during hot weather. Therefore lower concentrations should be used during hot weather (85 degrees F and higher) and higher concentrations should be used during cooler and wet weather (85 degrees to 95 degrees) The effectiveness of Fruit Fix SUPER CONCENTRATE 800 diminishes rapidly below 85 deg rees. The entire fruiting area of the tree should receive a thorough coverage of the spray solution.

DO NOT APPLY MORE THAN 1 APPLICATION PER YEAR
DO NOT USE ON VARIETIES WHERE BOTH YOUNG FRUITS AND FRUITS CLOSE TO MATURITY ARE PRESENT ON TREE.
DO NOT APPLY WITHIN 150 DAYS OF HARVEST.

NOTICE: This product conforms to its chemical description & is reasonably fit for the purpose stated on the label, when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the manufacturer and/or seller and the buyer assumes the risk of any such use.

LEPAVAMVAC/5481-129-AA 03-07-88

ACCEPTED
APR 12 1988
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 5481-130