



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON D C 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

Ms Kelly K Naujock Amvac Chemical Corporation 4695 MacArthur Court, Suite 1250 Newport Beach, CA 92660

AUG 31 2012

Subject

Label Amendment

Product Name Alco Citrus Fix

EPA Reg No 5481-145

Application dated March 21, 2012

Dear Ms Naujock

The proposed label amendment of the product referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable

Please note that marketing claims made on the pesticide label must be substantiated by data maintained in your files. If data supporting marketing claims made on the product label is not available, then those claims must be removed

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E) 40 CFR 156 10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance

Products shipped after 18 months from the date on this notice or the next printing of the label whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA. Your release for shipment of this product constitutes acceptance of these conditions. This label supersedes all previously accepted labels. You must submit one copy of the final printed label.

<u>before the product is released for shipment</u> One copy of the label stamped "Accepted" is enclosed for your records. If you have any questions, please contact Tracy White by phone at (703) 308-0042 or via email at white tracy@epa.gov

Sincerely,

Kathryn V Montague

Product Manager 23 Herbicide Branch

Registration Division (7505P)

Enclosure-Stamped Label and Stamped Supplemental Label



CITRUS FIX®

Plant Growth Regulator

wie de Act,

AUG 3 1 2012

ACTIVE INGREDIENT

Isopropyl Ester of 2 4 Dichlorophenoxyacetic Acid*

45 0%

INERT INGREDIENTS

55<u>0%</u>

TOTAL

100 0%

*2 4 Dichlorophenoxyacetic Acid Equivalent 38% Contains petroleum distillates

3 36 lbs 2 4 D acid equivalent per gallon

(, t,

The State of California Department of Agriculture classifies this ester of 2,4-D as "Highly Volatile" and therefore does not permit its use as a herbicide in California

KEEP OUT OF REACH OF CHILDREN **CAUTION**

	FIRST AID		
If swallowed	Call a poison control center or doctor immediately for treatment ad	vice	
	Do not give any liquid to the person		
Do not induce vomiting unless told to do so by the poison control center or doct			
	Do not give anything by mouth to an unconscious person		
If on skin or	Take off contaminated clothing		
clothing	Rinse skin immediately with plenty of water for 15 20 minutes		
	Call a poison control center or doctor for treatment advice	_	
If inhaled	Move person to fresh air		
	If person is not breathing call 911 or an ambulance then give artificial respiration preferably by		
	mouth to mouth if possible		
	Call a poison control center or doctor for further treatment advice		
If in eyes	Hold eye open and rinse slowly and gently with water for 15 20 mi	inutes	
	Remove contact lenses if present, after the first 5 minutes then con	ntinue rinsing eye	
	Call a poison control center or doctor for treatment advice		
	EMERGENCY INFORMATION		
Have the product treatment	container or label with you when calling a poison control center or	doctor, or going for	
FOR THE FOLLO	NING EMERGENCIES PHONE 24 HOURS A DAY		
For Medical Emergencies Phone		1 888 681-4261	
For Transportation Emergencies including spill leak or fire phone CHEMTREC		1 800 424 9300	
		1 888 462 6822	
	NOTE TO PHYSICIAN	- AND THE STREET	
Camtaina anamatia			

Contains aromatic solvents May pose an aspiration pneumonia hazard Contains petroleum distillates Vomiting may cause aspiration pneumonia

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE

EPA REG NO 5481 145 EPA EST NO 5481 CA 1 Net Contents As Marked on Container



145 20120627 combo label

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION Causes moderate eye irritation Harmful if swallowed absorbed through the skin or inhaled Avoid contact with skin eyes or clothing Avoid breathing vapor or spray mist

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are made of barrier laminate or viton
If you want more options follow the instructions for Category (G) on an EPA chemical resistance category selection chart

All mixers loaders applicators and other handlers must wear

Long sleeved shirt and long pants

Shoes and socks

Chemical resistant gloves

Chemical resistant apron when applying postharvest dips or sprays to citrus mixing or loading cleaning up spills or equipment or otherwise exposed to the concentrate

Follow manufacturer s instructions for cleaning/maintaining PPE

If no such instructions for washables exist use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

Aerial application is prohibited

User Safety Recommendations

User should

- Wash hands before eating drinking chewing gum using tobacco or using the toilet
- Remove clothing/PPE immediately if pesticide gets inside Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product Wash the outside of gloves before removing. As soon as
 possible wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

PHYSICAL OR CHEMICAL HAZARDS

Combustible Do not use, pour spill or store near heat or open flame

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training decontamination notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is

- Coveralls
- Chemical resistant gloves made of any waterproof material
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms forests nurseries or greenhouses. Do not enter or allow others to enter the treated area until sprays have dried

USE PRECAUTIONS

- ALCO CITRUS FIX is a plant growth regulator for use on citrus only Do not use as a herbicide!
- Do not use on citrus trees less than 6 years old
- Do not apply during a flush of leaf growth
- Do not apply within 7 days of harvest
- Do not allow drift to susceptible plants which include but are not limited to cotton grapes roses beans peas alfalfa lettuce ornamentals and broadleaf plants. This product may injure cotton beans peas grapes ornamentals etc (coarse sprays are less likely to drift)
- Do not use equipment that has been used to spray ALCO CITRUS FIX to spray 2 4 D sensitive plants Always use caution in disposing of spray solutions as they can cause extreme injury or kill sensitive plants (See Storage and Disposal section)
- Before using spray equipment for any other purpose thoroughly clean same with hot soap suds followed by soaking and washing with ammonia or baking soda

Spray Drift Management

A variety of factors including weather conditions (e.g. wind directions wind speed temperature relative humidity) and method of application (e.g. ground airblast chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product

Droplet Size

When applying sprays that contain 2 4 D as the sole active ingredient or when applying sprays that contain 2 4 D mixed with active ingredients that require a coarse or coarser spray apply only as a coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles

When applying sprays that contain 2 4 D mixed with other active ingredients that require a medium or more fine spray apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on target deposition and there are not sensitive areas (including but not limited to residential areas bodies of water known habitat for nontarget species nontarget crops) within 250 feet downwind. If applying a medium spray leave one swath unsprayed at the downwind edge of the treated field

Temperature Inversions

If applying at wind speeds less than 3 mph the applicator must determine if a) conditions of temperature inversion exist or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food forage or other plantings that might be damaged or crops thereof rendered unfit for sale use or consumption Susceptible crops include but are not limited to cotton okra flowers grapes (in growing stage) fruit trees (foliage) soybeans (vegetative stage) ornamentals sunflowers tomatoes beans and other vegetables or tobacco Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2 4 D herbicides Where states have more stringent regulations they must be observed

Equipment

All application equipment must be properly maintained and calibrated using appropriate carriers or surrogates

2 4 D esters may volatilize during conditions of low humidity and high temperatures. Do not apply during conditions of low humidity and high temperatures

For ground boom application

Do not apply with a nozzle height greater than 4 feet above the crop canopy

Increase fruit size on growing Navel oranges, Valencia oranges and grapefruit Limited to one application per crop cycle Maximum of 45 grams ae per acre (0 1 lbs ae per acre)

To reduce pre-harvest fruit drop on growing Navel oranges, Valencia oranges and grapefruit Limited to one application per crop cycle Maximum rate of 200 ppm per application

To prevent pre-harvest drop of mature fruit and leaves on lemons, Navel oranges, Valencia oranges and Tangelos Limited to one application per copycle Maximum rate of 24 ppm per application

Postharvest packing house application to lemons Limited to one application per crop Maximum rate of 500 ppm per application

USE DIRECTIONS

The following Use Directions as well as the above Use Precautions are taken from data developed and published by the University of California Experiment Station, Riverside California For additional Uses and Precautions of ALCO CITRUS FIX in California consult the UC IPM Pest Management Guidelines CITRUS Plant Growth Regulator Section or visit www.ipm.ucdavis.edu/pdf/pmg/pmgcitrus.pdf

MIXING DIRECTIONS

Carefully measure and accurately add the required quantity of ALCO CITRUS FIX to the spray mixture Add only during the agitation of the spray material

PRE HARVEST FRUIT DROP CONTROL
Add the specified quantity of ALCO CITRUS FIX to spray tank and spray to give adequate coverage
acre for mature trees
DO NOT USE ON CITRUS TREES YOUNGER THAN SIX YEARS DO NOT APPLY WITHIN 7 DAYS OF HARVEST

TRE-DARVEST FRO	UII DROF CONTROL S	gram acıd equivalent/a	OF 500 GALLONS PER ACRE OR LESS Application Rates of ALCO CITRUS FIX Per Acre		
Variety/Species	Time	(gae/a)	ml/acre	oz /acre	
Navel Orange ¹	Sept Nov *	30	75	2 5	
_	Sept Nov (in hydrated lime)	45	112	3 75	
	Dec Jan **	15	37 5	1 25	
Valencia Orange ¹	Young fruit at least ½ dia	15	37 5	1 25	
Grapefruit ¹ (Summer Drop)	Young fruit at least ¾ dia	30	75	2 5	
Grapefruit ¹ (Winter Drop)	Sept Jan **	30	75	2 5	
(···	Sept Jan ** (in hydrated lime)	45	112 5	3 8	
Lemon and Limes***	Oct Dec	15	56 25	19	
Tangelo, Minneola and other Citrus Hybrids, Tangerines (mandarins, mandarin oranges and Satuma mandarins)	Sept Nov	30	75	2 5	

¹ Limited to one application for pre harvest fruit drop control per crop cycle

PRE-HARVEST FRUIT DROP CONTROL SPRAY SOLUTIONS ABOVE 500 GALLONS PER ACRE							
			Amount of ALCO CITRUS FIX Per 100 Gallons				
Variety/Species	Time	ppm	ml	Maximum rate per acre (ml)****	fl oz	Maximum rate per acre (fl oz)****	
Navel Orange ¹	Sept Nov * Sept Nov (in hydrated lime) Dec Jan **	16 24 8	15 2 22 8 7 6	(122) (182) (61)	0 51 0 77 0 26	(4 1) (6 2) (2 1)	
Valencia Orange ¹	Young fruit at least ½ dia	8	7 6	(61)	0 26	(2 1)	
Grapefruit ¹ (Summer Drop)	Young fruit at least ¾ dia	16	15 2	(122)	0 51	(4 1)	
Grapefruit ¹ (Winter Drop)	Sept Jan **	16	15 2	(122)	0 51	(4 1)	
• •	Sept Jan ** (in hydrated lime)	24	22 8	(182)	0 77	(6 2)	
Lemon and Limes*** ¹	Oct Dec	8	76	(61)	0 26	(2 1)	
Tangelo, Minneola and other Citrus Hybrids, Tangerines (mandarins,	Sept Nov	16	15 2	(122)	0 51	(4 1)	

mandarın oranges and Satuma mandarıns)¹

- * Sept /Oct May be too early to provide good control when conditions favor fruit drop (i.e. warm winter protracted harvest) Consult your local ALCO CITRUS FIX dealer local farm advisor or AMVAC representative
- ** January sprays may be somewhat risky especially when environmental factors favor and earlier than usual Spring flush of growth Consult your local ALCO CITRUS FIX dealer local farm advisor or AMVAC representative
- *** Applications to coastal lemons are risky because of their everbearing nature Before use, consult with your local ALCO CITRUS FIX dealer local farm advisor or AMVAC representative
- **** Use the maximum amount of ALCO CITRUS FIX per acre regardless of water volume used
- Limited to one application for per harvest drop control per crop cycle

PREVENTION OF LEAF AND FRUIT DROP CAUSED BY A PESTICIDE OIL SPRAY

For use in pesticide oil sprays to counteract leaf drop and fruit drop caused by the oil Add 2 2 ml (0 07 fl oz) to 3 8 ml (0 12 fl oz) per gallon of oil Apply thorough coverage

The addition of ALCO CITRUS FIX to the pesticide oil spray provides some degree of preharvest fruit drop control But because of the long time span between most pesticide oil sprays and the onset of mature fruit drop the ALCO CITRUS FIX + oil spray mixture may not give adequate mature fruit drop control especially if the per acre ALCO CITRUS FIX dosage is less than 24 grams acid equivalent (gae/acre) or 60 ml/acre (2 02 fl oz/acre) ALCO CITRUS FIX

Under these conditions an aqueous ALCO CITRUS FIX preharvest spray may be needed If needed follow the guidelines in the chart below

ADDITIONAL PREHARVEST ALCO CITRUS FIX SPRAY REQUIREMENTS (in addition to ALCO CITRUS FIX + oil spray)				
Time from ALCO CITRUS FIX + oil spray application to mature fruit drop period	Additional preharvest ALCO CITRUS FIX application*			
more than 6 months	Apply the full rate for preharvest fruit drop control			
4 to 6 months	Apply the 50% rate for preharvest fruit drop control			
3 to 4 months	If per acre dosage (in oil spray) was less than 60 ml/acre acre (2 02 fl oz/acre) apply the 25% rate for preharvest fruit drop control			
3 to 4 months	If per acre dosage (in oil spray) was at least 60 ml/acre acre (2 02 fl oz/acre) NO additional product should be applied			

^{*} Apply to trees 6 years and older DO NOT apply within 7 days of harvest

INCREASING FRUIT SIZE

Fruit sizing sprays require excellent coverage. Use a minimum of 500 gallons per acre for mature trees. Apply water spray as an outside coverage using the recommended milliliters or fluid ounces of ALCO CITRUS FIX shown in Table 2. Increase fruit size on growing Navel oranges, Valencia oranges and grapefruit. Limited to one application per crop cycle. Maximum of 45 grams at per acre (0.1 lbs at per acre).

To reduce pre harvest fruit drop on growing Navel oranges, Valencia oranges and grapefruit Limited to one application per crop cycle Maximum rate of 200 ppm per application

To prevent pre harvest drop of mature fruit and leaves on lemons, Navel oranges, Valencia oranges and Tangelos Limited to one application per crop cycle Maximum rate of 24 ppm per application

Postharvest packing house application to lemons Limited to one application per crop Maximum rate of 500 ppm per application

TABLE 2					
Variety	Fruit Size at Time of Spraying (Inches in Diameter)	Milliliters Per Acre	Fluid Ounces Per Acre	Gram Acid Equivalent (GAE)	
	3/16 to 1/4	56	19	23	
Navel &	1/4 to 1/2	75	2 5	30	
Valencia Oranges	1/2 to 5/8	94	3 2	38	
	5/8 to 3/4	112 5	3 8	45	
	1/4 to 3/8	56	19	23	
Conv. C. A	3/8 to 5/8	75	2 5	30	
Grapefruit	5/8 to 3/4	94	3 2	38	
	3/4 to 1	112 5	3 8	45	

DIRECTIONS FOR PACKING HOUSE USE

- 1 To keep buttons alive and healthy and therefore prevent Alternaria rot
- 2 In lemon packing houses where a water wax emulsion is applied to the fruit prior to storage the growth regulator should be added to this wax emulsion
- If a water wax emulsion is not used then a separate treatment must be arranged to follow the final fresh water rinse of the fruit. Such a treatment could be designed so as to flush a growth regulator solution over the fruit and re circulate the surplus or apply a very fine spray of growth regulator to the fruit and not re circulate the small surplus. The amount of growth regulator adhering to the fruit would be about the same in either case, so no change in concentration would be necessary.
- 4 A 500 ppm concentration of 2 4 D is required. This is obtained by adding 1 5/8 fluid ounces (48 ml) of ALCO CITRUS FIX to each 10 gallons of water.

ADDITIONAL PRECAUTIONS

- Under certain circumstances which seem to be related to the use of chromates in the airwasher 2 4 D is not compatible with nitrogen trichloride. Decco as used for control of decay. Packing houses using Decco should avoid using a growth regulator in the wax emulsion or use it with extreme caution. Rind injury appearing as reddish or brownish discoloration has been greatly intensified by these growth regulators in some packing houses using Decco.
- When a minimum storage period is desired it would be a disadvantage to use 2.4 D because of the delay in changing to a vellow color. Also the percentages of juice may not increase as rapidly
- 3 Late summer use of 2 4 D in the packing house may necessitate more careful culling for weak fruit (blossom end decline) because of the retention of green buttons in such fruit

Use of this product in certain portions of California Oregon and Washington is subject to the January 22 2004 Order for injunctive relief in Washington Toxics Coalition, et al v EP, C01 01320 (W D WA) For further information please refer to http://www.epa.gov/espp/wtc

IMPORTANT Any equipment used in spraying any product containing 2,4 D should not be used to spray 2,4 D sensitive plants. In disposing of solutions of 2,4 D, it should be borne in mind that they can cause extreme injury or death to sensitive plants.

STORAGE AND DISPOSAL Do not contaminate water food or feed by storage or disposal

PESTICIDE STORAGE This product may crystallize if stored at cold temperatures If crystallization occurs call AMVAC for instructions before using Store product in original container in a secure dry and temperate area. Keep container closed

when not in use Do not store near food feed fertilizers seeds insecticides or fungicides. Do not store 30 gallon drums over two high. Do not use or store around the home. Avoid contact with water. In case of spill or leak soak up with sand earth or synthetic absorbent (do not use alkaline absorbents) and dispose of wastes in compliance with local State and Federal regulations.

PESTICIDE DISPOSAL Pesticide wastes are toxic Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA regional office for guidance. Do not contaminate water food or feed

CONTAINER DISPOSAL

Triple rinse as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container ¼ full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration or if allowed by State and local authorities by burning. If burned stay out of smoke Nonrefillable container. Do not reuse or refill this container.

ALCO CITRUS FIX may be used as a Plant Growth Regulator on Citrus in Florida Refer to CITRUS FIX Florida labels EPA SLN No FL 890040 and EPA SLN No FL 980006 for complete directions

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label (b) that this product is reasonably fit for the purposes set forth in the directions for use subject to the inherent risks referred to herein when it is used in accordance with such directions and (c) that the directions warnings and other statements on this label are based upon responsible experts evaluations of reasonable tests of effectiveness of toxicity to laboratory animals and to plants and residues on food crops and upon reports of field experience. Tests have not been made on all varieties of food crops and plants or in all states or under all conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THERE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED IN NO EVENT, SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT

AMVAC offers this product and Buyer accepts it subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC

ALCO[®] is a registered trademark of AMVAC Chemical Corporation CITRUS FIXTM is a trademark of AMVAC Chemical Corporation

AMVAC Chemical Corporation 4100 E Washington Boulevard Los Angeles CA 90023 U S A 1 888 462 6822 www amvac chemical com

Page 9 of 9
145 20120627 combo label