

ACCEPTED

FEB 9 1979

This material is for use only in the United States and is not recommended for the pest control of insects under 5481-165

NALED 8 INSECTICIDE

Active Ingredients:

*Naled

By Wt.

58.0%

Aromatic Petroleum Derivative Solvent

27.5%

Inert Ingredients

14.5%

*1,2-dibromo-2,2-dichloroethyl dimethyl phosphate.

Contains 8 pounds technical naled per gallon equivalent to 7.2 pounds naled.

READ THE LABEL

DANGER: Keep out of reach of children.
See back panel for additional cautions.

NET CONTENTS: ONE GALLON

OTHER ORNAMENTAL PLANTS TOMATOES CUCUMBERS Spider Mites Adult Whiteflies Aphids Leafhoppers
Apply undiluted on heat pipes for open pens on hot plates. Apply continuously or to intermittent sections of
perimeter of each range depending upon width. Application to cold pipes should be followed by immediately heating
at least 1 hour after pipes are hot. With hot water systems, close house for at least 3 hours. House may remain closed
if hot pipes wear goggles and approved respirator. Ventilate house before workers reenter. Wear respirator when
cucumbers and other houseplants are covered with plastic or rubber sheet during treatment and until house is vented.
Do not harvest. Spider Mites (Resistant and Non-Resistant) Make 3 to 4 treatments at 3 or 4 day intervals. Repeat
treatments as necessary. Aphids Apply when aphids appear. Repeat as necessary. WARNING: Avoid
contact with White Butterfly and Golden Rapture roses. Pink Champagne Chrysanthemums. Green Wandering Jew
and Mimosa.

Feeding Plants: Leafing Docks, Cull Piles and Refuse Areas, Cider Mills - Use 2 oz. in 2 1/2 gal. water (15 pts. to 100 pts. l.
water) and cull piles where insects congregate. Apply every 5 to 7 days as necessary. Do not apply to cull fruit or refuse
in food processing machinery. Do not apply when plants are in operation or when foods are present or
in contact with food. Cover food containers during spraying.

WATER AND HOUSEFLY CONTROL - RESIDENTIAL AREAS CROPS LISTED ON LABEL, TIDAL MARSHES
AND PASTURES INCLUDING DAIRY CATTLE Consult your State Fish and Game Agency before
applying to buildings and dairy barns. Make applications during peak of infestation and repeat as necessary. See crop
recommendations for use on shrimp. Do not apply to tidal or marsh waters which are important shrimp producing areas.
With water. Apply 2 to 8 qt. of diluted spray per acre. Repeat. Apply 1.5 to 4 fl. oz. per acre diluted with water.
COLD FOG Use 3 to 5 qt. per 100 gal. water. Calibrate equipment (rate of flow and output) to apply 0.1 to 0.25 lb.

1. Dairy and Beef Cattle Hogs Sheep Horses Hens Chickens - Airplane Application Use 1.5 to 4 fl. oz. per
100 sq. ft. of spray per acre. Apply over areas with animal present.

CRICKETS Outside Dwellings and in Lawns Use 1 fl. oz. in 3 gal. water. Apply to turf and to soil surfaces around

(in label) and Pastures including dairy pastures. Range Caterpillars on Rangeland and Pastures Use 1.5 to 4 fl. oz. per acre by
air or ground. Armyworms (except root) 1 pt. per acre by air or ground. By Air Use in 1 to 5 gal. water per acre
by air. Animals may be present during treatment. See crop recommendations for use in pastures near harvest. Forage
may be soiled or spoiled interstate with 14 days of application.

Do not get on skin, eyes or clothing. Causes eye damage. Use waterproof gloves and face shield
if swallowed. Avoid breathing spray mist. In case of contact, immediately remove contaminated
water, for eyes. Get medical attention. NOTE TO PHYSICIANS: Atropine is antidote. 2 PAM is
antidote with atropine. Avoid contamination of feed, foodstuffs and drinking water. DO NOT USE
IF SAME. Keep children and animals off treated areas. Do not spray on Nectarines. Ornamental
jury may occur. Do not store diluted spray. Make new dilution for each use. Wash equipment

Treated areas may be killed. Keep out of reach of game or ponds. Shrimp and crab may be killed at application rates
of important resources. Do not apply when weather conditions favor drift of spray from areas treated. Rinse equipment
and wash hands from water supplies.

Prevent or reduce on crops. Protective information may be obtained from your Cooperative Agricultural Extension
office or local health department.

THOROUGHLY AND DESTROY. NEVER RE-USE

Warranties that this material is suitable for the chemical and that it is the sole and exclusive liability for use as directed
by the representative to make any other warranty of FITNESS or MERCHANTABILITY. Guarantee

Guarantee from the manufacturer is voided by the use of chemicals. Such risks include but are not
limited to: loss of crop, loss of life and damage caused by drift to other plants or crops. Such risks occur even
when a label through label drift is used. Buyer and user acknowledge and assume all risks and liability
for handling, storage and use of this material.

CHEMICAL CORPORATION, Los Angeles, Calif. 90023

is a registered trademark of Chevron Chemical Company

0179