

WETTABLE SULFUR

AGRICULTURAL INSECTICIDE-FUNGICIDE

WETTABLE SULFUR — FOR MIXING WITH WATER

ACTIVE INGREDIENT:

Sulfur 97.0%

INERT INGREDIENTS:

..... 3.0%

100.0%

CAUTION

Keep Out of Reach of Children

USE PRECAUTIONS

Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops. Do not use on apricots, Anjou and comice pears, cranberries, cucurbits (cucumbers, squash and some varieties of melons), filberts, spinach, tung trees, walnuts and other sensitive plants.

Do not apply when shade temperatures exceed or are likely to exceed 90°F.

Sulfur may burn foliage and fruit during periods of high temperature and under certain climatic conditions.

Do not use sulfur within 30 days of an oil spray (60 days on citrus).

If the crop to be treated will be used for canning or freezing purposes, consult food processor before applying sulfur.

DIRECTIONS FOR USE

Pour recommended amount of this material through screen on surface of water in nearly filled spray tank, or pre-mix in a bucket half filled with water and pour mixture into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, and spraying operations. Do not allow mixture to stand. Do not use in low-volume gear-type spray equipment. Do not combine with emulsifiable liquids in the same spray tank.

EPA Reg. No. 5481-167-AA

EPA Est. 476-CA-5


0774

Manufactured for:

AMVAC CHEMICAL CORPORATION

LOS ANGELES, CA 90023

50 LBS. NET



RECOMMENDATIONS

COTTON: Atlantic mite, cotton rust—By air: Apply 3-10 lbs. per 5 to 10 gallons (45/50 lbs. per acre). By ground equipment: Apply 3-10 lbs. per 40-50 gallons water (45/50 lbs. per acre).

APPLES: Scab, powdery mildew—Apply 4-5 lbs. per 100 gals. from pink or pre-blossom spray through second cover spray. Add 1 gal. lime-sulfur in pink and calyx sprays. Use on sulfur-tolerant varieties only.

CHERRY: Blossom blight brown rot, leaf spot, powdery mildew—Apply 4-6 lbs. per 100 gals. starting at shuck-full.

CITRUS: Six-spotted and citrus rust mites—Apply 4 lbs. per 100 gals. (80 lbs. per acre); and Yuma spider mites—Apply 4 lbs. per 100 gals. (100 lbs. per acre).

GRAPES: Powdery mildew (eradicator spray)—Apply 2 lbs. per 100 gals. plus a wetting agent. Care should be taken to wet all surfaces from which the fungus is to be eradicated.

PEAR: Scab, powdery mildew—Apply 4 lbs. plus 1 gal. lime-sulfur per 100 gals. in the pink or pre-blossom spray. On Bartlett pears, apply 4-6 lbs. at petal fall through second cover sprays.

PEACH, NECTARINE: Brown rot, coryneum blight, rust, peach silver mite, powdery mildew—Apply 5-6 lbs. per 100 gals. from popcorn stage up to harvest. For brown rot control in summer sprays apply up to 10 lbs. per 100 gals. starting about 4-5 weeks before harvest and continuing at weekly intervals.

PLUM, PRUNE: Brown rot, leaf spot, prune rust—Apply 5-6 lbs. per 100 gals. in summer sprays, starting about 5 weeks before harvest.

STRAWBERRIES: Powdery mildew—Apply 1-5 lbs. per 100 gals. Discontinue applications during hot weather.

ORNAMENTALS

ROSES, PERENNIALS, SHRUBS: Black spot, powdery mildew, rust—Apply 3 table-spoons per gallon (6 lbs. per 100 gals.) and spray for full coverage wetting both sides of leaves. Begin applications when new growth appears in the spring and repeat every 7 days, if necessary.

CAUTION: Some burning may occur when used during warm weather.

NOTICE: AMVAC CHEMICAL CORPORATION makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on AMVAC's label.