

INDUSTRIAL

INSTITUTIONAL • INDUSTRIAL • COMMERCIAL • AUTOMOTIVE • INSTITUTIONAL • INDUSTRIAL • COMM

INDUSTRIAL

INSTITUTIONAL • INDUSTRIAL • COMMERCIAL • AUTOMOTIVE • INSTITUTIONAL • INDUSTRIAL • COMM

Dymon

Patent  
Pending

# DRY FOG<sup>®</sup> ULD

## Swat Pyrethrin Insecticide

FOR BAKERIES, FOOD PLANTS,  
RESTAURANTS, SCHOOLS, PLANTS,  
MOTELS, HOTELS & OFFICE BUILDING S.

## ACTIVE INGREDIENTS:

Pyrethrins	0.20%
Piperonyl butoxide, technical	0.72%
N-octyl bicycloheptene dicarboximide	0.40%
Petroleum distillate	8.68%
<b>INERT INGREDIENTS:</b>	<b>90.00%</b>
	<b>Total 100.00%</b>

\*Equivalent to 576% (butylcarbityl) (6-propylpiperonyl) ether and  
144% other related compounds  
•MGK-264 Insecticide Synergist

Net Wt. 8 oz. 227 gms.

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**CONTENTS UNDER PRESSURE**  
See side panel for additional precautions.

For Commercial/Industrial Use Only

EPA Reg. No. 11894-70 EPA Est. No. 11894-KS-1

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

## SHAKE BEFORE USE

**To Kill Flies, Mosquitoes, Gnats, Wasps, Hornets, Sandflies, Small Flying Moths:** Close all doors and windows and direct spray upward into center of room with a slow sweeping motion. Spray 5-10 seconds for average room (1,000 cu ft). Keep nozzle three feet from surfaces. Vacate premises. Keep room closed for 15 minutes after spraying. Ventilate room thoroughly before re-entry. Sweep up and destroy the fallen insects and repeat treatment as necessary.

**To Kill Wasps and Hornets:** Applications should be made in late evening when insects are at rest. Spray into hiding and breeding places contacting as many insects as possible.

**To Kill Roaches, Waterbugs, Silverfish, Ants, Scorpions, Centipedes, Sowbugs, Spiders, Fleas:** Spray baseboards, cracks, crevices, moist areas and openings around sinks and pipes and other hiding places, hitting the insects directly. Spray ant hills and ant trails around windows and door frames and other locations. For fleas, thoroughly spray animal quarters and sleeping area. Repeat as necessary.

**To Kill Clothes Moths:** Spray all articles to be protected, giving particular attention to folds and seams. Spray interior of storage containers, such as closets, chest and trunks, regularly. Treatment should be repeated every thirty days, unless the articles are stored in a moth-tight container.

**To Kill Bedbugs:** Spray mattress thoroughly especially around seams and tufts. Spray bedsteads, baseboards and floorboards. Repeat frequently.

## STORAGE AND DISPOSAL

**Storage:** Contents under pressure. Do not store near fire, sparks or flame. Do not puncture. Exposure to temperatures above 120° F may cause bursting.

**Pesticide Disposal:** Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Do not re-use the empty container. Do not puncture or incinerate. Wrap and discard in trash.

**General:** Consult federal, state or local disposal authorities for approved alternative procedures.


## PRECAUTIONARY STATEMENTS: Hazards to Humans and Domestic Animals

## CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in plants, bakeries and other food processing plants, equipment and utensils must be washed with an effective cleaning compound and thoroughly rinsed with potable water before operations are resumed. All food processing surfaces and utensils should be covered during treatment, or removed from area before use. Remove pets, birds and cover fish aquariums before spraying.

MANUFACTURED BY  
**Dymon, Inc.** 3401 KANSAS AVENUE  
KANSAS CITY, KANSAS 64106

DA-179



ACCEPTED

AUG 22 1990

Under the Federal Insecticide, Fungicide, and  
Rodenticide Act, as amended.

5481-212

75% Reduction

ALCO

## PCNB 2-E LIQUID EMULSIFIABLE CONCENTRATE SOIL FUNGICIDE

FOR CONTROL OF SOIL BORNE DISEASES  
AS LISTED  
ON COTTON - BEANS - GARLIC - POTATOES  
TURFGRASS and ORNAMENTALS

ACTIVE INGREDIENT:  
Pentachloronitrobenzene(PCNB) - 24.0%  
INERT INGREDIENTS: - - - - - 76.0%  
TOTAL 100.0%  
Contains 2 pounds of Technical PCNB per Gallon

### KEEP OUT OF REACH OF CHILDREN CAUTION

"PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que  
le expliquen como usarlo correctamente"

#### STATEMENTS OF PRACTICAL TREATMENTS

**IF SWALLOWED** Call a physician or Poison Control Center immediately. Do not induce vomiting unless  
under the supervision of a physician. Vomiting may cause aspiration pneumonia. This product contains  
an aromatic solvent. **IF IN EYES** Flush eyes with plenty of water. Get medical attention immediately.  
**IF ON SKIN** Remove contaminated clothing and wash affected area with soap and water. Get medical  
attention if irritation persists. **IF INHALED** Remove victim to fresh air. Apply artificial respiration if  
necessary.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG NO. 5481-212

EPA EST NO. 5481-CA-1

NET CONTENTS 1 GALLON

AMVAC CHEMICAL CORPORATION  
4100 EAST WASHINGTON BLVD.  
LOS ANGELES, CA. 90023

**SNAP & DRY BEANS:** For White Mold caused by *Sclerotinia sclerotiorum* APPLY AS A BAND ROW TREATMENT in an 8 inch wide band centered on the row, immediately after or at the time of planting by using 1 gallon mixed in 8 to 10 gallons of water to treat 14,500 linear feet of row for BUSH BEANS or 8,340 linear feet for POLE BEANS. Avoid application directly to seed since delayed emergence may occur. When Dinitro type pre-emergence herbicides are used, add 1 gallon of this product to the final spray mix of the herbicide and apply simultaneously. If disease is severe additional applications may be made at 2 to 3 week intervals using 1 or 2 nozzles per row and directing the spray to the soil at the base of the plants. Apply at the same rate using 1 gallon in 8 to 10 gallons of water to treat 14,500 linear feet of row. Do not apply after the first blooms appear. Do not feed treated vines to livestock.

#### GARLIC

For White Rot caused by *Sclerotium cepivorum* AS A FURROW TREATMENT use 10 gallons of this product mixed in 100 gallons water and apply to 21,800 linear feet of row at planting time. Utilize an appropriate "Z Nozzle" spray rig attached to the planter. Direct the front nozzle to treat the bottom of the seed furrow and the seed and the rear nozzle to treat the soil as the seed is covered.

#### POTATOES

For *Rhizoctonia solani* AS A PRE-PLANTING BROADCAST SOIL TREATMENT use 8 to 12 1/2 gallons per acre mixed with sufficient water to total 25 gallons of finished spray. Apply to the soil surface prior to planting and thoroughly disc to a depth of 4 inches by cross discing. Attach a harrow on the last discing. It is desirable to make this treatment when the soil is slightly moist. AS A FURROW TREATMENT use 5 gallons mixed in 15 gallons of water per 12,400 linear feet of row. Using three nozzles per row place the front "Fan" type nozzle ahead of the furrow opening disc with two "Cone" type nozzles placed above and slightly ahead of the closing disc in the rear. Select nozzles that will deliver about 1/3 of the spray mix per acre through each nozzle at the desired planting speed and pressure.

#### BROWN PATCH IN LAWNS AND ORNAMENTAL TURF

**SOUTHERN GRASSES:** For St. Augustine, Bermuda and similar warm season grasses apply in the fall or spring at the first indication of disease by mowing 1 quart in 10 to 15 gallons of water to treat 1,000 Sq Ft. Select a suitable liquid applicator to give even coverage. Sprinkle the area after treatment to move the chemical to the soil level. If heavy rains or flooding occur retreat the area in 3 to 4 weeks. A temporary discoloration may occur but will disappear in a short time. Do not graze treated turf nor feed clippings to livestock.

#### DICHONDRA

Treat as for grass however use 1 1/2 quarts per 40 gallons of water / 1,000 Sq Ft.

#### ORNAMENTAL AZALEAS AND CAMELLIAS

For Azalea Petal Blight & Camellia Flower Blight use AS A DRENCH TREATMENT to the soil beneath the plants using 1 1/2 quart in 30 gallons of water per 150 Sq Ft. Begin application prior to opening of buds and repeat every 3 to 4 weeks through the bloom period.

NOTICE: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label, when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for use of product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the manufacturer and/or seller and buyer assumes the risk of any such use.

12-28-88

**PRECAUTIONARY STATEMENTS**  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
**CAUTION:** Harmful if swallowed. Avoid inhaling spray mist. This material may cause skin irritation. Wash hands and face after using and before eating or smoking. Avoid contact with skin by wearing the following protective clothing during mixing/loading operations: gloves, long sleeved shirt, long pants, socks and shoes (in addition professional applicators must wear an approved respirator). Use with adequate ventilation. Do not contaminate feed or feedstuffs.  
**ENVIRONMENTAL HAZARDS**  
This pesticide is toxic to fish and aquatic organisms. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water or wetlands when disposing of equipment washwaters. Do not apply directly adjacent to potable water supplies.  
**CHEMICAL AND PHYSICAL HAZARDS**  
Do not use, pour, spill, or store near heat or open flame.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not plant root crops in PCNB treated fields within 12 months of broadcast and banding applications unless PCNB is registered for use on those crops. Do not apply this product through any type of irrigation system.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original container in a cool, dry, locked place out of reach of children. Protect from temperatures below 20 degrees F. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved procedures by State and local authorities.

#### COTTON

For Damping-Off caused by *Rhizoctonia solani* AS A ROW TREATMENT use 1/2 gallon for sandy soil and 1 gallon for organic soils per acre in a minimum of 15 gallons of water for 13,000 linear feet of row based on a 40 inch spacing. Direct the spray toward the seed and surrounding soil at planting time. Do not allow livestock to graze or feed on cotton foliage in PCNB treated fields.

#### BEANS

For Root and Stem Rot caused by *Rhizoctonia solani* AS A FURROW TREATMENT apply 2 to 3 quarts mixed in 8 to 10 gallons of water and apply to 14,500 linear feet of BUSH BEANS or 8,340 linear feet of POLE BEANS. Direct the spray to the base of planting furrow in front of seed drop and to the covering soil at planting time. Do not directly spray bare seed since delayed emergence may result. IN CALIFORNIA & ARIZONA use 3 to 4 parts in 8 to 10 gallons of water and apply to the same linear feet as above indicated for BUSH & POLE BEANS. Direct the spray to the furrow and to the covering soil at planting time. Use the lower rates for sandy and light soils. Do not apply to bare seed.  
(continued on next panel) ->

5481-212  
Pg. 1 of 1