

COMFILE B:54PCB2LF.LAB
EPA FILE SYMBOL: 5481-ERL



5481-215 PM21 1062
PAGE 1 OF 2
This is response
to EPA Letter 12/9/85

ACCEPTED

JUN 2 1986

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 5481-215

P C N B 2 L F

A L I Q U I D F L O W A B L E

S E E D T R E A T M E N T F U N G I C I D E

USE FOR BARLEY, OATS, WHEAT AND COTTON

FOR CONTROL OF SMUT DISEASE AS LISTED AND DAMPING-OFF CAUSED BY RHIZOCTONIA

ACTIVE INGREDIENT:

Pentachloronitrobenzene	20.0%
INERT INGREDIENTS:	80.0%
Total	100.0%

OK This Product Contains 1.70 lbs. PCNB per Gallon
Contains Dye
AN AQUEOUS BASE FLOWABLE LIQUID PRODUCT

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

STATEMENTS OF PRACTICAL TREATMENTS

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and warm water. Get medical attention.

IF INHALED: Remove victim to fresh air and give artificial respiration if indicated. Get medical attention.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 5481-ERL

EPA EST. NO. 5481-CA-1

NET CONTENTS 5 GALLONS

AMVAC CHEMICAL CORPORATION
4100 EAST WASHINGTON BLVD
LOS ANGELES CA 90023

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. This material may cause skin irritation. Do not get in eyes, on skin or on clothing. Wear rubber gloves when loading and mixing. Wash thoroughly after handling and before eating or smoking. Thoroughly wash all contaminated clothing before re-use. Do not contaminate food or feedstuffs. Use only as directed on this label. Do not use treated seed for food, feed or oil purposes.

ENVIRONMENTAL HAZARDS

This material is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, feed or food by storage or disposal. **STORAGE:** Store product in its original container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in sanitary landfill, or by other procedures approved by State and local authorities.

IMPORTANT: Treated seed must be labeled: **WARNING—THIS SEED HAS BEEN TREATED WITH PENTACHLORONITROBENZENE FUNGICIDE. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES. STORE AWAY FROM FEED OR FOODSTUFFS.**

APPLICATION RATES**SEED TREATMENT OF BARLEY, OATS, WHEAT AND COTTON**

Use PCNB 2 LF with suitable Liquid or Slurry type seed treating equipment such as Panogen or Mist-O-Matic and apply at the rates as indicated below. Seed to be used for planting should be sound, fully cured, cleaned and have an excellent germination record. Use the following Seed Treatment Rates/100 LBS.

TYPE OF SEED	AMOUNT PER 100 LBS. SEED		FOR CONTROL OF DISEASE
	FLUID OUNCES	CC'S	
*BARLEY	5 TO 10	148 TO 296	COVERED SMUT (U. kollerii)
*OATS	7 1/2 TO 15	222 TO 444	OAT SMUT (U. avenae)
WHEAT	4	118	COMMON SMUT or BUNT (T. caries/foetida) RHIZOCTONIA Damping-Off
COTTON			SEED ROT & PRE-EMERGENCE DAMPING-OFF caused by RHIZOCTONIA.
Acid Delinted	14	414	
Mech. Delinted	19	562	

*Use higher rate when weather and soil conditions have been or are known to favor the development of soil borne diseases in your locality and when additional seed protection is needed under those soil conditions which benefit the development of these diseases that infect the seed & plant.

By using calibrated liquid seed treatment equipment apply this product to the seed undiluted, directly from the drum. However as long as the above amounts of Active Ingredient are applied per 100 lbs. of seed the product may be diluted using 1 part water with 5 parts of PCNB 2LF. Recalibrate the slurry treater. Thoroughly mix in tank. Do not use treated seed for feed, food or oil purposes. **NOTICE:** Buyer assumes all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.