

COMFILE B:54PRNTRX.LAB #9  
CA STATE REG. NO. 5481-50154-AA  
EPA REG. NO. 5481-03854  
NEW EPA FILE SYMBOL: 5481-



ACCEPTED

APR 30 1986

PRINTREX 2-5G  
GRANULAR HERBICIDE 5481-239

For Weed Control on Non-cropland areas such as Utility, Highway, Pipeline, and Railroad Rights-of-way, Petroleum-Tank Farms, Lumberyards, Storage Areas, Industrial Sites, Firebreaks and around Buildings.

CONTROLS MANY GRASSES AND BROADLEAF WEEDS

ACTIVE INGREDIENTS:

Simazine; 2-chloro-4,6-bis(ethylamino)s-triazine.....	2.00%
Atrazine; 2-chloro-4-ethylamino-6 isopropylamino-s-triazine.	4.75%
Atrazine Related Compounds.....	0.25%
INERT INGREDIENTS:.....	93.00%
Total	100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.  
IF INHALED: Remove victim to fresh air. Get medical attention.  
IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water.  
IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 5481-

EPA EST NO. 5481-CA-1

NET WEIGHT \_\_\_\_\_ POUNDS

AMVAC CHEMICAL CORPORATION  
4100 EAST WASHINGTON BLVD.  
LOS ANGELES CA 90023

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation of dust. Do not get in eyes. Avoid contamination of food and feed. Wash thoroughly after handling and before eating or smoking.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. Simazine is known to leach through the soil, and has been found in ground water as a result of agricultural use. Users are advised not to apply in areas where soils are permeable; i.e. well drained, and which overlie shallow aquifers, particularly those currently being used for drinking water. Consult with the pesticide state lead agency for information regarding soil permeability and aquifer locations in your area. Do not apply directly to water or wetlands.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store product in its original labeled container in a cool, dry, locked place out of reach of children. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

FOR GRASSY AND BROADLEAF WEED CONTROL  
ON INDUSTRIAL SITES, FIREBREAKS, AND RIGHTS-OF-WAY

WHEN TO APPLY

PRINTREX 2-5G Granular Herbicide should be applied before or shortly after weeds emerge. For best weed control, schedule the application just prior to expected rainfall. Since the active ingredients in PRINTREX 2-5G enters the plants mainly through their roots, rainfall or irrigation are needed to move them into the root zone. Very dry soil conditions following application may necessitate follow-up use of a contact herbicide or shallow cultivation. PRINTREX 2-5G is non-corrosive to equipment, non-flammable and has a low electrical conductivity.

WEEDS CONTROLLED

A wide variety of annual and perennial weeds will be controlled such as Barnyard Grass, Bluegrass, Burdock, Canada Thistle, Cheatgrass, Crabgrass, Dog Fennel, Foxtails, Lambsquarter, Orchardgrass, Plantain, Puncture Vine, Purple Top, Quackgrass, Ragweed, Redtopgrass, Smooth Brome, and Turkey Mullein.

HOW MUCH TO APPLY

Apply at rates of from 100 lbs. to 200 lbs. per acre. Where areas to be treated have perennial growth or where soil is heavy-textured, the higher rate should be used. Light soils or where growth of perennials is generally absent, lighter rates may be used. Extensive deep rooted growth areas may require treatment at rates up to 600 lbs per acre. For small areas 4.6 lbs. per 1,000 Sq. Ft. is equivalent to 200 lbs. per acre.

NOTICE: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to manufacturer and /or seller, and buyer assumes the risk of any such use.