\AMVAC\5481-239.LAB HD#9
CA STATE REG. NO. 5481-50154-AA
EPA REG. NO. 5481-239



PAGE 1 OF 2 This Copy is in Response to Changes suggested in Notice of Reg. Dated April 30, 1986

 $A_{\mathcal{O}_{\mathcal{O}_{\mathcal{O}_{\mathcal{D}}}}}$

PRINTREX 2-5G

GRANULAR HERBICIDE

For Weed Control on Non-cropland areas such as Utility, Highway, Pipeline, and Railroad Rights-of-way, Petroleum-Tank Farms, Lumberyards, Storage Areas, Industrial Sites, Firebreaks and around Buildings.

CONTROLS MANY GRASSES AND BROADLEAF WEEDS

ACTIVE INGREDIENTS:

Simazine: 2-chloro-4,6-bis(ethylamino)s-triazine	2.00%
Atrazine: 2-chloro-4-ethylamino-6 isopropylamino-s-triazine.	4.75%
Atrazine Related Compounds	0.25%
INERT INGREDIENTS:	93.00%
Total :	100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with inger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF INHALED: Remove victim to fresh air. Get medical attention.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS EPA REG. NO. 5481-239 EPA EST NO. 5481-CA-1

NET WEIGHT ___50__ POUNDS

AMVAC CHEMICAL CORPORATION 4100 EAST WASHINGTON BLVD. LOS ANGELES CA 90023

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation of dust. Do not get in eyes. Avoid contamination of food and feed. Wash thoroughly after handling and before eating or smoking.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. Simazine and Atrazine are chemicals which can travel (seep or leach) through soil and can contaminate ground water which may be used as drinking water. These chemicals have been found in ground water. Users are advised not to apply this product where the water table (ground water) is close to the surface and where the soils are very permeable, i.e. well-drained soils such loamy sands. Your local agricultural agencies can provide further information on the type of soil in your areas and the location of ground water. Do not apply directly to water or wetlands.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

not contaminate water, food or feed by storage or disposal. STORAGE: Store product in its original labeled container in a cool, dry, locked place out of reach of children. PESTICIDE DISPOSAL: Wastes resulting from the use of this |product may be disposed of on site or at an approved waste disposal facility. | CONTAINER DISPOSAL: Completely empty bag into application equipment. Then |dispose of empty bag in a sanitary landfill or by incineration, or, if allowed| by State and local authorities, by burning. If burned stay out of smoke.

FOR GRASSY AND BROADLEAF WEED CONTROL ON INDUSTRIAL SITES, FIREBREAKS, AND RIGHTS-OF-WAY

WHEN TO APPLY PRINTREX 2-5G Granular Herbicide should be applied before or shortly after weeds emerge. For best weed control, schedule the application just prior to expected rainfall. Since the active ingredients in PRINTREX 2-5G enters the plants mainly through their roots, rainfall or irrigation are needed to move them into the root zone. Very dry soil conditions following application may

cessitate follow-up use of a contact herbicide or shallow cultivation. rRINTREX 2-5G is non-corrosive to equipment, non-flammable and has a low electrical conductivity.

WEEDS CONTROLLED

A wide variety of annual and perennial weeds will be controlled such as Barnyard Grass, Bluegrass, Burdock, Canada Thistle, Cheatgrass, Crabgrass, Dog Fennel, Foxtails, Lambsquarter, Orchardgrass, Plantain, Puncture Vine, Purple Top, Quackgrass, Ragweed, Redtopgrass, Smooth Brome, and Turkey Mullein.

HOW MUCH TO APPLY
Apply at rates of from 100 lbs. to 200 lbs. per acre. Where areas to be treated have perennial growth or where soil is heavy-textured, the higher mate should be used. Light soils or where growth of perennials is generally absent, lighter rates may be used. Extensive deep rooted growth areas may require treatment at rates up to 500 lbs. per acre. For small areas 4.6 lbs. per 1,000 Sq. Ft. is equivalent to 200 lbs. per acre.

NOTICE: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to manufacturer and /or seller, and buyer assumes the risk of any such use.

BEST AVAILABLE COPY