

ACCEPTED
MAR 15 1980
United States Environmental Protection Agency
Pesticide Registration Unit
EPA Reg. No. 5481-258

4% MALATHION DUST

**FOR INSECT CONTROL ON BEANS AND CABBAGE
FOR THE CONTROL OF POULTRY PESTS**

| | |
|---|--------|
| ACTIVE INGREDIENT: | |
| Malathion: 0,0 dimethyl phosphorodithioate of diethyl mercaptosuccinate | 4.0% |
| INERT INGREDIENTS: | 96.0% |
| Total | 100.0% |

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

"PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente"

STATEMENTS OF PRACTICAL TREATMENTS

IF SWALLOWED: Call a physician of Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. Apply Artificial Respiration if indicated. Get medical attention.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 5481-258

EPA Est. No. 5481-CA-1

Net Weight 25 Pound

AMVAC CHEMICAL CORPORATION
4100 EAST WASHINGTON BLVD.
LOS ANGELES CA 90023

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapor or spray mist. Avoid contact with skin. Wash thoroughly after handling and before eating or smoking. Wash contaminated clothing before re-use. Do not contaminate food or feed. **NOTE TO PHYSICIAN:** Malathion may cause cholinesterase inhibition. Atropine is antidotal.

RE-ENTRY AND FARM WORKER SAFETY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons... Do not enter treated areas until dusts have settled. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in treated areas or in an area to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information: "CAUTION: Area treated with 4% MALATHION DUST...on (Date). "Do not enter treated crops without appropriate clothing until dust has settled". When a mixture of 2 or more organophosphate pesticides is applied, the re-entry interval shall be extended by adding to the longest applicable interval, an additional 50% of that interval. In case of accidental exposure refer to STATEMENTS OF PRACTICAL TREATMENT or other Precautionary Statements on this product label. Apply this product only as specific; on this label or on the attached Supplemental Labeling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates and aquatic life stages of amphibians. Do not apply directly to wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original labeled container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay of our smoke.

BEANS: For control of Mexican Bean Beetle, Bean Leafhopper, Spider Mites, Thrips, and Aphids apply at the rate of 30 to 35 pounds per acre. Begin applications early and continue every 7 to 10 days. For best results apply to the undersides of leaves. Do not apply to beans within 3 day of harvest.

CABBAGE: For control of Aphids and Imported Cabbageworms, apply at the rate of 30 to 35 pounds per acre. Begin applications when Aphids or Worms first appear and continue every 7 to 10 days. Do not apply within 7 days of harvest.

POULTRY: For Lice and Mites in Poultry Houses, dust 1 lb. per 40 Sq.Ft. of litter, giving thorough coverage of house, roosts and nests. Replace nesting material after dusting. Do not apply while birds are in house.

NOTICE: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the manufacturer and/or seller, and buyer assumes the risk of any such use.