

# SULFUR FUNGICIDE FOR APPLES

ALSO FOR USE ON CHERRIES, PEACHES, PLUMS & PRUNES

ACTIVE INGREDIENT:  
Sulfur ..... 98.15%  
INERT INGREDIENTS: ..... 1.85%  
Total 100.00%

SEP 27 1989  
5481-300  
EPA Reg. No. 5481-300

**ENVIRONMENTAL HAZARDS**  
Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where run-off is likely to occur.

**DIRECTIONS FOR USE**  
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Sulfur will cause severe fruit and leaf injury to "sulfur sensitive" crops. Do not use on apple varieties such as McIntosh, Golden Delicious, and Jonathan. Do not use on any crop variety unless sulfur has been shown to be safe in your locality. If crop to be treated will be used for canning or freezing purposes or processing consult the food processor before applying SULFUR.

Do not apply when shade temperatures during or within 3 days, exceed, or are likely to exceed, 90°F. Sulfur may burn foliage and fruit during periods of high temperatures (90°F+) and under sunny & high humidity weather conditions.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
STATEMENT OF PRACTICAL TREATMENT  
IF ON SKIN: Wash with soap and water. Get medical attention.  
IF IN EYES: Flush eyes with flowing water for at least 15 minutes.  
Get medical attention if irritation persists.

"PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente"  
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS  
EPA Reg. No. 5481-300 EPA EST. NO. 5481-CA-1

NET WEIGHT 50 POUNDS

**AMVAC CHEMICAL CORPORATION**  
4100 EAST WASHINGTON BLVD  
LOS ANGELES, CA 90023

**STORAGE AND DISPOSAL**  
Do not contaminate water, food or feed by storage or disposal.  
**STORAGE:** Store product in its original labeled, closed container in a cool, dry, locked place out of reach of children.  
**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.  
**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

**WHEN TO USE**  
Apply to crops as needed in season when diseases first appear and plant disease damage is eminent. Repeat every 7 to 14 days or as directed for continued control. Contact your State Agricultural Extension Service for crop dusting schedules giving the proper timing to make applications in your locality. Do not apply within 10 to 14 days of an oil spray.

**HOW MUCH TO USE**  
Use 35 to 50 pounds of dust per acre. Apply dust uniformly and thoroughly to fruit and foliage. May be applied by aircraft or ground equipment. Make applications when air is quiet.  
1. Use the lower rate - as indicated, during periods of minimal to moderate disease infestation, for regular season scheduled sprays and as treatments are necessary when temperatures are at, and above 90°F  
2. Use the higher rate, as indicated, when diseases are damaging the crop and weather conditions are favorable for their continued plant damage and when temperatures will not exceed 90°F within the following 3 days after spraying. Sulfur will burn most plants in sunny 95°F+ temperatures.

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION:** May be harmful if swallowed, inhaled, or absorbed through the skin. Avoid inhalation of dust. Avoid contact with eyes, skin and clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.  
**RE-ENTRY AND WORKER SAFETY STATEMENT**  
Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas for 24 hours unless appropriate protective clothing is worn. Because certain states may require more restrictive re-entry intervals for various crops treated with this product consult your State Agricultural Extension Service for further information.  
\*Hat, long sleeved shirt, long pants, shoes, socks.  
Written or oral warnings must be given to workers who are expected to be in treated areas or in an area to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure.)  
When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information  
**CAUTION:** Area treated with SULFUR FUNGICIDE on (Date) . Do not enter treated areas for 24 hours unless appropriate protective clothing is worn. In case of accidental exposure refer to STATEMENTS OF PRACTICAL TREATMENT or other Precautionary Statements on SULFUR FUNGICIDE for APPLES product label. EPA Reg. No. 5481-300.

**APPLES:** For Powdery Mildew and Apple Scab use in delayed dormant, pink, petal fall, calyx and cover applications as required to maintain control. See paragraph on WHEN TO USE and follow schedules furnished by your State Agricultural Extension Service.

**CHERRY:** For \*Brown Rot, and Powdery Mildew apply at pink bud. Repeat at full bloom and before shuck shed or every 4 days in cool, wet weather. For \*\*Blossom blight, apply at popcorn, full bloom and petal fall. For Powdery Mildew use at 7 to 10 day intervals for cover applications.

**PEACHES:** For \*Brown Rot and Powdery Mildew: For \* "Blossom Blight" control apply at pink, full bloom and just before shuck fall. For \*\*"Fruit rot" and Powdery Mildew control apply at 7 to 10 day intervals the remainder of the season as necessary.

**PLUMS AND PRUNES:** For \*Brown Rot - For \*\*Blossom Blight control apply at pink, full bloom and pre-shuck fall or at 4 day intervals during cool wet weather. For \*Fruit rot control apply 5 weeks before harvest at 7 to 10 day intervals the remainder of the season as necessary.

**NOTICE:** This product conforms to its chemical description and is reasonably fit for the purpose stated on the label, when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or

**BEST AVAILABLE COPY**