

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

AUG 04 1992

CERTIFIED MAIL

Anvac Chemical Corp.  
4100 E. Washington Blvd.  
Los Angeles, CA 90023

S-412413 335  
38

Attention: Mr. Jack Prieur

*Computer notes*  
*only take 12*

Subject: Phosdrin 4 EC  
EPA Reg. No. 5481-412  
Deletion of Greenhouse Use

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable, subject to the following provisions:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Submit one copy of your final printed labeling before you release the product for shipment.
3. As provided for in the enclosed Federal Register Notice (57 FR 18146-7), the registrant is not authorized to sell or distribute any existing stocks of this product with the previously approved labeling.

This registration will be subject to cancellation in accordance with FIFRA sec. 6(e) if you do not comply with these conditions. Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

*RAF*


Robert A. Forrest  
Product Manager (14)  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

- Enclosures:
1. Stamped label
  2. A-79 Enclosure
  3. Federal Register Notice (18146-7)

RAF:dbp:07/31/92:Peacock WPS ENCLOSURES 2:305-5407,-6600

SYMBOL							
SURNAME							
DATE							

**BEST AVAILABLE COPY**



**RESTRICTED USE PESTICIDE**  
 Due to Very High Acute Toxicity to Humans and  
 Severe Effects on Fish, Mammals and Aquatic  
 Species, For Retail Sale, Use Only by Cer-  
 tified Applicators or Persons Under Their Direct  
 Supervision and Only for Those Uses Covered by  
 the Certified Applicators Certificate. Do not use  
 when the product is defined as "the restricted  
 applicator being physically present during applica-  
 tion, mixing, loading, repair and cleaning of ap-  
 plication equipment." Certified applicators must  
 also ensure that all persons involved in these ac-  
 tivities are informed of the precautionary  
 statements.

# Phosdrin 4 EC

INSECTICIDE

Emulsifiable Concentrate - Contains 4 lbs. active ingredients per Gallon

**ACTIVE INGREDIENTS:**

Mevinphos (2-Carbomethoxy-1-methylvinyl dimethyl phosphite, alpha isomer).....	28.3%
Related Compounds.....	18.8%
<b>INERT INGREDIENTS:</b> .....	52.9%

\*Equivalent of 47.1% w MEVINPHOS Insecticide Total 100.0%

Phosdrin is a Registered Trade Mark of the AMVAC CHEMICAL CORPORATION

EPA REG. NO. 5481-412

EPA EST. NO. 5481-CA-1

**KEEP OUT OF REACH OF CHILDREN**



**DANGER - POISON**  
**PELIGRO**



"PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado completamente"

**STATEMENTS OF PRACTICAL TREATMENT**

ATROPINE IS AN ANTIDOTE, CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING

**IF SWALLOWED:** drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. NEVER induce vomiting or give anything by mouth to an unconscious or non-alert person.  
**IF INHALED:** remove from exposure. Have patient lie down and keep quiet. If patient is not breathing start artificial respiration immediately. Get medical attention.  
**IN CASE OF CONTACT,** wash skin with plenty of soap and water; for eyes, flush with water and get medical attention; remove and wash contaminated clothing before re-use.  
**IF WARNING SYMPTOMS APPEAR:** (See WARNING SYMPTOMS Under "NOTE TO PHYSICIAN") Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest hospital.

For medical emergencies involving this product, call 1-(213) 264-3910

SEE ANTIDOTE AND OTHER PRECAUTIONS ON SIDE PANEL

**PRECAUTIONARY STATEMENTS**

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**DANGER!** Poisonous if swallowed, inhaled or absorbed through the skin. Rapidly absorbed through the skin. Repeated inhalation or skin contact may without symptoms, progressively increase susceptibility to PHOSDRIN insecticide poisoning. Do not swallow or get in eyes, on skin or on clothing. Do not breathe vapors. Do not contaminate food or feed products. Keep away from heat or open flame.

**PRECAUTIONS IN USING**

**USE ONLY WHEN WEARING THE FOLLOWING PERSONAL PROTECTIVE EQUIPMENT DURING MIXING/LOADING, APPLICATION, REPAIR AND CLEANING OF MIXING, LOADING, AND APPLICATION EQUIPMENT, AND DISPOSAL OF PESTICIDE:** protective suit of one or two pieces that covers all parts of the body except head, hands, and feet; chemical resistant gloves; chemical resistant shoes (or chemical resistant shoe coverings or chemical resistant boots); goggles or face shield; hood or wide brimmed hat; NIOSH or MSHA approved respiratory protective device. In the case of aerial application, the pilot should not assist in the mixing and loading operation.  
**IF MIXING/LOADING IS PERFORMED USING A CLOSED SYSTEM, THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT MAY BE WORN AS AN ALTERNATIVE:** Long sleeved shirt; long legged pants; chemical resistant gloves; chemical resistant apron; shoes and socks; Goggles or face shield must be worn when the system is under pressure. All other protective clothing and equipment required for use with open systems must be available nearby.  
**IF APPLICATION IS PERFORMED USING AN ENCLOSED CAB OR COCKPIT, THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT MAY BE WORN AS AN ALTERNATIVE:** Long sleeved shirt and long legged pants; shoes and socks. Chemical resistant gloves must be available in the cab or cockpit and must be worn during entry to and exit from the application vehicle. All other protective clothing and equipment required for use during application must be available in the cab and must be worn when exiting the cab into the treated area. When used for this purpose, compressed clothing may not be brought back into the cab unless in an enclosure such as a plastic bag.  
**REMEMBER - THIS CLOTHING IS NOT ADEQUATE TO PROTECT YOU DURING REPAIR AND CLEANING OF APPLICATION EQUIPMENT AND EARLY REENTRY TO TREATED AREAS REFER TO PROTECTIVE CLOTHING AND EQUIPMENT REQUIREMENTS ABOVE.**  
**IMPORTANT!** If pesticide comes in contact with skin, wash off with soap and water and contact a physician immediately. Always wash hands, face, and arms with soap and water before smoking, eating, drinking, or sleeping.  
**AFTER WORK:** Before removing gloves, wash them with soap and water. Take off all work clothes and shoes. Shower using soap and water. Wear only clean clothes when leaving job - do not wear contaminated clothing. Personal and protective clothing worn during work must be stored and laundered separately from household articles. Clean or launder protective clothing after each use. Respirators must be cleaned and filters replaced according to instructions included with the respirator. Protective clothing and protective equipment that becomes heavily contaminated or discolored with mevinphos must be destroyed according to State and local regulations. **HEAVILY CONTAMINATED OR DISCOLORED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED.**

The National Pesticide Telecommunication Network is available for recommendations regarding pest management, emergency treatment, and other information regarding the toxicity of mevinphos. The toll free number for the National Pesticide Telecommunication Network is 1-800-868-7378.

**IF HANDLED INDOORS** provide mechanical exhaust ventilation. Keep all unprotected persons, children, livestock and pets away from treated areas or where there is danger of drift. Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get medical help right away. See the First Aid (Practical Treatment) section on this label.

Do not contaminate or store near food or feedstuff. Note Reentry Statements Section. **IMPORTANT:** Not for use or storage in or around the home. Keep animals and unprotected persons out of operational areas and while there is a danger of drift.

In case of spillage, decontaminate areas and equipment by washing with a dilute solution of alkali (less than 5%) and detergent and rinse with water. In case of significant spill, call CHEMTREC (800) 424-9300.

**NOTE TO PHYSICIAN**

**WARNING SYMPTOMS:** Symptoms include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

**TREATMENT:** Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation Atropine Sulfate should be injected at 30 minute intervals in doses of 1 to 2 milligrams until complete atropinization has occurred.

Prothiames chloride (2-PAM chloride) may also be used as an effective antidote in addition to and while maintaining full atropinization. In adults, an initial dose of 1 gram of 2-PAM should be injected, preferably as an infusion, in 250 cc of saline over a 15 to 20 minute period. If this is not practical, 2-PAM may be administered slowly by intravenous injection as a 5% solution in water over not less than 2 minutes. After about an hour, a second dose of 1 gram of 2-PAM will be indicated if muscle weakness has not been relieved. For infants and children, the dose of 2-PAM is 0.25 grams.

Morphine is an improper treatment.

Clear effect by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase regeneration has been attained as determined by a blood test.

**FOR MEDICAL EMERGENCIES INVOLVING THIS PRODUCT, CALL 1-(213) 264-3910.**

**ENVIRONMENTAL HAZARDS**

**Terrestrial Food and Nonfood Uses.**

This product is toxic to fish and wildlife. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

Soils feeding on treated areas may be killed.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur.

In case of a significant spill, call CHEMTREC 1-(800) 424-9300.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**REENTRY STATEMENT**

Do not apply this product in a manner so directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

**HUMAN FLAGGERS ARE PROHIBITED** during aerial application of this product unless in totally enclosed vehicles.

**Do not treat CITRUS GROVES, GRAPE VINEYARDS, and NECTARINE and PEACH orchards** for 24 hours (4 days) after the end of application, unless the protective clothing specified on this label for early reentry is worn. **Reentry into all other treated areas is prohibited for 48 hours after the end of application, unless the protective clothing specified on this label for early reentry is worn.**

**FOR EARLY REENTRY INTO TREATED AREAS BEFORE SPRAYS HAVE DRIED,** Wear all protective clothing specified on this label for an applicator.

**FOR EARLY REENTRY INTO TREATED AREAS AFTER SPRAYS HAVE DRIED,** Wear protective suit of one or two piece covering all parts of the body (shirt, pants, shoes, socks, gloves, chemical resistant shoes (or chemical resistant shoe coverings or chemical resistant boots).

Some States may require more restrictive reentry intervals for various crops treated with this product. Contact your State Department of Agriculture for further information.

Oral warnings must be given to workers who are expected to be in treated areas or in an enclosure with this product. (Oral warnings which require workers of areas treated with this product to take specific protective clothing, period of time limits must be given to take in case of accidental exposure.) When oral warnings are given they shall be given in a language customarily understood by workers. Oral warnings must include the following information:

Do not enter without protective clothing until (insert date/time reflecting end of reentry interval set forth on label) in case of accidental exposure see STATEMENTS OF PRACTICAL TREATMENT found on AEC product label.

**USE AND APPLICATION DIRECTIONS**

Do not apply this product through any type of irrigation system.  
 Phosdrin 4EC can be used in conventional hydraulic sprayers, low-volume ground applicators, and backpack sprayers. Use the product only for recommended purposes and at recommended dosage. For more information consult the State Agricultural Extension Service for exact dosage and timing recommendations. Mix with water by vigorous agitation; agitate the mixture while spraying. If allowed to settle, agitate before use. Apply when insects first appear and repeat application as often as necessary for pest control. Apply in a suitable volume of water to give uniform coverage of foliage. For control of caterpillars, dilute this product with 3 to 10 gallons of water per acre.

**READ THE DIRECTIONS CAREFULLY AND FOLLOW THEM AT ALL TIMES**

BEST AVAILABLE COPY

100 . 11

999

ACCEPTED  
AUG 04 1992  
Under the Federal Information  
Processing and Statistics Act  
in accordance with the provisions  
of 5 U.S.C. 552, EPA Reg. No. 5491-412

htz

494

**FRUIT AND NUT CROPS:** Use recommended dosages in sufficient water to obtain thorough spray coverage, but do not exceed indicated maximum gallons for that crop. If concentrate ground application is employed, do not apply less than 10 gallons of spray per acre, and adjust dosage of Phosdrin 4EC to obtain same dosage per acre as recommended for full coverage application.

**FIELD AND VEGETABLE CROPS:** Use 10 to 125 gallons of spray per acre when using ground application equipment.

Avoid drift of this product to crops for which it is not registered or to crops for which its use may be registered but application would be closer to harvest than recommended.

## FRUIT AND NUT CROPS

**APPLES, PEACHES, PEARS, PLUMS:** For control of Aphids, and Mites use 1/4 to 1/2 pt. per 100 gals. water. Do not exceed 1,200 gallons of finished spray per acre on apples and pears when 1/2 pt. is used. For control of Grasshoppers, Lygus Bugs, and Red-Banded Leaf Roller Use 1/2 to 3/4 pt. per 100 gals. water. Do not exceed 800 gals. of finished spray per acre on apples and pears when 1/2 pt. is used. Do not treat within 1 day of harvest. Do not exceed 750 gals. finished spray per acre on peaches or plums.

**CHERRIES (SOUR ONLY):** For control of Aphids and Mites use 1/2 to 3/4 pt. per 100 gals. water. Do not treat within 2 days of harvest. Do not exceed 750 gals. finished spray per acre.

**CITRUS (ORANGES, LEMONS, GRAPEFRUIT):** For control of Aphids use 1 to 2 pts. per acre in 200 or more gals. water. For control of Fruit Tree Leafroller, Orange Tortrix and Omnivorous Leafroller, use 1 qt. per acre in 500 gals. water. For control of Western Tussock Moth larvae, Citrus Cutworm, Variegated Cutworm, and Pink Scavenger Caterpillar use 2 qt. per acre in 1,200 gals. water. Allow at least 7 days between applications. Do not treat within 1 day of harvest.

**GRAPES:** For control of Aphids use 1/2 to 1 pt. per acre. Do not treat within 2 days of harvest. For control of Leaf Folder, Leafhoppers, Mites, Red-Banded Leafroller and Lygus Bugs use 1 to 2 pts. per acre. At the 2 pt. dosage do not treat within 5 days of harvest.

**RASPBERRIES:** For control of Aphids, use 1/4 to 1/2 pt. per 100 gals. water. For control of Mites, Leafhoppers, Fruit Tree Leafroller and Orange Tortrix, use 1/2 to 3/4 pt. per 100 gals. water. Apply both dosages in up to 200 gals. water per acre. Do not treat within 3 days of harvest.

**STRAWBERRIES:** For control of Aphids and Mites use 1/4 to 1/2 pt. per 100 gals. water. For control of Grasshoppers, Strawberry Leafroller, Salt-marsh Caterpillar and Lygus Bugs use 1/2 to 1 pt. per 100 gals. water. Do not treat within 1 day of harvest. For hard-to-kill insects, including Aphids and Mites, use 1 qt. per 100 gals. water, but do not treat within 2 days of harvest. Do not exceed 100 gals. finished spray per acre.

**WALNUTS:** For control of Aphids, use 1/4 to 1/2 pt. per 100 gals. water. For control of Mites, Omnivorous Leafroller, Orange Tortrix, Fruit Tree Leafroller, and Western Tussock Moth use 1/2 pt. per 100 gals. water. Do not treat within 1 day of harvest. Do not exceed 1,000 gals. finished spray per acre.

## FIELD CROPS

**ALFALFA, CLOVER:** For control of Aphids and Alfalfa Caterpillar, use 1/4 to 1/2 pt. per acre. For control of Grasshoppers, Leafhopper, Cutworms (climbing), Mites, Alfalfa Weevil Larvae and Lygus Bugs use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

**CORN (FIELD & POPCORN) FOR FORAGE & GRAIN:** For control of Aphids, use 1/4 to 1 pt. per acre. Do not treat within 1 day of harvest.

**PHOSDRIN 4EC** may cause a red or purple discoloration of the foliage on corn plants similar to other organophosphate compounds.

**MUSTARD GREENS, TURNIP TOPS:** For control of Aphids use 1/4 to 1/2 pt. per acre. For control of Cabbage Looper, Imported Cabbage Worm, False Chinch Bug, Dipterous Leafminer (adult), Grasshoppers, Leafhoppers and Mites use 1/2 to 1 pt. per acre. Do not treat within 3 days of harvest.

**PEA VINES (FOR FORAGE ONLY):** For control of Aphids, use 1/4 to 1/2 pt. per acre. For control of Grasshoppers, Leafhoppers, Mites and Cutworms (climbing), use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

**SORGHUM (FOR FORAGE AND GRAIN):** For control of Aphids use 1/4 to 1/2 pt. per acre. For control of Corn Earworm, and Webworm use 1/2 to 1 pt. per acre. For control of Fall Armyworms use 1 pt. per acre. Do not treat within 3 days of harvest.

**PHOSDRIN 4EC** may cause a red or purple discoloration of the foliage on sorghum plants similar to other organophosphate compounds.

## VEGETABLE CROPS

**ARTICHOKES:** For control of Aphids use 1/4 to 1/2 pt. per acre. For control of Plume Moths, use 1 to 2 pts. per acre. Do not treat within 2 days of harvest.

**BEANS:** For control of Aphids use 1/4 to 1/2 pt. per acre. For control of Grasshoppers, Leafhoppers, Mites, and Mexican Bean Beetle use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

**BEETS (INCLUDING TOPS):** For control of Aphids, use 1/4 to 1/2 pt. per acre. For control of Cabbage Looper, Cutworm (climbing), Dipterous Leafminer (adult), False Chinch Bug, Grasshoppers, Imported Cabbage worm, Leafhoppers, Mites and Salt-marsh caterpillars, use 1/2 to 1 pt. per acre. Do not treat within 3 days of harvest.

**BROCCOLI, CABBAGE:** For control of Aphids use 1/4 to 1/2 pt. per acre. For control of Cabbage Looper, Imported Cabbage worm, Grasshoppers, Leafhoppers, Salt-marsh Caterpillar, Mites, Cutworms (climbing), Dipterous Leafminer (adult), and Lygus Bugs use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest. For hard-to-kill insects, including Aphids, use 1 qt. per acre, but do not treat within 3 days of harvest.

**BRUSSEL SPROUTS, CAULIFLOWER, COLLARDS, KALE:** For control of Aphids, use 1/4 to 1/2 pt. per acre. For control of Cabbage Looper, Imported Cabbage worm, Grasshoppers, Leafhoppers, Salt-marsh Caterpillar, Mites, Cutworms (climbing), Dipterous Leafminer (adult), and Lygus Bugs use 1/2 to 1 pt. per acre. Do not treat within 3 days of harvest. For hard-to-kill insects, including Aphids, use 1 qt. per acre, but do not treat COLLARDS and KALE within 7 days of harvest.

**CARROTS:** For control of Aphids, use 1/4 to 1/2 pt. per acre. For control of Leafhoppers, Lygus Bugs, Mites, Cabbage Looper, Dipterous Leafminer (adult), Cutworms (climbing), and Salt-marsh Caterpillar use 1/2 to 1 pt. per acre. Do not treat within 2 days of harvest.

**CELERY:** For control of Aphids, use 1/4 to 1/2 pt. per acre. For control of Dipterous Leafminer (adult), Lygus Bugs, Salt-marsh Caterpillar, Leafhoppers, Cabbage Loopers and Mites, use 1/2 to 1 pt. per acre. Do not treat within 3 days of harvest. For hard-to-kill insects, including Aphids, use 1 qt. per acre, but do not treat within 5 days of harvest.

**CORN, (SWEET) FOR GRAIN & FORAGE:** For control of Aphids use 1/4 to 1 pt. per acre. Do not treat within 1 day of harvest.

**PHOSDRIN 4EC** may cause a red or purple discoloration of the foliage on corn plants similar to other organophosphate compounds.

**CUCUMBER (OUTDOOR):** For control of Aphids, use 1/4 to 1/2 pt. per acre. For control of Grasshoppers, Leafhoppers, and Mites, use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

**EGGPLANT, PEPPERS:** For control of Aphids, use 1/4 to 1/2 pt. per acre. For control of Grasshoppers, Leafhoppers, and Mites, use 1/2 to 1 pt. per acre. Do not treat within 2 days of harvest.

**LETTUCE:** For control of Aphids, use 1/4 to 1/2 pint per acre. For control of Corn Earworm, Cutworms (climbing), Dipterous Leafminer (adult), Cabbage Looper, Imported Cabbage Worm, Grasshoppers, Mites, Lygus Bugs, Salt-marsh Caterpillar, False Chinch Bug, and Thrips, use 1/2 to 1 pint per acre.

Keep a 7 day interval between applications.

**Head Lettuce:** When applying 1 pint or less per acre, do not treat within 2 days of harvest. For hard-to-kill insects, including Aphids, use 1 quart per acre but do not treat within 4 days of harvest.

**Leaf Lettuce (including Butter or Boston, Greenleaf, Redleaf and Romaine):** When applying 1 pint or less per acre, do not treat within 7 days of harvest. For hard-to-kill insects, including Aphids, use 1 quart per acre but do not treat within 10 days of harvest.

**MELONS (INCLUDING CANTALOUPE, HONEYDEW MELONS, MUSKMELONS, WATERMELONS):** For control of Aphids, use 1/4 to 1/2 pt. per acre. For control of Cabbage Looper, Dipterous Leafminer (adult), Leafhoppers, Lygus Bugs, Mites, False Chinch Bug, Salt-marsh Caterpillar, Cutworms (climbing), and Grasshoppers use 1/2 to 1 pt. per acre. To control *Rhinoceros* (Cabbage Looper, Cutworms, Salt-marsh Caterpillars, Tobacco Budworms) on WATERMELONS apply 1/2 pt. per acre. Do not treat CANTALOUPE, HONEY-DEW MELONS, MUSKMELONS OR WATERMELONS within 1 day of harvest.

**OKRA:** For control of Aphids, use 1/4 to 1/2 pt. per acre. For control of Cutworms (climbing), Corn Earworm, Green Stink Bug, Mites and Velvet Bean Caterpillar, use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

**ONIONS (INCLUDING GREEN ONIONS):** For control of Thrips and Cutworms (climbing) use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

**PARSLEY:** For control of Aphids, apply 1 to 2 pt. (0.5 lb. to 1.0 lb. a.i./acre) as foliage application using a minimum of 40 gallons of water per acre by ground application or 10 to 15 gallons of water per acre by air application. For light infestations use lower dosage of 1 pt. (0.5 lb. a.i./acre). Do not treat later than 5 days before harvest. For hard-to-kill insects and increased populations use 2 pt. (1.0 lb. a.i./acre). Do not treat later than 8 days before harvest for higher dosage. Begin application when insects first appear and repeat as often as necessary to maintain control. Do not treat more than 3 times between harvest.

**PEAS (INCLUDING VINES):** For control of Aphids, use 1/4 to 1/2 pt. per acre. For control of Grasshoppers, Leafhoppers, Mites and Cutworms (climbing), use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

**POTATOES:** For control of Aphids, use 1/4 to 1/2 pt. per acre. For control of Grasshoppers, Leafhoppers, and Mites use 1/2 pt. per acre. Do not treat within 1 day of harvest.

**SPINACH:** For control of Aphids, use 1/4 to 1/2 pt. per acre. For control of Cabbage Looper, Imported Cabbage worm, Grasshoppers, Leafhoppers, Mites, Dipterous Leafminer (adult), Cutworms (climbing), Salt-marsh Caterpillar, and False Chinch Bug use 1/2 to 1 pt. per acre. Do not treat within 4 days of harvest. For hard-to-kill insects, including Aphids, use 1 qt. per acre but do not treat within 7 days of harvest.

**SUMMER SQUASH:** For control of Aphids, use 1/4 to 1/2 pt. per acre. For control of Cabbage Looper, Dipterous Leafminer (adult), Leafhoppers, Lygus Bugs, Mites, False Chinch Bug, Salt-marsh Caterpillar, Cutworms (climbing), and Grasshoppers, use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

**TOMATOES (OUTDOOR):** For control of Aphids, use 1/4 to 1/2 pt. per acre. For control of Grasshoppers, Leafhoppers, and Mites use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

**TURNIPS:** For control of Aphids, use 1/4 to 1/2 pt. per acre. For control of Cabbage Looper, Imported Cabbage worm, Grasshoppers, Leafhoppers, Mites, False Chinch Bug and Dipterous Leafminer (adult), use 1/2 to 1 pt. per acre. Do not treat within 3 days of harvest.

**BEST AVAILABLE COPY**

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Open dumping is prohibited.

**STORAGE:** Do not store in or around the home. Store in a secure, dry and temperate area. Store in original container only. Product is flammable. Keep away from heat, sparks or open flame. If spilled, avoid exposure. If significant spill, call CHEMTREC (800) 424-9300. Spill can be contained by covering with a sufficient amount of absorbent material such as clay, diatomaceous earth, sand, sawdust, or sweeping compound. Place in drum, label contents, and dispose of according to the instructions below in PESTICIDE DISPOSAL. . . . .

**CAUTION:** Wear all personal protective equipment specified in the PRECAUTIONS IN USING section of this label including a pesticide respirator when tending dust clouds during clean-up.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If the wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** (Metal) Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture & dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. (Plastic) Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke. . . . .

**NOTICE:** This product conforms to its chemical description and is reasonably fit for the purpose stated on the label, when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the manufacturer and / or seller and the buyer assumes the risk of any such use. 899