

ALCO

K-SALT FRUIT FIX 800

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be harmful if swallowed, inhaled or absorbed through the skin. Causes substantial but temporary eye irritation. Avoid contact with skin, eyes or clothing. Do not breathe spray mist. Wear goggles or safety glasses, full length trousers, long-sleeved shirt and head covering. In case of eye contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse.

RE-ENTRY AND WORKER SAFETY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in treated areas or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written and oral warnings must include the following information. **WARNING:** Area treated with K-SALT FRUIT FIX 800 on (Date) _____. Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, refer to the Precautionary Statements found on the K-SALT FRUIT FIX 800 product label.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters. Avoid drift of spray mist.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original container, in a cool, (not in direct sunlight), dry, locked place out of reach of children. Do not allow product to freeze. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

K-SALT FRUIT FIX 800 controls preharvest drop of APPLES and PEARS, leading to larger harvest and a reduction of losses from windfall and "knockdown".

A HORMONE TYPE PRODUCT FOR CONTROL OF PREHARVEST DROP OF APPLES AND PEARS

ACTIVE INGREDIENT:

1-Naphthalene Acetic Acid, Potassium Salt - 24.20%

INERT INGREDIENTS: - - - - - 75.80%

Total 100.00%

Equivalent to 20.2% 1-Naphthaleneacetic Acid
or 800 grams (1.76 lbs.) per gallon

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicado ampliamente

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED - Call a physician or Poison Control Center immediately. Induce vomiting by drinking 1 or 2 glasses of water and touching back of throat. Do not give anything by mouth to an unconscious or convulsing person. **IF IN EYES** - Flush eyes with plenty of water. Get medical attention if irritation persists. **IF ON SKIN** - Wash thoroughly with soap and water. **IF INHALED** - Remove to fresh air.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

EPA Reg. No. 5481-URG(413)

EPA Est. No. 5481-CA-1

NET CONTENTS 1 GALLON

AMVAC CHEMICAL CORPORATION
1100 EAST WASHINGTON BLVD
LOS ANGELES, CALIFORNIA 90023

GENERAL DIRECTION FOR USE ON APPLES AND

Spray trees thoroughly, wetting all fruit stems and adjacent leaves 7 to 10 days before harvest on APPLES and 5 to 7 days before harvest on PEARS. K-SALT FRUIT FIX 800 becomes effective in 3 to 4 days after application and remains effective for 3 to 4 weeks. Separate sprays should be applied to different varieties. Do not make more than 2 applications and do not spray within 2 days of optimum maturity. Do not spray within 2 days of harvest.

HOW TO APPLY WITH GROUND EQUIPMENT FOR APPLES, thoroughly mix 3 fluid ounces with 250 (sp. per 100 gals. water) to prepare a 20 ppm spray. The trees using up to 465 gallons per acre of this 20 ppm spray equivalent of 35 grams of 1-Naphthaleneacetic acid per acre.

FOR PEARS, thoroughly mix 3 fluid ounces with 500 gal (per 100 gals. water) to prepare a 10 ppm spray. Thoroughly wet trees using up to 670 gallons of this 10 ppm spray dilution per acre, which is equivalent of 25 grams of 1-Naphthaleneacetic acid per acre. Apply to trees 10 to 15 days before harvest. Do not make more than 2 applications.

HOW TO APPLY AS AN AERIAL APPLICATION

Aircraft should utilize spray equipment which gives thorough coverage of all portions of trees. Apply only when weather conditions favor spraying. Use lower rates for normal foliage and fruiting; the higher rates for heavy fruiting.

APPLES and D'ANJOU PEARS: Use 1/4 to 1/2 pint (per 100 gals. water) at least 5 gallons of water per acre. Do not apply within 2 days of harvest.

BARTLETT AND BOSCH PEARS: Use 1/8 to 1/4 pint (per 100 gals. water) at least 5 gallons of water per acre. Do not apply within 2 days of harvest. If BARTLETT and BOSCH are interplanted, spray only once. Use 1/2 pint per acre on BARTLETT to prevent pre-mature ripening.

FOR USE IN PRESERVING ORNAMENTAL HOLLY BOUGHS

To delay leaf drop of Holly boughs make a "dip" solution of 33 (2 to 4 fluid ounces) in 100 gallon of water; (equals 33 ppm) as soon as possible after cutting. Allow excess solution to drain from boughs in dip solution. Replenish dip solution as needed to maintain concentration. Replace with fresh solution every 3 to 4 days to keep boughs fresh until used. Maintain a 34 to 40 °C bough storage area.

NOTICE: This product conforms to its chemical description for the purpose stated on the label, when used in accordance with the normal conditions of use. Manufacturer is not responsible for product contrary to the label instructions, or under abnormal conditions not reasonably foreseeable to the manufacturer. Buyer assumes the risk of any such use. (07-01-90 Revision)

