



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

July 9, 2021

Julie Porter
Director of Regulatory Compliance
AMVAC Chemical Corporation
4695 MacArthur Court, Suite 1200
Newport Beach, CA 92660

Subject: Label Amendment – Incorporate 24(c) Language (EPA SLN Nos.: OR-010002 and OR-010003) and for Remove Non-marketed Uses [Mandolin (EPA Reg. No. 628)] on the Section 3 Registration
Product Name: K-Salt Fruit Fix 800
EPA Registration Number: 5481-413
Application Date: February 24, 2021
Decision Number: 572835

Dear Ms. Porter:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 5481-413
Decision No. 572835

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Ernest Kraka by phone at 7033478455, or via email at kraka.ernest@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Shaja B. Joyner". The signature is written in a cursive style with a large initial 'S'.

Shaja B. Joyner, Product Manager 20
Fungicide-Herbicide Branch
Registration Division 7505P

Enclosure

ACCEPTED

07/09/2021

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 5481-413

K-SALT® FRUIT FIX® 800

A HORMONE TYPE PRODUCT FOR THINNING AND CONTROL OF PRE-HARVEST DROP OF APPLES AND PEARS

ACTIVE INGREDIENT:

1-Naphthaleneacetic Acid, Potassium Salt 24.2%

INERT INGREDIENTS: 75.8%

TOTAL: 100.0%

Equivalent to 20.2% 1-Naphthaleneacetic Acid (1-NAA) or 830 grams (1.83 lbs.) 1-NAA per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

EMERGENCY INFORMATION

[Emergency Information for K-Salt Fruit Fix 800]

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY:

For Medical Emergencies phone:.....1-800-892-0099

For Transportation Emergencies, including spill, leak or fire, phone: CHEMTREC®1-800-424-9300 or 1-202-483-7616

For Product Use Information phone: VALENT.....1-800-682-5368

[Alternate Emergency Information for Alternate Brand Names]

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY:

For Medical Emergencies phone:.....1-888-681-4261

For Transportation Emergencies, including spill, leak or fire, phone: CHEMTREC®1-800-424-9300

For Product Use Information phone: AMVAC®1-888-462-6822

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE.

EPA Reg. No. 5481-413

EPA Est. No. _____

413-20210505

Net Contents:

As Marked on Container



AMVAC Chemical Corporation
4695 MacArthur Court, Suite 1200
Newport Beach, CA 92660 U.S.A.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Harmful if swallowed, absorbed through skin or inhaled. Do not get in eyes, or on skin or clothing. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of any waterproof material.

All mixers, loaders, applicators, flaggers and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks
- Chemical-resistant gloves (except for pilots), and
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters. Do not contaminate irrigation ditches or water used for irrigation or domestic purposes. Do not apply when conditions favor drift from treated areas. Do not use in a greenhouse.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected persons may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

K-SALT FRUIT FIX 800 thins the fruit of apples and pears to provide an adequate return bloom the following season. K-SALT FRUIT FIX 800 controls pre-harvest drop of apples and pears leading to a larger harvest and a reduction of losses from windfall and knockdown. Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with this labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

K-SALT FRUIT FIX 800 is an auxin-type hormone product for use in fruit production. The response of fruit trees to application may be varied in different years. In reviewing the results, the grower needs to consider the wide variation in varieties, climatic factors, dosage, tree vigor, and fruit set before choosing the desired rate and correct timing for application.

Apply in sufficient water for good even coverage with your specific equipment. Application can be made in as little as 50 gallons of water per acre and up to 400 gallons of water per acre and is highly dependent on type of equipment used and density of the foliage. Uniform and even distribution of the spray is imperative. For rates and application timing for your specific variety and location, consult your local Extension Pomologist for recommendations.

MANDATORY SPRAY DRIFT

Aerial Applications:

- Do not apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Applicators must select nozzle and pressure that deliver medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 641 (ASABE S641). If the windspeed is 10 miles per hour or less, applicators must use ½ swath displacement upwind at the downwind edge of the field. When the windspeed is between 11-15 miles per hour, applicators must use ¾ swath displacement upwind at the downwind edge of the field.
- Do not apply during temperature inversions.

Airblast applications:

- Sprays must be directed into the canopy.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- Do not apply during temperature inversions.

Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Select nozzle and pressure that deliver medium or coarser droplets as indicated in nozzle manufacturers' catalogues and in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- Do not apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters. Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.
- IMPORTANCE OF DROPLET SIZE: An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.
 - Controlling Droplet Size - Ground Boom
 - Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
 - Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
 - Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.
 - Controlling Droplet Size – Aircraft
 - Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.
- BOOM HEIGHT – Ground Boom: For ground equipment, the boom should remain level with the crop and have minimal bounce.
- RELEASE HEIGHT – Aircraft: Higher release heights increase the potential for spray drift.
- SHIELDED SPRAYERS: Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.
- TEMPERATURE AND HUMIDITY: When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.
- TEMPERATURE INVERSIONS: Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.
- WIND: Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.
- Boomless Ground Applications: Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.
- Handheld Technology Applications: Take precautions to minimize spray drift.

The maximum application rate per year or crop cycle is not to exceed 0.33 pound per acre of 1-NAA equivalent.

The minimum interval between applications is to be no less than 5 days.

K-SALT FRUIT FIX 800 COMPATIBILITY WITH OTHER PRODUCTS

K-SALT FRUIT FIX 800 is compatible with a wide range of pesticides. The mixture must be field tested on a small scale to ensure the product performance and plant safety before large-scale field use. Observe all cautions and limitations on labeling of all products used in mixtures. Do not apply tank mix combination unless your previous experience indicates the mixture is effective and will not result in application problems or plant injury.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

CHEMICAL THINNING

Apples

Because of differences in varieties, climatic factors, tree vigor and fruit set, the application rate for thinning can range from 2 to 20 ppm and application timing can range from petal fall to 30 days after full bloom (0.6 fl. oz. of K-SALT FRUIT FIX 800 in 100 gallons of water will make 10 ppm 1-NAA spray). The higher concentrations must be limited to vigorous trees with high fruit set potential. Use the lower concentrations for weaker, less vigorous trees and less food reserves and lower fruit set potential. Use 2 to 5 ppm when a surfactant (wetting agent) is added. Use surfactant (wetting agent) according to manufacturer's recommendations, but not more than 1 pint per 100 gallons of spray. Mid-season to late blooming varieties such as Red Delicious, Golden Delicious, Fuji, Gala, Granny Smith and Romes must be treated 5 to 21 days after full bloom depending on locality and other factors. For concentrate spray, use K-SALT FRUIT FIX 800 at the rate of 0.5 to 1.0 fl. oz. per acre. Application in sufficient water for uniform and even distribution of the spray is imperative.

Thinning sprays are most effective when the king apple fruitlets are between 5 to 10 mm in diameter. During cool springs, delay application as necessary until the largest fruits are 7 mm to 10 mm in diameter and forecasted daytime temperatures are likely to rise above 60°F within the next five days. The preferred application temperature is between 70°F and 75°F. Applications are not advised below 60°F or above 80°F. When daytime temperatures exceed 85°F, over-thinning may occur. Good thinning can be obtained under the latter conditions with lower rates of chemicals.

Tank mix combinations of K-SALT FRUIT FIX 800 and Carbaryl (e.g. Sevin® 4F & Sevin® XLR Plus) have successfully thinned several early-maturing, heavy-setting varieties, as well as hard-to-thin apple varieties in accordance with the most restrictive of label limitations and precautions. Also a petal fall application of K-SALT FRUIT FIX 800 followed 7 to 10 days later by an application of K-SALT FRUIT FIX 800 + Carbaryl has improved thinning on hard-to-thin varieties such as Golden Delicious Fuji, Gala, and Rhode Island Greening. No label dosage rates must be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

IMPORTANT: K-SALT FRUIT FIX 800 must not be used on early season varieties (those that ripen before McIntosh) including Yellow Transparent, Early Williams Red and Oldenburg since severe injury or over-thinning may occur. Amid-Thin®W is specified for these early season varieties.

Pears - Bartlett, Bosc and Comice

Apply K-SALT FRUIT FIX 800 at a spray concentration of between 10 to 15 ppm for dilute spray in up to 400 gallons of water per acre or at a rate of 2 to 3 fl. oz. of product (13 to 19 grams of 1-NAA) per acre for concentrated spray (0.6 fl. oz. of K-SALT FRUIT FIX 800 in 100 gallons makes 10 ppm of spray solution). Apply K-SALT FRUIT FIX 800 once per year 15 to 28 days after full bloom with a surfactant (wetting agent). Apply in sufficient water for good, uniform and even distribution of the spray. K-SALT FRUIT FIX 800 may not adequately thin Bartlett pears in some seasons. The preferred application temperature is between 70°F and 75°F. Application is not advised below 60°F or above 80°F.

CONTROLS PRE-HARVEST DROP

Apples

Apply to apples 7 to 14 days prior to harvest. Separate sprays should be applied to early and late maturing varieties. Do not make more than two applications and do not delay picking beyond optimum maturity. Do not spray within two days of harvest. K-SALT FRUIT FIX 800 becomes effective in 3 to 4 days after application and controls drop for two weeks.

Ground Application: Use 4 to 8 fl. oz. of K-SALT FRUIT FIX 800 (25 to 50 grams of 1-NAA) per acre in sufficient water to ensure good coverage. Under normal conditions, the 8 fl. oz. rate per acre must be used.

Aerial Application: Use 4 to 8 fl. oz. in at least 5 gallons of water per acre. Aircraft must utilize spray equipment which gives thorough coverage to all portions of the trees. Apply only when weather conditions favor minimal spray drift. Use lower rates for normal foliage and fruiting and higher rates for dense foliage and heavy fruiting. Do not apply within two days of harvest.

Bartlett, Bosc, and D'Anjou Pears

Apply to pears 5 to 7 days prior to harvest. Separate sprays must be applied to early and late maturing varieties. Do not make more than two applications and do not delay picking beyond optimum maturity. Do not spray within two days of harvest. K-SALT FRUIT FIX 800 becomes effective in 3 to 4 days after application and controls drop for two weeks.

Ground Application: Use 2 to 4 fl. oz. of K-SALT FRUIT FIX 800 (12.5 to 25 grams of 1-NAA) per acre in sufficient water to ensure good coverage. Under normal conditions, the 4 fl. oz. rate per acre must be used. When Bartlett and Bosc are interplanted, spray only once per year. Use a maximum of 4 fl. oz. of product per acre per year on Bartlett to prevent premature ripening.

Aerial Application: Use 2 to 4 fl. oz. in at least 5 gallons of water per acre. Aircraft must utilize spray equipment which gives thorough coverage to all portions of the trees. Apply only when weather conditions favor minimal spray drift. Use lower rates for normal foliage and fruiting and higher rates for dense foliage and heavy fruiting. Do not apply within two days of harvest.

D' Anjou Pears

FOR USE ONLY WITHIN THE STATES OF OREGON AND WASHINGTON

K-Salt Fruit Fix 800 becomes effective in 3 to 4 days after application and controls drop for two (2) weeks. K-Salt Fruit Fix 800 may be applied by ground or air. For optimum results, apply 7 to 14 days before harvest. For maximum effectiveness, sprays must be applied when the temperature is 70°F or higher. The lower rate of K-Salt Fruit Fix 800 can be used under normal weather conditions. If warmer temperature conditions (i.e. 90-95°F) are expected to occur for a period greater than 3 consecutive days following the application, then the higher rate of K-Salt Fruit Fix 800 (up to 8 fl. oz. per acre) may be used. A repeat application of K-Salt Fruit Fix 800 is advised only if the efficacy of the first application of the product is declining, such as due to high temperatures. Consult your local Extension Pomologist as to time of application and rates to be used in your area for your specific stop drop situation.

1. Do not make more than two (2) applications of K-Salt Fruit Fix 800 at 4.0 fl. oz (25 grams 1-NAA/A) or one (1) application of K-Salt Fruit Fix 800 at 8.0 fl. oz. (50 grams 1-NAA/A) per stop drop season. Allow 7 to 14 days between applications.
2. Do not delay picking beyond optimum maturity.
3. PHI - Do not apply within two (2) days of harvest.
4. Do not apply this product through any type of irrigation system.

Ground Application: For dilute spray, mix 0.9 to 1.2 fl. oz. K-Salt Fruit Fix 800 per 100 gallons of water and apply 300 to 600 gallons of the spray mixture per acre of trees. For concentrate spray, use 4.0 to 8.0 fl. oz. K-Salt Fruit Fix 800 (25 to 50 grams of 1-NAA) per acre in sufficient water to ensure good coverage of trees.

Aerial Application: Use 4.0 to 8.0 fl. oz. K-Salt Fruit Fix 800 (25 to 50 grams of 1-NAA) in 5 to 10 gallons of water per acre. Aircraft must utilize spray equipment which gives thorough coverage to all portions of the trees. Apply only when weather conditions favor minimal spray drift.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store product in original container in a cool (not in direct sunlight), dry, locked place out of reach of children. Do not allow product to freeze.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning if available, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; and (b) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions. THIS WARRANTY DOES NOT EXTEND TO THE USE OF THIS PRODUCT CONTRARY TO LABEL INSTRUCTIONS, OR UNDER CONDITIONS NOT REASONABLY FORESEEABLE.

THERE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN TORT, CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

AMVAC offers this product, and Buyer accepts it, subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC.

© 2021 AMVAC Chemical Corporation. is a wholly owned subsidiary of American Vanguard Corporation. All Rights Reserved. AMERICAN VANGUARD, AMVAC, K-SALT, FRUIT FIX, and their respective logos are trademarks owned by AMVAC Chemical Corporation. Chemtrec is a service mark of the American Chemistry Council, Inc. Dreft is a trademark owned by Procter & Gamble.

AMVAC Chemical Corporation
4695 MacArthur Court, Suite 1200
Newport Beach, CA 92660 U.S.A.