

BEST AVAILABLE COPY

ACCEPTED



K-SALT FRUIT FIX 200

A HORMONE TYPE PRODUCT
FOR CONTROL OF PREHARVEST DROP
OF APPLES AND PEARS

ACTIVE INGREDIENT	
1 Naphthalene Acetic Acid, Potassium Salt	8.25%
INERT INGREDIENTS	91.75%
Total 100.00%	
Equivalent to 5.18% 1-Naphthaleneacetic Acid or 200 grams (0.44 lbs.) per gallon	

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by drinking 1 or 2 glasses of water and touching back of throat. Do not give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists. **IF ON SKIN:** Wash thoroughly with soap and water.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 5481 EPA Est. No. 5481 CA 1
NET CONTENTS 1 GALLON

AMVAC CHEMICAL CORPORATION
4100 EAST WASHINGTON BLVD
LOS ANGELES, CALIFORNIA 90023

Registered under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide described under EPA Reg. No. 5481 CA 1

HOW TO APPLY WITH GROUND EQUIPMENT

FOR APPLES, thoroughly mix 12 fluid ounces with 250 gallons of water (4.8 fluid ounces/100 gals. water) to prepare a 20 ppm spray. Thoroughly spray APPLE trees using up to 465 gallons per acre of this 20 ppm spray, which provides the equivalent of 35 grams of 1-Naphthaleneacetic acid per acre.

FOR PEARS, thoroughly mix 12 fluid ounces with 500 gallons of water (2.4 fluid ounces/100 gals. water) to prepare a 10 ppm spray. Thoroughly spray trees using up to 670 gallons of this 10 ppm spray dilution per acre, which provides the equivalent of 25 grams of 1-Naphthaleneacetic acid per acre. Apply to fruit 5 to 7 days before harvest. Do not make more than 2 applications.

HOW TO APPLY AS AN AERIAL APPLICATION

Aircraft should utilize spray equipment which gives thorough spray coverage to all portions of trees. Apply only when weather conditions favor minimal spray drift.

APPLES and D'ANJOU PEARS Use 22 fluid ounces in 5 gallons of water per acre. Do not apply within 2 days of harvest.

BARTLETT AND BOSCH PEARS Use 11 fluid ounces of this product in 5 gallons of water per acre. Do not apply within 2 day of harvest. When BARTLETT and BOSCH are interplanted, spray only once. Use a maximum of 11 fluid oz per acre on BARTLETT to prevent premature ripening.

FOR USE IN PRESERVING ORNAMENTAL HOLLY BOUGHS

To delay leaf drop of Holly boughs make a "dip" solution containing 8 to 16 fluid ounces in 100 gallon of water, (equals 33 to 66 ppm). Dip Holly as soon as possible after cutting. Allow excess solution to drain. Do not "soak" or leave boughs in dip solution. Replenish dip solution as needed however to maintain concentration. Replace with fresh solution every 3 to 4 days. Package while moist to keep boughs fresh until used. Maintain a 34 to 40 degree temperature in bough storage area.

NOTICE: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the manufacturer and/or seller and the buyer assumes the risk of any such use. 07-07-89

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. In case of eye contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse.

RE-ENTRY AND WORKER SAFETY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in treated areas or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written and oral warnings must include the following information: **CAUTION:** Area treated with K-SALT FRUIT FIX 200 on (Date) _____. Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, refer to the Precautionary Statements found on this product label.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters. Avoid drift of spray mist.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconsistent with its labeling. Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original container in a cool (not in direct sunlight), dry, locked place out of reach of children. Do not allow product to freeze. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then check for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned stay out of smoke.

K-SALT FRUIT FIX 200 controls preharvest drop of APPLES and PEARS, helping to a larger harvest and a reduction of losses from windfall and "knockdown".

GENERAL DIRECTIONS FOR USE ON APPLES AND PEARS

Spray trees thoroughly, wetting all tree stems and adjacent leaves. Apply 7 to 14 days before harvest on APPLES and 5 to 7 days before harvest on PEARS. K-SALT FRUIT FIX 200 becomes effective in 3 to 4 days after application and continues to drop for 3 to 4 weeks. Separate sprays should be applied to early and late maturing varieties. Do not make more than 2 applications and do not delay picking beyond optimum maturity. Do not spray within 2 days of harvest.

BEST AVAILABLE COPY