

PM: S 5481-441 P5 135

U.S. ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (WH-567)
WASHINGTON, D.C. 20460

EPA REGISTRATION NO.
5481-441

DATE OF ISSUANCE
1972-9-13

TERM OF ISSUANCE

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NAME OF PESTICIDE PRODUCT

PCNB 75 DG

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

AMVAC Chemical Corporation
4100 East Washington Blvd.
Los Angeles, CA 90023

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA Section 3(c)(5) or 4(a) when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment:

a. Add the phrase "EPA Registration No. 5481-441".

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

If these conditions are not complied with, this registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Robert J. Taylor
Product Manager (25)
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

RD:RTaylor:eja:9/9/92:3056800

BEST AVAILABLE COPY

PCNB 75 DG

A Soil Fungicide

For Turf and Ornamentals - For Field and Vegetable Crops

FOR PREVENTION AND CONTROL OF SOIL-BORNE DISEASES WHICH ATTACK TURFGRASS, NURSERY AND GREENHOUSE PLANTS, FIELD GROWN ORNAMENTALS, BEDDING AND FLOWERING PLANTS, HERBACEOUS AND WOODY ORNAMENTALS AND FOR FIELD AND VEGETABLE CROPS

ACTIVE INGREDIENT:
 Para-chloronitrobenzene(PCNB) - - 75.0%
 INERT INGREDIENTS: - - - - - 25.0%
 TOTAL 100.0%

EPA Registration No. 5461-

EPA Establishment No. 5461-CA-1

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENTS OF PRACTICAL TREATMENT
REPRODUCTION: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and pushing back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or comatose person. **RESPIRATORY:** Flush eyes with plenty of water. Get medical attention if irritation persists. **EYES:** Remove contaminated clothing and wash affected area with soap and warm water. Get medical attention if irritation persists. **SKIN:** Remove victim to fresh air and give artificial respiration if indicated.

PRECAUTIONARY STATEMENTS
HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed. Avoid inhaling dust. May cause skin irritation. Wash hands and face thoroughly after using and before eating or smoking. Avoid contact with skin. The following protective clothing is required during mixing/loading operations: (An approved respirator, for professional applicators only), gloves, long-sleeved shirt, long pants, socks and shoes. Thoroughly wash all contaminated clothing before reuse. Do not recontaminate food or feedstuffs.
ENVIRONMENTAL HAZARDS
 Do not contaminate water when disposing of equipment washwaters. This product is toxic to fish and aquatic organisms. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms.

In adjacent aquatic areas. Do not apply directly to water, to areas where surface water is present or to terrestrial areas before the mean high water mark. Apply this product only as specified on this label. Do not apply directly adjacent to potable water supplies.

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not plant root crops in PCNB treated soils within 12 months after treatment and banding applications unless PCNB is registered for use on these crops. Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL
 Do not contaminate water, food or feed by storage or disposal. **CONTAINER:** Store product in its original labeled container in a cool, dry, well-ventilated area out of reach of children. **RESIDUAL:** Fertilizer wastes are toxic. Improper disposal of this product, spray residue, or waste from the violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. (64-7008.PM) (7-15-82)

GENERAL INFORMATION
 This product is a wettable powder and is recommended for use in water. When mixing make sure that the water mixture is continuously agitated to give uniform fungicide coverage. PCNB 75 DG is a preventive fungicide recommended for control of certain soil-borne diseases as listed for turf and ornamental crops therefore best disease control results when applied as a preventive program. Carefully follow the recommended rates and the application remarks.

FOR TURFGRASS
 Use as a fungicide in the prevention and control of the listed soil-borne diseases which attack turfgrass on golf courses, sod farms, home lawns, institutional, ornamental, and recreational areas where fine turf is under a maintenance program.

DISEASES CONTROLLED
 Brown Patch (*Rhizoctonia solani*)
 Leaf Spot (*Helminthosporium spp.*)
 Dollar Spot (*Sclerotinia homocarpa*)
 Gray Snow Mold (*Typhula spp.*)
 Pink Snow Mold (*Fusarium nivale*)
 *Do not use in California.

For best results use PCNB 75 DG as a preventive application before diseases appear, or apply at the first sign of the infection. After application the treated areas should be watered to wash the fungicide down to the soil level. If heavy rainfall or flooding occurs after treatment, or if disease is severe or reappears, treat area again. Follow the rates as outlined below for the respective disease.

Disease/Fest	Rate/Gal. water per 1,000 sq. ft.	How and When to Apply
Brown Patch (<i>Rhizoctonia solani</i>) St. Augustine, Bermudagrass and similar warm season grasses	1 lb./10-15 Gals.	Apply once in Spring or Fall or when disease appears. If it is severe or reappears, repeat 3 to 4 weeks later.
Dollar Spot (<i>Sclerotinia homocarpa</i>) Bluegrass, Ryegrass, and similar cool season grasses	1 lb./40 Gals.	Treat as above.
Leaf Spot (<i>Helminthosporium spp.</i>) Dollar Spot (<i>Sclerotinia homocarpa</i>) *Do not use in California Cool & Warm season turf varieties	7-10 oz./5-10 Gals.	For preventive apply in Spring or Fall of each year. If disease is present apply at 10 oz. rate as soon as possible. If disease is severe or reappears repeat every 3 to 4 weeks later.
Gray Snow Mold (<i>Typhula spp.</i>) Pink Snow Mold (<i>Fusarium nivale</i>) Cool season turf varieties	8 oz./5-10 Gals.	1st treatment should be made prior to first snowfall.

Do not graze treated areas or lead fresh droppings to livestock. Under certain growing conditions, a temporary discoloration of the grass occasionally may occur. This causes no harm and will disappear in a short time.

NOTICE: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the manufacturer and/or seller, and buyer assumes risk of any such use.

ORNAMENTALS
 Mixtures of PCNB 75 DG and water must be mixed with the soil as a drench treatment for protection against the listed diseases on nursery, greenhouse, field grown and established plants. A preliminary trial is suggested first before a full treatment is given to any plant species not listed in these directions.

DISEASES CONTROLLED
Common Name
 Root and stem rot
 Botrytis blight
 Root rot
 Stem rot
 Root rot
 Stem rot
 Root rot
 Stem rot
 Root rot
 Stem rot
Fungal Name
 Rhizoctonia spp.
 Botrytis spp.
 Sclerotinia glauca
 Sclerotinia glauca
 Sclerotinia glauca
 Sclerotinia glauca
 Sclerotinia glauca
 Sclerotinia glauca
 Sclerotinia glauca
 Sclerotinia glauca
 Sclerotinia glauca
 Sclerotinia glauca

NURSERY & GREENHOUSE
 For *Rhizoctonia* control apply as a soil drench at time of seeding and/or transplanting.

Tropical Foliage Plants
 Agave, Aechmea, Dieffenbachia, Dracaena, Ficus, Philodendron, Gynura, Hoy, Peperomia, Petiole, Schefflera, Syngonium, Tradescantia and similar tropical foliage plants. Mix 4 to 8 oz. in 100 gallons water and apply to 400-800 sq. ft. of bed or bench area. Coverage: For soil depths of 4 inches or less, apply 100 gallons to 800 sq. ft. For soil depths of more than 4 inches apply 100 gallons to 400 sq. ft. One repeat application can be made 4 to 6 weeks later if necessary. Do not apply to Philodendron or Ficus since injury may occur.

Bedding and Flowering Plants
 Carnation, Chrysanthemum, Dahlia, Geranium, Marigold, Petunia, Polka Dot, Portulaca, Salvia, Vinca, Zinnia and similar plants. Use 4 oz. with 100 gallons of water and apply to 800 sq. ft. of flat or bench area. Apply at seeding or transplanting time.

Herbaceous and Woody Plants
 Herbaceous Plants: Kalimeris, Lily and similar plants. Woody Plants: Azalea, Boxwood, English Ivy, Holly, Juniper, Pine, Rhododendron and similar plants. Use 4 to 8 oz. with 100 gallons of water and apply to 400-800 sq. ft. of bed or bench area.

DIP TREATMENT for BULBS and DORMANT ROSES

Use before planting or storing. It is important to maintain continuous circulation of dip solution to assure uniform application of fungicide. Add 1% "Sticker" to solution. Use wire mesh for dipping and drain boards to return excess solution to dip tank. Dispose of unused dip solution according to "Storage and Disposal" and observe use of protective clothing (gloves etc.) as listed under "Hazardous to Humans"; both sections found on this label.

For control of Crown Rot and Black Rot on Iris, Narcissus, Hyacinth and Tulip dip bulbs for 5 minutes in a solution containing 2 lbs. PCNB 75 DG in 3.2 gallons of water.

For Root Rot (*Rhizoctonia*) of Easter Lilies dip in a solution containing 4 to 6 lbs. PCNB 75 DG in 100 gallons of water for a period of 5 to 15 minutes.

For Botrytis Blight on Roses: Dip or spray dormant plants before packaging or storage, with a solution of 2 lbs. PCNB 75 DG in 100 gals. of water.

FIELD GROWN ORNAMENTAL CROPS

HOW TO MIX AND APPLY
 Mix the following amounts of PCNB 75 DG in sufficient water and apply before planting to give uniform ground coverage. Incorporate the spray drench solution to depth of soil as indicated.
AS A PRE-PLANT BROADCAST TREATMENT
 For Rhizoctonia Root Rot on Easter Lilies, and for Crown Rot and Black Rot on Iris, Hyacinth, Narcissus and Tulip use 3 1/4 to 6 1/2 lbs. per 1,000 sq. ft. Thoroughly incorporate to a depth of 6 to 7 inches. Use the lower rate for light soils; the higher rate for heavy soils.

For Bulb Rot and Stem Rot (*Sclerotinia rot*) on Dutch Iris use 3 1/4 to 4 1/2 lbs./1,000 sq. ft. Apply prior to planting. Thoroughly incorporate solution to depth of 2 to 3 inches.

For Neck or Bulb Dry Rot (*Stemelia* or *Sclerotinia glauca*) on Gladiolus use 3 2/3 lbs./1,000 sq. ft. Apply prior to planting. Thoroughly incorporate to depth of 4 to 6 inches. Bulbs may be planted immediately after solution is incorporated.

For Stem Rot (*Rhizoctonia*) on California Larkspur, Snapdragon and Sweet Pea use 3 1/2 lbs. per 1,000 sq. ft. Apply prior to planting. Thoroughly incorporate to depth of 4 inches.

For Rhizoctonia Needle Blight on Southern Pine seedlings use 80 lbs. per acre. Mix and apply in 100 gallons of water to give uniform ground coverage. An additional 1/2 cup of water per acre should be applied by irrigation prior to planting. Seed may be planted immediately after irrigation or within one week of this date.

AS A BAND TREATMENT
 For Neck or Bulb Dry Rot (*Stemelia* or *Sclerotinia glauca*) on Gladiolus apply 5 1/8 lbs. per 1,000 linear feet of row as a 1/2 inch band centered on the row at planting time. Incorporate to depth of 4 inches. Bulbs may be planted immediately after incorporation.

For Bulb Rot and Stem Rot (*Sclerotinia rot*) on Dutch Iris use 1/2 lb. per 1,000 linear feet of row as a 1/2 inch band centered on the row at planting time. Follow with 3 additional applications at the same rate at 15 day intervals.

AS AN IN FURROW TREATMENT AT PLANTING TIME
 For control of Rhizoctonia Root Rot on Easter Lilies, and for Crown Rot and Black Rot on Iris, Hyacinth, Narcissus and Tulip mix 4 1/2 to 6 lbs. in 5 to 10 gallons of water and apply per 1,000 linear feet of row. Direct the spray into open furrow to thoroughly cover entire furrow and bulbs. Close furrow in normal manner. Use lower rate for lighter soils and higher rate for heavier soils. For Black Rot control, use high rate.

GROUND SPRAY for ESTABLISHED AZALEAS and CAMELIAS

For Azalea Leaf Blight (*Dothidea azaleae*) on Azaleas and Camellia Flower Blight on Camellias use 1 lb./150 sq. ft. Mix and apply in sufficient water (5 to 10 gallons per 100 sq. ft.) to give uniform coverage to soil beneath bushes and in the plant bed. Begin application prior to opening of buds and repeat every 3 to 4 weeks during bloom season.

For the use of PCNB 75 DG on the following Field and Vegetable crops refer to the PCNB 75 DG Supplemental/Additional Label Booklet FOR COTTON, PEANUTS, BEANS, BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, GARLIC, PEPPERS, TOMATOES

NET WEIGHT 80 POUNDS
AMVAC CHEMICAL CORPORATION
 4160 EAST WASHINGTON BLVD. • LOS ANGELES, CA 90023 U.S.A.

Supplemental / Additional Labeling for Field and Vegetable Crops

PCNB 75 DG

A Soil Fungicide

For Turf and Ornamentals - For Field and Vegetable Crops

FOR COTTON, PEANUTS, BEANS, BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, GARLIC, PEPPERS, TOMATOES

Refer to the label affixed to the container for instructions in using this product for Turfgrasses and Ornamentals

ACTIVE INGREDIENT: Pentachloronitrobenzene(PCNB) - - 75.0%
 INERT INGREDIENTS: - - - - - 25.0%
 TOTAL 100.0%

EPA Registration No. 5481-

EPA Establishment No. 5481-CA-1

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENTS OF PRACTICAL TREATMENT
IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving 1 or 2 glasses of water and flushing back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing patient. **IF EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists. **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and warm water. Get medical attention if irritation persists. **IF INHALED:** Remove victim to fresh air and give artificial respiration if indicated.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed. Avoid inhaling dust. May cause skin irritation. Wash hands and face thoroughly after using and before eating or smoking. Avoid contact with skin. The following protective clothing is required during manufacturing operations: (An approved respirator, for professional applicators only), gloves, long-sleeved shirt, long pants, socks and shoes. Thoroughly wash all contaminated clothing before reuse. Do not contaminate food or feedstuffs.

ENVIRONMENTAL HAZARDS
 Do not contaminate water when disposing of empty wet containers. This product is toxic to fish and aquatic organisms. DRI and runoff from treated areas may be hazardous to fish and aquatic organisms.

In untreated aquatic sites. Do not apply directly to water, to areas where surface water is present or to low tidal areas below the mean high water mark. Apply this product only as specified on this label. Do not apply directly adjacent to potable water supplies.

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not plant crop crops in PCNB treated fields within 12 months of broadcast and banding applications unless PCNB is registered for use on these crops. Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL
 Do not contaminate water, food or feed by storage or disposal. **RECYCLE:** Place product in its original labeled container in a cool, dry, well-ventilated place out of reach of children. **NON-RECYCLABLE:** Practice proper waste disposal. Improper disposal of empty containers, equipment, or waste is a violation of Federal law. If these wastes cannot be disposed of by the user according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA regional office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. (EPA 700P-PM-1 (07-16-82))

Use and Rates Chart for Field and Vegetable Crops

CROP	DISEASE	POUNDS/ACRE	OUNCE/1000 FEET OF ROW	RECOMMENDED APPLICATION
BROCCOLI BRUSSELS SPROUTS CABBAGE CAULIFLOWER	Clubroot (Fusarium wilt)	2 to 6 pounds Apply in 100 gallons of water. Use 1/2 to 1/4 part of this solution per plant.		Preventive Solution: The described disease should be based on the severity of the disease infection. Maintain solution to full growth in suspension.
		10 Apply in 25 gallons of water (40 inch row spacings - broccoli, Brussels sprouts and cabbage - 48 inch row spacings - cauliflower)	Brccoli, Brussels Sprouts, Cabbage 36.6 Cauliflower 44	Band Application: Spray in a 12 to 15 inch band centered on the row and incorporate to a depth of 4 to 6 inches immediately prior to planting.
		10 Apply in 10 gallons of water		Spot Application: Apply as a 6 inch horizontal spray prior to planting. Thoroughly incorporate to a depth of 4 to 6 inches using a disc or other suitable equipment.
Note: For alternate control of clubroot on broccoli, Brussels sprouts, cabbage and cauliflower. It is extremely important to thoroughly mix PCNB with the soil.				
Wilt Signs of Black Rot (Crown rot)		15 to 20 Apply in 50 gallons of water		Preventive Crown Application: Apply as a soil drench at the time of or immediately after seeding.
		For smaller areas use 1 level tablespoon per gallon of water in a soil drench at 50 to 80 lb of soil per acre. Apply with water can or similar equipment.		
Row Crotch Treatment		10 to 15 Apply in 25 gallons of water (40 inch row spacings - broccoli, Brussels sprouts and cabbage - 48 inch row spacings - cauliflower)	Brccoli, Brussels Sprouts, Cabbage 12.2 to 18.3 Cauliflower 14.6 to 22	Row Crotch Treatment: Spray on an 8 inch band centered on the row at the time of or immediately after seeding.
GARLIC	Wilt Rot (Sclerotium)	2 1/2 Apply in 100 gallons of water (24 inch row spacing)	20	In-Furrow Application: Apply 2 1/2 pounds per acre in a 12 inch band centered on the row at the time of or immediately after seeding. Attach a spray rig to the machine planting and spray @ 12 inch bands using 2 gallons per acre. Check band width to assure thoroughness. As it is difficult to spray the row, it may be necessary to spray the row with a good mechanical applicator in the spray tank to insure a uniform application.
		1 1/2 pounds per 1000 lb of seed cloves		Crown Treatment: Mix in suspension to apply directly to a uniform and spray for complete coverage of treated cloves. For optimum results add a commercial sticker to the suspension.
PEPPERS TOMATOES	Southern Blight (Sclerotium rot)	3 to 5 pounds in 100 gallons of water Use 1/2 part of the solution per plant.		Apply at the time of transplanting. The solution should be applied often to maintain a uniform suspension and prevent disease.
		10 Apply in 100 gallons of water (36 inch row spacings - pepper 72 inch row spacing. Red tomatoes 48 inch row spacing - standard levelness)	Peppers 11 Flat Tomatoes 22 Sliced Tomatoes 14.6	Apply as an in-furrow spray to the row at the time of or immediately after seeding. When cultivating, set planter depth to 2 to 3 inches to avoid getting arrested soil against stems of plants.
COTTON	Damping off (Fusarium rot)	1 1/2 to 2 1/2 (40 inch row spacing) Apply in at least 15 gals. of water	1.8 to 3.1	Apply spotted drench as a directed in-furrow spray to the seed and surrounding soil at planting time. Do not allow the loading or plowing of cotton clumps by harvest.
PEANUTS	Southern stem and root rot (Sclerotium rot)	3 1/2 inch row spacing 12 1/2 Apply in 8 to 10 gallons of water	14.7	PRE-PLANT SOIL DRENCH: Apply spotted drench as a spray in an 8 to 12 inch band centered on the row and incorporate to a depth of 2 to 4 inches. This drench may be made simultaneously with planting using suitable application equipment.
		4 1/2 Apply in 8 to 10 gallons of water	4.8	AT-PLANTING BAND: Apply spotted drench as a spray in a 12 inch band centered on the row at 48 to 60 to 72 inch spacing and plowing areas of the plant. If the row widths are not uniform at the same time, slash the row evenly between rows to depth of 2 to 3 inches of soil. Do not make any undisturbed soil into the row. If necessary remove fields areas.
				AT CULTIVATION: Apply spotted drench at cultivation. Do not apply more than 3 times per row season. Slash the row evenly between rows of 2 to 3 inches between rows on each side and adjust to 12 to 15 inch band of 12 inches centered on the row. Do not allow undisturbed soil on crown of plant.
NOTE: Do not exceed 10 lb actual PCNB per acre in any one season regardless of method of application. Do not feed treated wheat to livestock. Do not allow nitrogen down of peanuts.				
BEANS	Root and Stem Rot (Fusarium rot)	1 1/2 to 2 Apply in 8 to 10 gallons of water (36 inch row spacing - bush type beans - 42 inch row spacing - pole beans)	Bush 1.8 to 2.2 Pole 2.5 to 3.8	Apply as a directed spray in the seed furrow and to the covering soil at planting time.
BEANS (Snap & Dry)	Wilt Rot (Sclerotium rot)	1 1/2 Apply in 8 to 10 gallons of water	Bush 1 Pole 5.2	Apply as an 8 inch band centered on the row at or immediately after planting. If the disease pattern is severe, reapplication may be required at 2 to 3 week intervals using directed spray application with one to two gallons per row spaced at the end of the bands. Soil should remain undisturbed after application.
Do not apply after first harvest. Do not exceed 10 lb actual PCNB per acre in any one season regardless of method of application. Do not feed treated wheat to livestock.				
ROTATION RESTRICTION: Do not plant row crops in PCNB treated fields within 12 months of broadcast and banding applications unless PCNB is registered for use on these crops.				

BEST AVAILABLE COPY

NET WEIGHT 50 POUNDS
AMVAC CHEMICAL CORPORATION
 4100 EAST WASHINGTON BLVD. - LOS ANGELES, CA 90023 U.S.A.

ACCEPTED

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 5481-441

25