Pm 25 5y81-443

PCNB 2 FLOWABLE

TURF & ORNAMENTAL SOIL FUNGICIDE

FOR CONTROL OF SOIL-BORNE DISEASES ON TURFGRASS and ORNAMENTALS

ACTIVE INGREDIENT:

Pentachloronitrobenzene(PCNB)	21.6%
INERT INGREDIENTS	78.4%
TOTAL	100.0%

Contains 2 pounds of PCNB per Gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED Call a physician or Poison Control Center immediately Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger Do not induce vomiting or give anything by mouth to an unconscious or convulsing person IF IN EYES If eyes irritate, pour plenty of water Get medical attention if irritation persists IF ON SKIN If skin affected area with soap and water Get medical attention IF INHALED Remove victim to fresh air Apply artificial respiration if indicated Get medical attention

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION Harmful if swallowed Avoid inhaling spray mist This material may cause skin irritation Use with adequate ventilation Do not contaminate food or foodstuffs

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below If you want more options, follow the instructions for category (C) on an EPA chemical-resistance category selection chart Applicators and other handlers must wear long-sleeved shirt and long pants, chemical resistant gloves made of barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or Viton, shoes plus socks and for exposures in addition to dustmist filtering respirator (MSHA/NIOSH approval No prefix TC-21C) For exposures in excess of 5 hours, a mixer gear with either an approved vapor removing cartridge(s) with a prefilter approved for pesticides (MSHA/NIOSH approval No prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval No prefix TC-14G) Follow manufacturer's instructions for cleaning/maintaining PPE If no such instructions for washables, use detergent and hot water Keep and wash PPE separately from other laundry

User Safety Recommendations

Users should Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet Remove clothing immediately if pesticide gets inside Then wash thoroughly and put on clean clothing Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible, wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters This product is toxic to fish Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark Do not apply when weather conditions favor runoff or drift from the treated areas Apply this product only as specified on this label Do not apply directly to estuaries

CHEMICAL AND PHYSICAL HAZARDS

Do not allow product to freeze

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling Do not apply this product through any type of irrigation system Strict agitation may be required to mix the material Do not mix treated seed for feed, food or oil purposes Do not apply this product in a way that will contact workers or other persons, either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170 This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides It contains requirements for training, decontamination, notification, and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: long-sleeved shirt and long pants, chemical-resistant gloves such as Barrier Laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or Viton, and shoes plus socks

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. NO. 5481-443 NET CONTENTS 5 GALLONS EPA EST NO. 5481-CA-1

AMVAC CHEMICAL CORPORATION

4100 EAST WASHINGTON BLVD. LOS ANGELES, CA 90023 U.S.A.

FOR ORNAMENTALS

Mixtures of this product and water must be mixed with the soil as a drench treatment for protection against the listed diseases on nursery, greenhouse, field grown and established plants. A preliminary test is suggested first before a full treatment is given to any plant species not listed in these directions.

DISEASES CONTROLLED

Common Name	Fungus Name	Common Name	Fungus Name
Root and Stem rot	Rhizoctonia spp	Root blight	Diplodia spp
Damping off/seedling rot	Botrytis spp	Bulb and stem rot	Sclerotium spp
Black dry rot	Stromatinia gladioli	Flower blight	Botrytis cinerea
Bulb dry rot	Sclerotium gladioli	Crown rot	Phytophthora spp
Needle blight	Dothistroma pini	Black rot	Sphaeropsis sapinea

NURSERY & GREENHOUSE

For Rhizoctonia control apply as a soil drench at time of seeding and/or transplanting

Tropical Foliage Plants

Aloe, Aphelandra, Dieffenbachia, Dracena, Ficus, Fittonia, Gynura, Hoya, Peperomia, Pothos, Schefflera, Syngonium, Tradescantia and similar tropical foliage plants Mix 3/4 to 1 1/2 pints in 100 gallons water and apply to 400-800 sq ft of bed or bench area Coverage For soil depths of 4 inches or less, apply 100 gallons to 800 sq ft For soil depths of more than 4 inches apply 100 gallons to 400 sq ft One repeat application can be made 4 to 5 weeks later if necessary Do not apply to Philodendron or Pilea since injury may occur

Bedding and Flowering Plants

Carnation, Chrysanthemum, Dahlia, Geranium, Marigold, Petunia, Poinsettia, Portulaca, Salvia, Vinca, Zinnia and similar plants Use 3/4 pint with 100 gallons of water and apply to 800 sq ft of bed or bench area Apply at seeding or transplanting time

Herbaceous and Woody Plants

Herbaceous Plants, Kalanchoe, Lily and similar plants Woody Plants Azalea, Boxwood, English Ivy, Holly, Juniper, Privet, Rhododendron and similar plants Use 3/4 to 1 1/2 pints with 100 gallons of water and apply to 400-800 sq ft of bed or bench area

DIP TREATMENT for BULBS and DORMANT ROSES

Use before planting or storing It is important to maximize simultaneous circulation of dip solution to assure uniform application of fungicide Add 1% "Sactipe" to tankmix Utilize wire mesh for dipping and drain bucket to return excess solution to dip tank Dispose of unused dip solution according to "Storage and Disposal" and observe use of protective clothing as listed under "Hazards to Humans", both sections found on this label Wear chemical resistant gloves when dipping

For control of Crown Rot and Black Rot on Iris, Hyacinth, Narcissus and Tulips dip bulbs for 5 minutes in a solution containing 3/4 gallon of this product in 3.2 gallons of water

For Root Rot (Rhizoctonia) of Easter Lilies dip in a solution containing 1 1/2 to 2 1/4 gal's PCNB 2 FLOWABLE TURF & ORNAMENTAL in 100 gallons of water for a period of 5 to 15 minutes

For Botrytis Storage Rot on Roses, Dip or spray dormant canes before packaging or storage, with a solution of 1 gallon PCNB 2 FLOWABLE TURF & ORNAMENTAL in 100 gals of water

FIELD GROWN ORNAMENTAL CROPS

HOW TO MIX AND APPLY

Mix the following amounts of PCNB 2 FLOWABLE TURF & ORNAMENTAL in sufficient water and apply before planting to give uniform ground coverage. Incorporate the spray/drench solution to depth of soil as indicated

AS A PRE-PLANT BROADCAST TREATMENT

For Rhizoctonia Root Rot on Easter Lilies, and for Crown Rot and Black Rot on Iris, Hyacinth, Narcissus and Tulips use 1 1/5 to 2 1/3 gals. per 1,000 sq. ft. Thoroughly incorporate to a depth of 6 to 7 inches. Use the lower rate for light soils; the higher rate for heavy soils.

For Crown Rot and Stem Rot (Sclerotinia rolfsii) on Dutch Iris use 1 1/5 to 1 2/3 gals /1,000 sq ft Apply prior to planting Thoroughly incorporate solution to depth of 2 to 3 inches

For Neck or Bulb Dry Rot (Stromatinia or Sclerotinia gladioli) on Gladiolus use 1 1/3 gal /1,000 sq.ft. Apply prior to planting. Thoroughly incorporate to depth of 4 to 6 inches. Bulbs may be planted immediately after solution is incorporated

For Stem Rot (Rhizoctonia) on Calendula, Larkspur, Snapdragon and Sweet Pea use 1 1/3 gal per 1,800 sq ft. Apply prior to planting. Thoroughly incorporate to depth of 4 inches

For Rhizoctonia Needle Blight on Southern Pine Seedlings mix 16 3/4 gals. per acre rate and apply in 100 gallons of water to give uniform ground coverage. An additional 1 1/2 inch of water per acre should be applied by irrigation prior to planting. Seed may be planted immediately after irrigation or within one week of this date

AS A BAND TREATMENT

For Neck or Bulb Dry Rot (Stromatinia or Sclerotinia gladioli) on Gladiolus apply 2 1/4 gallons per 1,000 linear feet of row as a 12 inch band centered on the row at planting time Incorporate to depth of 4 inches Bulbs may be planted immediately after incorporation

For Bulb Rot and Stem Rot (Sclerotinia rolfsii) on Dutch Iris use 1 ct. per 1,000 linear feet of row as a 12 inch band centered on the row at planting time Follow with 3 additional applications at the same rate at 15 day intervals

AS AN IN FURROW TREATMENT AT PLANTING TIME

For control of Rhizoctonia Root Rot on Easter Lilies, and for Crown Rot and Black Rot on Iris, Hyacinth, Narcissus and Tulips, use 1 3/4 gal. to 2 3/4 gals. in 5 to 10 gallons of water and apply per 1,000 linear feet of row Direct the spray into open furrows to thoroughly cover entire furrow and bulbs Cover furrow in normal manner Use lower rate for lighter soils and higher rate for heavier soils For Bulb Rot control, use high rate

SPRAY for ESTABLISHED AZALEAS & CAMELLIAS

For Azalea Petal Blight (Conidia stage) on Azaleas and Camellia Flower Blight on Camellia use 3 pints /100 sq ft Mix and apply in sufficient water (5 to 10 gallons per 100 sq ft) to give uniform coverage to soil beneath branches and to thin plant bed Begin application prior to opening of buds and repeat every 3 to 4 weeks during bloom season.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal STORAGE Store product in its original container in a cool, dry, locked place out of reach of children Protect from temperatures below 20 °F PESTICIDE DISPOSAL Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility CONTAINER DISPOSAL Triple rinse (or equivalent) Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved procedures by State and local authorities

FOR TURFGRASS

Use as a fungicide in the prevention and control of the listed soil-borne diseases which attack turfgrass on golf courses, sod farms, home lawns, institutional, ornamental, and recreational grass. Adjust rate for turf in under a maintenance program Do not use for Turf Leaf or Dollar Spot or Cottonwood

DISEASES CONTROLLED

Brown Patch (Rhizoctonia solani)	Leaf Spot (Helminthosporium (spp.)*
Dollar Spot (Sclerotinia homeocarpa)*	Gray Snow Mold (Typhula spp.)
Pink Snow Mold (Fusarium nivale)	*Do not use in California

For best results use as a preventive application before diseases appear, or apply at the first sign of the infection If heavy rainfall or flooding occurs after treatment, or if diseases is severe or reappears, treat area again. Follow the rates and comments on listed below for control of turfgrass diseases NOTE Apply the high rate when disease is severe and low rate when disease pressure is light Treated areas should be watered lightly as soon as possible after the application to wash the fungicide down to the soil level

Disease/Host	Amt./Gals. water per 1,000 Sq Ft	How and When to Apply
Brown Patch (Rhizoctonia solani) St. Augustine Bermuda & similar warm season grasses Dichondra	32-48 fl oz /1 to 5 Gals. of water	Apply once in spring or fall or when disease appears If it reappears, re-treat 3 to 4 weeks later
	32-48 fl oz /5 to 10 Gals. of water	Treat as above
Bluegrass Rye Fescue Bent & similar cool season grasses	16-32 fl oz /1 to 5 Gals. of water	At only at 1st sign of disease Repeat every 7-10 days during warm damp weather
Leaf Spot (Helminthosporium spp.) Dollar Spot * (Sclerotinia homeocarpa) *Do not use in California	22-32 fl oz /1 to 5 Gals. of water	For preventative apply in spring or fall of each year If disease is present apply at 32 fl. oz. rate as soon as possible If disease is severe or reappears repeat in 3 to 4 weeks later
Gray Snow Mold (Typhula spp.) Pink Snow Mold (Fusarium nivale)	24-32 fl oz /1 to 5 Gals. of water	1st treatment should be made prior to first snow fall

Do not graze treated areas or feed grass clippings to livestock Under certain growing conditions a temporary discoloration of the grass occasionally may occur This appears no harm and will disappear in a short time

IMPORTANT: PLEASE READ BEFORE USE

By using this product, the user accepts the following LIMITED WARRANTY

AMVAC warrants that (a) this product conforms to the chemical description on its label (b) this product is reasonably fit for the purposes stated on its label, subject to the inherent risks referred to above, when used in accordance with the directions and (c) that the directions, warnings, cautions, and other statements on this label are based upon responsible analysis, evaluations of reasonable tests of effectiveness, of toxicity to humans, domestic animals, of agricultural crops and plants, and upon records of field experience Testing has not been performed on all varieties of listed crops and plants, in all states, or under all applications, weather and pest conditions

There are no warranties, representations either than those set forth herein Amvac neither makes nor intends, nor does it authorize any agent or representative to make, any other warranty, express or implied Amvac expressly disclaims and does not intend, representations of merchantability Because for particular purpose or any other warranty of quality or performance

This warranty does not extend to, and the user shall be solely responsible for, any loss or damage which results from the use of this product in any manner which is inconsistent with this label's directions, warnings or cautions User's exclusive remedy and Amvac's exclusive obligation liability for any other loss, damage, or injury resulting from the use or handling of this product, whether or not based on contract, negligence, strict liability in tort, or other legal shall be limited, at Amvac's option, to replacement of or repayment of the purchase price for the quantity of product with respect to which damages are claimed In no event shall Amvac or Seller be liable for special, indirect, or consequential damages resulting from the use or handling of this product

(443-amvl Pkg#08-88-85)

FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY

TRANSPORTATION CHEMTREC 1-800-424-9300

MEDICAL HAZARD INFORMATION SERVICES (H.I.S.) 1-800-228-5635 Ext. 188

JUN 28 1995

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 5481-443

BEST COPY AVAILABLE