

TERM OF ISSUANCE

NAME OF PESTICIDE PRODUCT

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION

(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

PCNB 20% WDG SOIL FUNGICIDE

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Reg # 5481-450
Amvac Chemical Corporation
4100 East Washington Blvd.
Los Angeles, CA 90023

PM-25

103
7

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA Section 3(c)(5) and sec. 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment:
 - a. Add the phrase "EPA Registration No. 5481-450"
3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, this registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

ATTACHMENT IS APPLICABLE

BEST AVAILABLE COPY

SIGNATURE OF APPROVING OFFICIAL

DATE

OCT 12 1994

293

PCNB

20% WDG SOIL FUNGICIDE

BEST AVAILABLE COPY PEANUTS

FOR CONTROL OF SOIL-BORNE DISEASES AS LISTED ON COTTON - BEANS - CRUCIFERS PEANUTS - POTATOES LAWNS and TURF ORNAMENTAL PLANTINGS

ACTIVE INGREDIENT:
 Pentachloronitrobenzene (PCNB) 20.0%
 INERT INGREDIENTS: 80.0%
 Total 100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush with plenty of water. Get medical attention immediately. **IF ON SKIN:** Wash affected area with soap & water. **IF INHALED:** Remove victim to fresh air. Apply artificial respiration if needed. Get medical attention.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhaling dust. This product may cause skin irritation. Do not contaminate feed or foodstuffs.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants; waterproof gloves; and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Cover or incorporate granules. Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply directly adjacent to potable water supplies.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not plant root crops in PCNB treated fields within 12 months of broadcast and banding applications unless PCNB is registered for use on those crops. **IMPORTANT:** Where minimum and maximum rates are given, use the lower rate for crops in sandy soils and when conditions are unfavorable for the development of disease. Use the maximum rate for heavy organic soils and when conditions are favorable for the disease such as moist conditions with warm days and cool nights.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves and shoes plus socks.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

COTTON

For the control of Damping-off, Seed and Seedling Rots — APPLY AS AN IN-FURROW SOIL TREATMENT for Rhizoctonia solani. Apply 5 to 10 lbs. per acre as an in-furrow soil application which is equivalent to 1 to 2 lbs. per 1300 linear feet of row based on a 40 inch spaced row. By using suitable application equipment, apply over the seed and immediately cover with the seed at planting time. Do not allow livestock to graze or feed on cotton foliage in treated fields.

BEANS

Root and Stem Rot - Rhizoctonia solani: APPLY AS AN IN-FURROW SOIL TREATMENT - Apply 5 to 7 1/2 lbs. per 14,500 linear feet of row for Bush Beans or per 8,340 sq.ft. of row for Pole Beans. Apply over the seed at planting time in the furrow and immediately cover with soil utilizing appropriate application equipment. Do not feed treated vines to livestock.

For Southern Blight (Sclerotium rolfsii) - AS A SURFACE BAND TREATMENT - Apply 50 lb per 12,400 linear feet of row, in an 8 to 12 inch wide band, centered on the row, at peggin time. - AS A PRE-PLANT SOIL TREATMENT apply 50 lbs./12,400 linear feet of row, in an 8 to 12 inch band, centered on the row. Mix to a depth of 2 to 4 inches. This should be done simultaneously with planting, using appropriate equipment. AS A SOIL MIX DURING CULTIVATION it may be applied in three split applications when cultivating the crop at the rate 16 1/2 lbs. per 12,400 sq.ft. per treatment by attaching two delivery tubes per row ahead inside cultivator sweeps (one on each side) and adjusted so as to treat a total band of 12 inches wide, centered on the row. Do not pile untreated soil on crown of plant. NOTE: Do not exceed 10 lbs. actual PCNB per acre in any one season, regardless of the application methods used. Do not feed treated vines to livestock. Do not allow hogging down of peanuts.

POTATOES

Rhizoctonia solani - AS AN OVER-ALL BROADCAST TREATMENT - Apply 90 to 125 lbs. p acre over the soil surface prior to planting and thoroughly disc to a depth of 4 to 6 inches cross discing if necessary. - AS AN IN-FURROW TREATMENT - Apply 50 lbs./12,400 linear feet of a 36 inch row by thoroughly working granules to a depth of 4 to 6 inches and into the tube forming area of the soil. It is best to apply when the soil is slightly moist.

IN LAWNS AND ORNAMENTAL TURF

FOR BROWN PATCH

(Rhizoctonia solani)

SOUTHERN GRASSES - For St. Augustine, Bermuda, etc. and Dichondra - Apply 3 3/4 lb / 1,000 sq.ft. in the Fall or the Spring at the first sign of infection. Lightly sprinkle the area 1 wash the granules to soil level. An additional application in 3 to 4 weeks may be needed if continued control if heavy rainfall or flooding occurs after this treatment.

NORTHERN GRASSES - For Bluegrass, Fescue, Rye, Bent, etc. - Apply 1 to 1 1/4 lbs. p 1,000 sq.ft. when disease first appears and repeat at 7 to 10 day intervals during warm, damp weather. Lightly sprinkle the area to wash granules to soil level. During lush growing period a temporary discoloration of the grass may occur, however, this is not harmful and disappears in a few days.

HELMINTHOSPORIUM LEAF SPOT

Apply at the rate of 2 1/2 to 3 3/4 lbs./1,000 sq. ft. Preventive applications should be made in the spring or fall of each year. For control of existing infections, apply at the higher rate as soon as symptoms are first noted. Following applications, the treated area should be light watered to insure movement through the grass to soil level. If the treated area is subjected to unusually heavy rainfall, or flooded, or if the disease is severe or reappears, the area should be retreated in 3 to 4 weeks.

SNOW MOLD

(Typhula incarnata, Fusarium nivale) Apply at the rate of 2 1/2 to 3 3/4 lbs./1,000 sq. ft. using a granular lawn spreader or comparable equipment. Application should be made prior to first snow fall. Do not graze treated areas or feed clippings to livestock.

ORNAMENTAL PLANTS BEDDING PLANTS

For African Violets, Begonias, Carnations, Chrysanthemums, Poinsettias, Snapdragons and similar bedding plants. Treat bench soil prior to planting with 7 1/2 lbs. granules / 1,000 sq.ft. for control of Rhizoctonia, Stem rot. Thoroughly work into the top 2 inches of soil.

BULBS

For Tulips, Narcissus, Hyacinth, and Bulbous Iris in production - Apply 500 to 1,000 lbs. p acre to the soil as a pre-plant broadcast application for control of Crown rot and Black rot (Equivalent to 1.15 to 2.25 lbs./100 sq.ft.). Mix with the top 6 to 7 inches of soil with appropriate equipment. Use the higher rate in heavy organic soils and especially for the control of Black Rot.

CALENDULAS, LARKSPUR, SNAPDRAGON

For control of Sclerotinia and Stem rot. Apply 500 lbs./acre as a broadcast application two weeks before planting; (Equiv. to 1.15 lbs. per 100. sq.ft.). Mix with soil to a depth of 4 inches.

DUTCH IRIS

For Bulb and Stem Rot caused by Sclerotium rolfsii disease. Make a PRE-PLANT IN BROADCAST SOIL APPLICATION of 500 to 1,000 lbs. per acre which is equivalent to 1.15 to 2.25 lbs./100 sq.ft. Mix the granules with the soil to a depth of 2 to 3 inches. FOR A ROW BAND TREATMENT apply 60 lbs. per 14,500 sq.ft. of linear row in a 12 inch band centered on the row, at planting time. Follow this treatment with 3 additional band applications at 15 d intervals.

EASTER LILY

(Lilium Longiflorum Eximium): For control of Rhizoctonia Root Rot. Apply AS A BROADCAST SOIL TREATMENT at the rate of 500 lbs. on light sandy soils and 1,000 lbs. per acre on heavy organic soils. Follow with a good double discing to mix granules with the soil to the depth 6 to 7 inches.

GLADIOLUS

For Neck or Bulb Dry Rot caused by Stromatinia or Sclerotium gladioli. Apply 600 lbs. per acre (equivalent to 1 1/3 lbs. per 100 sq.ft.) AS A BROADCAST SOIL TREATMENT. Thorough mix with soil to depth of 4 to 6 inches. Corms may be planted immediately.

IMPORTANT: PLEASE READ BEFORE USE

By using this product, the user accepts the following **LIMITED WARRANTY:** AMVAC warrants that (a) this product conforms to the chemical description on its label; (b) this product is reasonably fit for the purposes stated on its label, subject to the inherent risk referred to herein, when used in accordance with its directions; and (c) that the direction warnings, cautions, and other statements on this label are based upon responsible expert evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and plant of residues on food crops, and upon reports of field experience. Testing has not been performed on all varieties of food crops, and plants, in all states, or under all application weather and crop conditions. There are no express warranties other than those set forth herein. Amvac neither makes nor intends, nor does it authorize any agent or representative to make, any other warranty express

ADDITIONAL
Web COMMENTS
In EPA Letter Dated:

001 | 2 | 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

NTS: 80.0%
Total 100.0%

OF REACH OF CHILDREN CAUTION

NTS OF PRACTICAL TREATMENT
If a person or Petal Control Center immediately. Induce vomiting by
water and touching back of throat with finger. Do not induce
vomiting if the person is unconscious or convulsing person. **IF IN EYES:** Flush
with water immediately. **IF ON SKIN:** Wash affected area
with soap and water. Remove victim to fresh air. Apply artificial respiration if
needed.

CAUTIONARY STATEMENTS FOR HUMANS AND DOMESTIC ANIMALS

Avoid inhaling dust. This product may cause skin irritation.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants; waterproof
gloves; and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining
PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE
separately from other laundry.

User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the
toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on
clean clothing. Remove PPE immediately after handling this product. Wash the outside of
gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Cover or incorporate granules. Do not
contaminate water when disposing of equipment washwaters. Do not apply directly to water,
to areas where surface water is present, or to intertidal areas below the mean high water mark.
Do not apply directly adjacent to potable water supplies.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do
not apply this product in a way that will contact workers or other persons, either directly or
through drift. Only protected handlers may be in the area during application. For any
requirements specific to your State or Tribe, consult the agency responsible for pesticide
regulation. Do not plant root crops in PCNB treated fields within 12 months of broadcast and
banding applications unless PCNB is registered for use on those crops.
IMPORTANT: Where minimum and maximum rates are given, use the lower rate for crops in
sandy soils and when conditions are unfavorable for the development of disease. Use the
maximum rate for heavy organic soils and when conditions are favorable for the disease such
as moist conditions with warm days and cool nights.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard,
40 CFR part 170. This Standard contains requirements for the protection of agricultural
workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesti-
cides. It contains requirements for training, decontamination, notification, and emergency
assistance. It also contains specific instructions and exceptions pertaining to the statements
on this label about personal protective equipment (PPE), and restricted-entry interval. The
requirements in this box only apply to uses of this product that are covered by the Worker
Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-
entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted
under the Worker Protection Standard and that involves contact with anything that has been
treated, such as plants, soil, or water, is: coveralls, waterproof gloves and shoes plus socks.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store product in its original container in a cool, dry, locked place out of reach of
children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be
disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:**
Completely empty bag into application equipment. Then dispose of empty bag in a sanitary
landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay
out of smoke.

COTTON

For the control of Damping-off, Seed and Seedling Rots — APPLY AS AN IN-FURROW SOIL
TREATMENT for *Rhizoctonia solani*. Apply 5 to 10 lbs. per acre as an in-furrow soil application
which is equivalent to 1 to 2 lbs. per 1300 linear feet of row based on a 40 inch spaced row.
By using suitable application equipment, apply over the seed and immediately cover with the
seed at planting time. Do not allow livestock to graze or feed on cotton foliage in treated fields.

BEANS

Root and Stem Rot - *Rhizoctonia solani*: APPLY AS AN IN-FURROW SOIL TREATMENT -
Apply 5 to 7 1/2 lbs. per 14,500 linear feet of row for Bush Beans or per 8,340 sq. ft. of row for
Pole Beans. Apply over the seed at planting time in the furrow and immediately cover with soil
utilizing appropriate application equipment. Do not feed treated vines to livestock.

CRUCIFERS

CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, AND BROCCOLI

Clubroot (*Plasmodiophora brassicae*): APPLY AS A PRE-TRANSPLANTING ROW TREAT-
MENT - Apply 100 lbs. per 13,100 feet of row for Cabbage, Broccoli and Brussels Sprouts or
to 10,900 linear feet of row for Cauliflower, in a 12 to 15 inch band. Apply and rototill the
granules to a depth of 4 to 6 inches just before transplanting.

TO APPLY AS A BROADCAST TREATMENT - Apply 150 lbs. per acre to the soil surface prior
to transplanting, then disc the granules into the soil 4 to 6 inches deep. Thorough mixing with
the soil will give best disease control. **CAULIFLOWER:** For control of Wire stem or Black Rot
(*Corticium solani*) APPLY AS A BAND ROW TREATMENT - Apply 55 to 125 lbs. per 10,900
linear feet of row in an 8 inch wide band centered on the row, immediately before planting. For
small areas use 6 lbs. / 1000 sq. ft. Mix granules into soil before planting to a depth of 2 inches.

Rhizoctonia solani - AS AN OVER-ALL BROADCAST TREATMENT - Apply 90 to 125 lbs. p
acre over the soil surface prior to planting and thoroughly disc to a depth of 4 to 6 inches cross
discing if necessary. - AS AN IN-FURROW TREATMENT - Apply 50 lbs./12,400 linear fe
of a 36 inch row by thoroughly working granules to a depth of 4 to 6 inches and into the tube
forming area of the soil. It is best to apply when the soil is slightly moist.

IN LAWN AND ORNAMENTAL TURF

FOR BROWN PATCH

(*Rhizoctonia solani*)

SOUTHERN GRASSES - For St. Augustine, Bermuda, etc. and Dichondra - Apply 3 3/4 lb
/ 1,000 sq. ft. in the Fall or the Spring at the first sign of infection. Lightly sprinkle the area i
wash the granules to soil level. An additional application in 3 to 4 weeks may be needed i
continued control if heavy rainfall or flooding occurs after this treatment.

NORTHERN GRASSES - For Bluegrass, Fescue, Rye, Bent, etc. - Apply 1 to 1 1/4 lbs. p
1,000 sq. ft. when disease first appears and repeat at 7 to 10 day intervals during warm, dem
weather. Lightly sprinkle the area to wash granules to soil level. During lush growing period
a temporary discoloration of the grass may occur, however, this is not harmful and w
disappear in a few days.

HELMINTHOSPORIUM LEAF SPOT

Apply at the rate of 2 1/2 to 3 3/4 lbs./1,000 sq. ft. Preventive applications should be mad
in the spring or fall of each year. For control of existing infections, apply at the higher rate a
soon as symptoms are first noted. Following applications, the treated area should be light
watered to insure movement through the grass to soil level. If the treated area is subjected t
unusually heavy rainfall, or flooded, or if the disease is severe or reappears, the area shoul
be retreated in 3 to 4 weeks.

SNOW MOLD

(*Typhula incarnata*, *Fusarium nivale*): Apply at the rate of 2 1/2 to 3 3/4 lbs./1,000 sq. ft. usin
a granular lawn spreader or comparable equipment. Application should be made prior to fir
snow fall. Do not graze treated areas or feed clippings to livestock

ORNAMENTAL PLANTS BEDDING PLANTS

For African Violets, Begonias, Carnations, Chrysanthemums, Polkaetias, Snapdragons an
similar bedding plants. Treat bench soil prior to planting with 7 1/2 lbs. granules / 1,000 sq. ft
for control of *Rhizoctonia*, Stem rot. Thoroughly work into the top 2 inches of soil.

BULBS

For Tulips, Narcissus, Hyacinth, and Bulbous Iris in production - Apply 500 to 1,000 lbs. p
acre to the soil as a pre-plant broadcast application for control of Crown rot and Black r
(Equivalent to 1.15 to 2.25 lbs./100 sq. ft.). Mix with the top 6 to 7 inches of soil with appropriat
equipment. Use the higher rate in heavy organic soils and especially for the control of Bla
Rot.

CALENDULAS, LARKSPUR, SNAPDRAGON

For control of Sclerotinia and Stem rot. Apply 500 lbs./acre as a broadcast application
before planting; (Equiv. to 1.15 lbs. per 100. sq. ft.). Mix with soil to a depth of 4 inches.

DUTCH IRIS

For Bulb and Stem Rot caused by *Sclerotium rot* disease. Make a PRE-PLANTING
BROADCAST SOIL APPLICATION of 500 to 1,000 lbs. per acre which is equivalent to 1.1
to 2.25 lbs. /100 sq. ft. Mix the granules with the soil to a depth of 3 to 4 inches. FOPAR O
BAND TREATMENT apply 60 lbs. per 14,500 sq. ft. of linear row in a 12 inch band center
on the row, at planting time. Follow this treatment with 3 additional band applications at 15 d
intervals.

EASTER LILY

(*Lilium Longiflorum Eximium*): For control of *Rhizoctonia* Root Rot - Apply AS A BROADCAST
SOIL TREATMENT at the rate of 500 lbs. on light sandy soils and 1,000 lbs. per acre on hea
organic soils. Follow with a good double disking to mix granules with the soil to the depth
6 to 7 inches.

GLADIOLUS

For Neck or Bulb Dry Rot caused by *Stromelia* or *Sclerotium gladioli* - Apply 600 lbs. per acr
(equivalent to 1 1/3 lbs. per 100 sq. ft.) AS A BROADCAST SOIL TREATMENT. Thorough
mix with soil to depth of 4 to 6 inches. Corms may be planted immediately.

IMPORTANT: PLEASE READ BEFORE USE

By using this product, the user accepts the following: **LIMITED WARRANTY:**
AMVAC warrants that (a) this product conforms to the chemical description on its label; (b)
this product is reasonably fit for the purposes stated on its label, subject to the inherent risk
referred to herein, when used in accordance with its directions; and (c) that the direction
warnings, cautions, and other statements on this label are based upon responsible expert
evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and plant
of residues on food crops, and upon reports of field experience. Testing has not been
performed on all varieties of food crops, and plants, in all states, or under all applicatio
weather and crop conditions.
There are no express warranties other than those set forth herein. Amvac neither makes n
intends, nor does it authorize any agent or representative to make, any other warranty, expres
or implied. Amvac expressly excludes and disclaims all implied warranties of merchantabil
fitness for particular purpose, or any other warranty of quality or performance.

This warranty does not extend to, and the user shall be solely responsible for, any loss
damage which results from the use of this product in any manner which is inconsistent with
this label's directions, warnings or cautions. User's exclusive remedy and Amvac's or seller's
exclusive liability for any claim, loss, damage, or injury resulting from the use or handling
this product, whether or not based in contract, negligence, strict liability in tort, or otherwise
shall be limited, at Amvac's option, to replacement of, or repayment of the purchase price to
the quantity of product with respect to which damages are claimed. In no event shall Amvac
or Seller be liable for special, indirect, or consequential damages resulting from the use
handling of this product. (PCNB20DG.PMS/09-22-94)

FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY
TRANSPORTATION: CHEMTREC 1-800-424-9300
MEDICAL: HAZARD INFORMATION SERVICES (H.I.S.) 1-800-228-5635 Ext. 169

EPA REG. NO 5481-

EPA EST. 5481-CA-1

BEST AVAILABLE COPY
AMVAC CHEMICAL CORPORATION

4100 EAST WASHINGTON BLVD. - LOS ANGELES, CA 90023 U.S.A.