

NOV 2 - 1994

Mr. H.R. McLane
H. R. McLane, Inc.
7210 Red Road, Suite 206
Miami, FL 33143

Dear Mr. McLane:

Subject: Final Printed Labels
PCNB 20% WDG SOIL FUNGICIDE
EPA Reg. No. 5481-450
Your submission dated Oct. 4, 1994

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable and a stamped copy is enclosed for your records.

Sincerely yours,

Robert J. Taylor
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENTS OF PRACTICAL TREATMENT

ALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention immediately. **IF ON SKIN:** Wash affected area with soap & water. **IF INHALED:** Remove victim to fresh air. Apply artificial respiration if needed. Get medical attention.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhaling dust. This product may cause skin irritation. Do not contaminate feed or foodstuffs.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants; waterproof gloves; and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Cover or incorporate granules. Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply directly adjacent to potable water supplies.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not plant root crops in PCNB treated fields within 12 months of broadcast and banding applications unless PCNB is registered for use on those crops.

IMPORTANT: Where minimum and maximum rates are given, use the lower rate for crops in sandy soils and when conditions are unfavorable for the development of disease. Use the maximum rate for heavy organic soils and when conditions are favorable for the disease such as moist conditions with warm days and cool nights.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves and shoes plus socks.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

COTTON

For the control of Damping-off, Seed and Seedling Rots — APPLY AS AN IN-FURROW SOIL TREATMENT for Rhizoctonia solani. Apply 5 to 10 lbs. per acre as an in-furrow soil application which is equivalent to 1 to 2 lbs. per 1300 linear feet of row based on a 40 inch spaced row. By using suitable application equipment, apply over the seed and immediately cover with the seed at planting time. Do not allow livestock to graze or feed on cotton foliage in treated fields.

BEANS

Root and Stem Rot - Rhizoctonia solani: APPLY AS AN IN-FURROW SOIL TREATMENT - Apply 5 to 7 1/2 lbs. per 14,500 linear feet of row for Bush Beans or per 8,340 sq.ft. of row for Pole Beans. Apply over the seed at planting time in the furrow and immediately cover with soil utilizing appropriate application equipment. Do not feed treated vines to livestock.

CRUCIFERS

CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, AND BROCCOLI

Clubroot (Plasmodiophora brassicae): APPLY AS A PRE-TRANSPLANTING ROW TREATMENT - Apply 100 lbs. per 13,100 feet of row for Cabbage, Broccoli and Brussels Sprouts or to 10,900 linear feet of row for Cauliflower, in a 12 to 15 inch band. Apply and rebill the granules to a depth of 4 to 6 inches just before transplanting.

TO APPLY AS A BROADCAST TREATMENT - Apply 150 lbs. per acre to the soil surface prior to transplanting, then disc the granules into the soil 4 to 6 inches deep. Thorough mixing with the soil will give best disease control. **CAULIFLOWER:** For control of Wire Stem or Black Rot (Corticium solani) APPLY AS A BAND ROW TREATMENT - Apply 55 to 125 lbs. per 10,900 linear feet of row in an 8 inch wide band centered on the row, immediately before planting. For small areas use 6 lbs. / 1000 sq. ft. Mix granules into soil before planting to a depth of 2 inches.

Rhizoctonia solani - ACN-11
To control the soil-borne fungus causing damping-off, apply 10 lbs. per acre as a pre-planting soil treatment. Apply 10 lbs. per acre as a pre-planting soil treatment in a 35 inch row by thoroughly working granules to a depth of 4 to 6 inches and into the tuber-forming area of the soil. It is best to apply when the soil is slightly moist.

IN LAWNS AND ORNAMENTAL TURF

FOR BROWN PATCH (Rhizoctonia solani)

SOUTHERN GRASSES - For St. Augustine, Bermuda, etc. and Dichondra - Apply 3 3/4 lbs. / 1,000 sq.ft. in the Fall or the Spring at the first sign of infection. Lightly sprinkle the area to wash the granules to soil level. An additional application in 3 to 4 weeks may be needed for continued control if heavy rainfall or flooding occurs after this treatment.

NORTHERN GRASSES - For Bluegrass, Fescue, Rye, Bent, etc. - Apply 1 to 1 1/4 lbs. per 1,000 sq.ft. when disease first appears and repeat at 7 to 10 day intervals during warm, damp weather. Lightly sprinkle the area to wash granules to soil level. During lush growing periods a temporary discoloration of the grass may occur, however, this is not harmful and will disappear in a few days.

HELMINTHOSPORIUM LEAF SPOT

Apply at the rate of 2 1/2 to 3 3/4 lbs./1,000 sq. ft. Preventive applications should be made in the spring or fall of each year. For control of existing infections, apply at the higher rate as soon as symptoms are first noted. Following applications, the treated area should be lightly watered to insure movement through the grass to soil level. If the treated area is subjected to unusually heavy rainfall, or flooded, or if the disease is severe or reappears, the area should be retreated in 3 to 4 weeks.

SNOW MOLD

(Typhula incarnata, Fusarium nivale): Apply at the rate of 2 1/2 to 3 3/4 lbs./1,000 sq. ft. using a granular lawn spreader or comparable equipment. Application should be made prior to first snow fall. Do not graze treated areas or feed clippings to livestock

ORNAMENTAL PLANTS BEDDING PLANTS

For African Violets, Begonias, Carnations, Chrysanthemums, Poinsettias, Snapdragons and similar bedding plants. Treat bench soil prior to planting with 7 1/2 lbs. granules / 1,000 sq.ft. for control of Rhizoctonia, Stem rot. Thoroughly work into the top 2 inches of soil.

BULBS

For Tulips, Narcissus, Hyacinth, and Bulbous Iris in production - Apply 500 to 1,000 lbs. per acre to the soil as a pre-plant broadcast application for control of Crown rot and Black rot (Equivalent to 1.15 to 2.25 lbs./100 sq.ft.). Mix with the top 6 to 7 inches of soil with appropriate equipment. Use the higher rate in heavy organic soils and especially for the control of Black Rot.

CALENDULAS, LARKSPUR, SNAPDRAGON

For control of Sclerotinia and Stem rot. Apply 500 lbs./acre as a broadcast application 1 week before planting; (Equiv. to 1.15 lbs. per 100. sq.ft.). Mix with soil to a depth of 4 inches.

DUTCH IRIS

For Bulb and Stem Rot caused by Sclerotium rots disease: make a PRE-PLANTING BROADCAST SOIL APPLICATION of 500 to 1,000 lbs. per acre which is equivalent to 1.15 to 2.25 lbs. /100 sq.ft. Mix the granules with the soil to a depth of 2 to 3 inches. FOR A ROW BAND TREATMENT apply 60 lbs. per 14,500 sq.ft. of linear row in a 12 inch band centered on the row, at planting time. Follow this treatment with 3 additional band applications at 15 day intervals.

EASTER LILY

(Lilium Longiflorum Eximium): For control of Rhizoctonia Foot Rot. Apply AS A BROADCAST SOIL TREATMENT at the rate of 500 lbs. on light sandy soils and 1,000 lbs. per acre on heavy organic soils. Follow with a good double disking to mix granules with the soil to the depth of 6 to 7 inches.

GLADIOLUS

For Neck or Bulb Dry Rot caused by Stromatolium gladioli. Apply 600 lbs. per acre (equivalent to 1 1/3 lbs. per 100 sq.ft.) AS A BROADCAST SOIL TREATMENT. Thoroughly mix with soil to depth of 4 to 6 inches. Corms may be planted immediately.

IMPORTANT: PLEASE READ BEFORE USE

By using this product, the user accepts the following LIMITED WARRANTY: AMVAC warrants that (a) this product conforms to the chemical description on its label; (b) this product is reasonably fit for the purposes stated on its label, subject to the inherent risks referred to herein, when used in accordance with its directions; and (c) that the directions, warnings, cautions, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and plants, of residues on food crops, and upon reports of field experience. Testing has not been performed on all varieties of food crops, and plants, in all states, or under all application, weather and crop conditions. There are no express warranties other than those set forth herein. Amvac neither makes nor intends, nor does it authorize any agent or representative to make, any other warranty, express or implied. Amvac expressly excludes and disclaims all implied warranties of merchantability, fitness for particular purpose, or any other warranty of quality or performance.

This warranty does not extend to, and the user shall be solely responsible for, any loss or damage which results from the use of this product in any manner which is inconsistent with this label's directions, warnings or cautions. User's exclusive remedy and Amvac's or seller's exclusive liability for any claim, loss, damage, or injury resulting from the use or handling of this product, whether or not based in contract, negligence, strict liability in tort, or otherwise shall be limited, at Amvac's option, to replacement of, or repayment of the purchase price for, the quantity of product with respect to which damages are claimed. In no event shall Amvac or Seller be liable for special, indirect, or consequential damages resulting from the use or handling of this product. (FCNB20DG.PM5/09-22-94)

FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY
TRANSPORTATION: CHEMTREC 1-800-424-9300
MEDICAL: HAZARD INFORMATION SERVICES (H.I.S.) 1-800-228-5635 Ext. 169

EPA REG. NO 5481- EPA EST. 5481-CA-1

NET WEIGHT 50 POUNDS

AMVAC CHEMICAL CORPORATION

4100 EAST WASHINGTON BLVD. - LOS ANGELES, CA 90023 U.S.A.

BEST AVAILABLE COPY