


MA 25 5481-453

1-53

5

show
app



AUG 1 1995

MA 25 5481-453

193

PCNB 75 WSP

Wettable Powder in a Water Soluble Packet
SOIL FUNGICIDE FOR COTTON, PEANUTS, BEANS* BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, GARLIC*, PEPPERS, TOMATOES****

HOW TO USE WATER SOLUBLE PACKETS
 Add the required amount of water to spray forming tank. Drop the required number of unopened soluble packets of PCNB 75 WSP into the tank at once with the agitator running. Maintain agitation while adding the packets and the water in the tank. Water to be added to the packets of PCNB 75 WSP must be completely dissolved in the water prior to any application. When tanks mixing insure that complete dissolving of the PCNB 75 WSP has occurred prior to the addition of other products. Continue agitation until all of the spray solution has been applied.

ACTIVE INGREDIENT:
 Pentachloronitrobenzene(PCNB) 75.0%
INERT INGREDIENTS: 25.0%
TOTAL 100.0%

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

EPA Registration No. 5481- EPA Est. No. 51036-GA-1

STATEMENTS OF PRACTICAL TREATMENT
 If SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 to 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or comatose person. If EYES: Flush eyes with plenty of water. Get medical attention. If SKIN: Remove contaminated clothing and wash affected area with soap and warm water. Get medical attention if irritation persists. If INHALED: Remove victim to fresh air and give first aid as required.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
 CAUTION: Irritant if swallowed. Avoid inhaling dust. May cause skin irritation. Avoid contact with skin. Do not contaminate food or feedstuffs.

PERSONAL PROTECTIVE EQUIPMENT (PPE)
 Workers and other handlers must wear long-sleeved shirt and long pants; waterproof gloves; goggles or eye protection; and for exposures outdoors, a dust/filtration respirator (MSHA/NIOSH approved type that meets or exceeds the requirements for protection against particulates, respirator with either an organic vapor/acid gas cartridge with a prefilter approved for pesticides (MSHA/NIOSH approved type that meets or exceeds the requirements for protection against particulates, respirator with either a TC-23C, or a canister approved for pesticides (MSHA/NIOSH approved type that meets or exceeds the requirements for protection against particulates, use detergent and hot water. Keep and wash PPE as directed from other labels.

Use Safety Recommendations
 Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Do not eat, drink, or use tobacco while using this product. If pesticide gets inside, then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of your clothing as soon as possible. Wash thoroughly and change into clean clothing. Do not contaminate water when disposing of equipment washwaters. This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to irrigation areas below the mean high water mark. Apply this product to surface water only. Do not apply directly adjacent to potable water supplies.

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not plant root crops in PCNB treated fields within 12 months of broadcast and banding applications unless PCNB is registered for use on those crops. Do not apply this product through any type of irrigation system. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. *Not for any use in California on Garlic (Solanum tuberosum) as a postplant application on Snap or Dry Beans for control of white mold (Sclerotinia sclerotiorum).

AGRICULTURAL USE REQUIREMENTS
 Use this product only in accordance with the labeling and with the Worker Protection Standard, 29 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves and shoes plus socks.

STORAGE AND DISPOSAL
 Do not contaminate water, food or feed by storage or disposal. STORAGE: Store product in its original labeled container in a cool, dry, locked place out of reach of children. PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or empty containers is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

Use and Rates Chart PCNB 75 WSP

CROP	DISEASE	POUNDS / ACRE	OZS./1,000 FT. OF ROW	RECOMMENDED APPLICATION
BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER	Clubroot (Plasmodium brassicae)	2 to 6 pounds / acre Apply in 100 gallons of water. Use 1/2 to 3/4 pint of this solution per plant		Transplant Solution: The specified dosage should be based on the severity of the disease infection. Maintain agitation to hold powder in suspension.
		30 pounds / acre Apply in 25 gallons of water (40 inch row spacing for broccoli, brussels sprouts, and cabbage. 48 inch row spacing for cauliflower.) 40 pounds / acre Apply in 30 gallons of water	Broccoli, Brussels Sprouts, Cabbage, 36.8 ounces Cauliflower, 44 ounces	Band Application: Spray as a 12 to 15 inch band centered on the row and incorporate to a depth of 4 to 6 inches immediately after planting. Broadcast Application: Apply as a preplant broadcast spray prior to planting. Thoroughly incorporate to a depth of 4 to 6 inches using a disk or other suitable equipment. Broadcast Drench Application: Apply as a soil drench at the time of or immediately after seeding.
COTTON	Furrow Rot (Sclerotium rolfsii)	15 to 20 lbs./acre. Apply in 50 gallons water For smaller areas use 1 level table spoonful per gallon of water as a soil drench per 50 sq. ft. of seed bed. Apply with watering can or similar equipment.		Row Drench Treatment: Spray as an 8 inch band centered on the row at the time of, or immediately after seeding.
		10 to 15 pounds / acre Apply in 35 gallons water (40 inch row spacing for broccol, brussels sprouts and cabbage. 48 inch row spacing for cauliflower.) 12 1/2 pounds / acre Apply in 100 gallons of water (24 inch row spacing)	Broccoli, Brussels Sprouts, Cabbage 12 to 16.3 ozs Cauliflower 14.6 to 22 ozs	In Furrow Application: Apply as an in-furrow spray at planting time. Attach a spray rig to the machine planter and spray 8 to 12 inch bands using 2 nozzles per furrow. Direct front nozzle to spray the "seed" as it is dropped, direct the rear nozzle to spray the covering soil. Use good mechanical agitation in the spray tank to maintain a uniform suspension. Clay Treatment: Mix in sufficient water to apply dosage as a uniform mist-spray for complete coverage of treated cloves. For optimum results add a commercial sticker to the spray solution.
PEPPERS, TOMATOES**	Southern Blight (Sclerotium rolfsii)	3 to 5 pounds in 100 gallons of water. Use 1/2 pint of this solution / plant	Peppers, 11 ounces Flat Tomatoes, 22 ounces Staked Tomatoes, 14.6 ozs.	Apply at the time of transplanting. The solution should be agitated often to maintain a uniform suspension and assure proper dosage. Apply as an in-furrow spray to the open "V" trench, just prior to planting. When cultivating, set plows as flat as possible to avoid getting untreated soil against stems of plants.
		10 pounds / acre Apply in 100 gallons of water (36 inch row spacing to 48 inch row spacing for flat tomatoes, 48 inch row spacing for staked tomatoes.)		
TOMATOES	Late Blight (Phytophthora infestans)	1 1/2 to 2 3/4 pounds / acre (40 inch spacing) Apply at least 15 gals. of water. 2 1/2 inch row spacing 15 1/2 pounds / acre Apply in 8 to 10 gallons of water	1.6 to 3.4 ounces	Apply specified dosage as directed in-furrow spray to the seed and surrounding soil at planting time. Do not allow the feeding or grazing of cotton bolls by Freckle.
		4 1/2 pounds / acre	14.7 ounces	PRE-PLANTING: Apply specified dosage as a spray in an 8 to 12 inch band centered on the row and incorporated to a depth of 2 to 4 inches. Application may be made simultaneously with planting using suitable application equipment. AT-PEGGING BAND: Apply specified dosage as a spray in a 12 inch band centered on the row so as to cover the crown and pegging zones of the plant. If the row middles are cultivated at the same time, attach two spray nozzles (one on each side) to the cultivator sweeps. Do not move any untreated soil into the area, if necessary remove weeds weekly. AT-CULTIVATION: Apply specified dosage at cultivation. Do not apply more than 3 times per crop season. Attach two spray nozzles per row of inside cultivator sweeps (one on each side) and adjust so as to treat a total band of 12 inches centered on the row. Do not pile untreated soil on crown of plant.
BEANS***	Root and Stem Rot (Rhizoctonia solani)	1 1/2 to 2 pounds / acre Apply in 8 to 10 gallons of water (36 inch row spacing for bush type beans 42 inch row spacing for pole beans)	Bush Beans 14 to 2.2 ozs Pole Beans 2.3 to 3.8 ozs	Apply as directed spray in the seed furrow and to the covering soil at planting time.
		2 3/4 pounds / acre Apply in 8 to 10 gallons of water	Bush Beans, 3 ounces Pole Beans, 3.2 ounces	Apply as an 8 inch band centered on the row at or immediately after planting. If the disease problem is severe, applications may be repeated at 2 to 3 week intervals using directed spray application with one or two nozzles per row aimed at the soil at the base of the plants. Soil should remain undisturbed after application.

Note: Do not exceed 10 lbs. actual PCNB per acre in any one season, regardless of application methods used. Do not apply within 45 days of harvest. Do not feed treated excess livestock. Do not allow hogs to roam on peanuts.

GENERAL INFORMATION
 PCNB 75 WSP is a soilborne fungicide for control of the listed soilborne diseases of field and garden crops. It is a preventive program following the rates and the recommended application techniques as described in the following chart. Carefully read, understand and follow label directions.
VEGETABLES
 PCNB 75 WSP is also applicable to soilborne fungicide for control of the listed diseases which attack various vegetable crops. It can be accomplished by applying the product as a soil drench with mechanical incorporation or as an in-furrow application. Under certain crop uses (snap & dry beans) treated soil conditions may be necessary. Refer to the following Use & Rates chart for specific disease control and recommendations for the seed crop.

EMERGENCIES PHONE 24 HOURS A DAY
 AMVAC CHEMICAL CORPORATION
 4100 EAST WASHINGTON BLVD. - LOS ANGELES, CA 90023 U.S.A.
 HAZARDOUS MATERIAL SERVICE'S 24 HOURS: 1 800 228-5635 Ext. 189

IMPORTANT: PLEASE READ BEFORE USE
 By using this product, the user accepts the following LIMITED WARRANTY:
 AMVAC warrants that (a) this product conforms to the chemical description on its label, (b) this product is reasonably fit for the purposes stated on its label, and (c) that the directions, warnings, cautions, and other statements on the label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and plants, of residues on food crops, and upon reports of field experience. Testing has not been performed on all varieties of food crops, and plants, in all states, or under all application, weather and crop conditions.
 There are no express warranties other than those set forth herein. Amvac neither makes nor intends, nor does it authorize any agent or representative to make, any other warranty, express or implied. Amvac expressly disclaims and disavows all implied warranties of merchantability, fitness for particular purpose, or any other warranty of quality or performance.
 This warranty does not extend to, and the user shall be solely responsible for, any loss or damage which results from the use of this product in any manner which is inconsistent with this label's directions, warnings or cautions. User's exclusive remedy and Amvac's or seller's exclusive liability for any claim, loss, damage, or injury resulting from the use or handling of this product, whether or not based in contract, negligence, strict liability in tort, or otherwise shall be limited, at Amvac's option, to replacement of, or refund of, the purchase price for the quantity of product with respect to which damages are claimed. In no event shall Amvac or Seller be liable for special, indirect, or consequential damages resulting from the use or handling of this product. (DC 75-wsp PMS 1/07-08-95)

Water Soluble Packets - Net Weight 1 Pound
AMVAC CHEMICAL CORPORATION
 4100 EAST WASHINGTON BLVD. - LOS ANGELES, CA 90023 U.S.A.

BEST COPY AVAILABLE

BROADCAST TREATMENT (Cont.)

For **Leaf and Stem Rot** (Sclerotinia sclerotiorum) on **Gladiolus** use 3 2/3 lbs. 1,000 sq ft. Apply prior to planting. Thoroughly incorporate to depth of 4 to 6 inches. Bulbs may be planted immediately after solution is incorporated.

For **Stem Rot** (Rhizoctonia) on **Calendula**, **Larkspur**, **Snapdragon** and **Sweet Pea** use 3 1/2 lbs. per 1,000 sq ft. Apply prior to planting. Thoroughly incorporate to depth of 4 inches.

For **Rhizoctonia** on **Houseplant** **Seedlings** use 50 lbs. per acre. Mix and apply in 100 gallons of water to give uniform ground coverage. An additional 1 1/2 inch of water per acre should be applied by irrigation prior to planting. Seed may be planted immediately after irrigation or within one week of this date.

AS A BAND TREATMENT

For **Leaf and Stem Rot** (Sclerotinia sclerotiorum) on **Gladiolus** apply 5 1/2 lbs. per 1,000 linear feet of row as a 12 inch band centered on the row at planting time. Incorporate to depth of 4 inches. Bulbs may be planted immediately after incorporation.

For **Leaf and Stem Rot** (Sclerotinia sclerotiorum) on **Dutch Iris** use 1/2 lb. per 1,000 linear feet of row as a 12 inch band centered on the row at planting time. Follow with 3 additional applications at the same rate at 15 day intervals.

AS AN IN FURROW TREATMENT AT PLANTING TIME

For control of **Rhizoctonia Root Rot** on **Easter Lilies**, and for **Crown Rot** and **Black Rot** on **Iris**, **Hyalanth**, **Narcissus** and **Tulips** mix 4 1/2 to 6 lbs. in 5 to 10 gallons of water and apply per 1,000 linear feet of row. Direct the spray into open furrow to thoroughly cover entire furrow and bulbs. Use **low pressure** normal manner. Use **coverrate** for lighter soils and higher rate for heavier soils. For **Black Rot** control use high rate.

6

Disease/Host	Am't/Gal water per 1,000 Sq Ft	How and When to Apply
Brown Patch (Rhizoctonia solani) St. Augustine, Bermudagrass and similar warm season grasses	1 lb/10-15 Gals.	Apply once in spring or fall or when disease appears if it is severe or reappears, retreat 3 to 4 weeks later.
Dichondra	1 lb/40 Gals.	Treat as above
Bluegrass, Rye, Fescue, Bent and similar cool season grasses	3-4 oz/13-6 Gals.	Apply at 1st sign of disease. Repeat every 7-10 days during warm damp weather.
Helminthosporium spp.* Leaf Spot Dollar Spot* (Sclerotinia homocarpa) *Do not use in California	7-10 oz/15-10 Gals.	For preventive apply in spring or fall of each year. If disease is present apply at 10 oz rate as soon as possible. If disease is severe or reappears retreat area 3 to 4 weeks later.
Gray Snow Mold (Typhula spp.) Pink Snow Mold	8 oz/15-10 Gals.	1st treatment should be made prior to first snow fall.

3

Page 6 of 34

incubate treated areas of leaf grass clippings to livestock. Under certain growing conditions, complete incorporation of the grass occasionally may occur. This causes no harm and will disappear in a short time.

PRECAUTION: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the manufacturer. Manufacturer and user assume the risk of any such use.

ORNAMENTALS

Solutions of PCNB 75 WSP and water must be mixed with the soil as a drench treatment for protection against the listed diseases on nursery, greenhouse, field grown and established plants. A preliminary trial is suggested first before a full treatment is given to any plant species not listed in these directions.

DISEASES CONTROLLED

- | | |
|----------------------|--------------------------|
| Common Name | Fungal Name |
| Root and stem rot | Rhizoctonia spp. |
| Bulb and storage rot | Botrytis spp. |
| Leaf blight | Sclerotinia gladioli |
| Leaf spot | Sclerotium gladioli |
| Seed blight | Sclerotinia gladioli |
| Leaf blight | Sclerotinia gladioli |
| Bulb and stem rot | Sclerotium gladioli |
| Leaf blight | Sclerotium gladioli |
| Stem rot | Sclerotium gladioli |
| Black rot | Sclerotinia sclerotiorum |

4

NURSERY & GREENHOUSE

For **Rhizoctonia** control apply as a soil drench at time of seeding and/or transplanting **Tropical Foliage Plants**

Agave, Apatandra, Daffodil, Dracena, Ficus, Fiddle Leaf, Gynura, Hoya, Peperomia, Pothos, Schefflera, Syngonium, Tradescantia and similar tropical foliage plants. Mix 4 to 6 oz. in 100 gallons water and apply to 400-800 sq ft. of bed or bench area.

Coverage: For soil depths of 4 inches or less, apply 100 gallons to 800 sq ft. For soil depths of more than 4 inches apply 100 gallons to 400 sq ft. One repeat application can be made 4 to 6 weeks later if necessary. Do not apply to **Philodendron** or **Pilea** since injury may occur.

Bedding and Flowering Plants

Carnation, Chrysanthemum, Dahlia, Geranium, Mangold, Pelonia, Poinsettia, Portulaca, Salvia, Vinca, Zenia and similar plants. Use 4 oz. with 100 gallons of water and apply to 800 sq ft. of bed or bench area. Apply at seeding or transplanting time.

Herbaceous and Woody Plants

Herbaceous Plants: Kalanchoe, Lily and similar plants. **Woody Plants:** Azalea, Boxwood, English Ivy, Holly, Juniper, Pine, Rhododendron and similar plants. Use 4 to 8 oz. with 100 gallons of water and apply to 400-800 sq ft. of bed or bench area.

FIELD GROWN ORNAMENTAL CROPS

HOW TO MIX AND APPLY

Mix the following amounts of PCNB 75 WSP in sufficient water and apply before planting to give uniform ground coverage. Incorporate the spray/drench solution to depth of soil as indicated.

AS A PRE-PLANT BROADCAST TREATMENT

For **Rhizoctonia Root Rot** on **Easter Lilies**, and for **Crown Rot** and **Black Rot** on **Iris**, **Hyalanth**, **Narcissus** and **Tulips** use 3 1/4 to 6 1/2 lbs. per 1,000 sq ft. Thoroughly incorporate to a depth of 6 to 7 inches. Use the lower rate for light soils, the higher rate for heavy soils.

For **Bulb Rot** and **Stem Rot** (Sclerotinia rot) on **Dutch Iris** use 3 1/4 to 4 1/2 lbs. per 1,000 sq ft. Apply prior to planting. Thoroughly incorporate solution to depth of 2 to 3 inches.

5

PRECAUTIONARY STATEMENTS

HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Irritation to eyes, nose, throat, and skin. Avoid inhaling dust. May cause sunburn-like. Avoid contact with skin. Do not contaminate feed or food crops.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

For all uses, operators must wear long-sleeved shirt and long pants, waterproof gloves, shoes and socks and for exposures outdoors, a Dust/Mist filtering respirator (MSHA/NIOSH approved type "N" or "P" for particulates in enclosed areas, a respirator with either an organic vapor cartridge or a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-11). For all uses, operators must wear eye protection (MSHA/NIOSH approved type "G" eye protection). For all uses, operators must wear protective clothing (MSHA/NIOSH approved type "G" eye protection). For all uses, operators must wear protective clothing (MSHA/NIOSH approved type "G" eye protection).

User Safety Recommendations

Wash hands thoroughly with soap and water after handling this product. Wash the outside of gloves before removing. As soon as possible, after handling this product, wash the outside of gloves before removing. As soon as possible, after handling this product, wash the outside of gloves before removing. As soon as possible, after handling this product, wash the outside of gloves before removing.

HAZARDOUS TO AQUATIC LIFE

Do not contaminate water when disposing of equipment washwaters. This product is toxic to fish and aquatic invertebrates. Do not discharge from treated areas into streams, rivers, lakes, ponds, or other bodies of water. Do not apply directly to water, to areas where surface waters exist, or to areas where water may be present. Do not apply to areas where water may be present. Do not apply to areas where water may be present.

STORAGE AND DISPOSAL

Do not store in water. Store in a cool, dry, locked place out of reach of children. **PESTICIDE:** Do not store in a way that allows access to the area during application. Do not store in a way that allows access to the area during application. Do not store in a way that allows access to the area during application. Do not store in a way that allows access to the area during application.

6

(pc75wsp2 pm5107-06-95)

DIRECTIONS FOR USE

Read and follow the label directions for use. Do not use in a manner inconsistent with its labeling. Do not use in a manner inconsistent with its labeling. Do not use in a manner inconsistent with its labeling. Do not use in a manner inconsistent with its labeling.

HOW TO USE WATER SOLUBLE PACKETS

Drop the required number of packets into the tank at once with the agitator running. Maintain the agitator running for 15 minutes. Do not stop the agitator until the packets are completely dissolved and dispersed in water prior to initiating spray application. When tank is empty, ensure that complete dispersion of the PCNB 75 WSP has occurred prior to the addition of other products. Continue agitation until all of the spray solution has been applied.

FOR TURFGRASS

Use as a preventive and control of the listed soil borne diseases which attack lawns on golf courses, sod farms, home lawns, institutional, ornamental and recreational areas where the turf is under a maintenance program.

DISEASES CONTROLLED

- Brown Patch (Rhizoctonia solani)
- Leaf Spot (Helminthosporium spp.)
- Dollar Spot (Sclerotinia homocarpa)
- Gray Snow Mold (Typhula spp.)
- Pink Snow Mold (Fusarium nivale)

For best results use PCNB 75 WSP as a preventive application before diseases appear, or apply at the first sign of infection. After application to the treated areas should be restricted to the time of down to the soil level. If re-infection or flooding occurs after treatment, or if disease is severe or reappears, retreat again. Follow the rates as outlined below for the respective disease.

(PC75WSP2 PM5 (7-6-95))

2

SUPPLEMENTAL AND ADDITIONAL LABELING FOR
See Basic Label for use of this product on Food & Fiber Crops
EPA REG. NO. 5481- EPA EST NO. 51036-GA-1

PCNB 75 WSP

Wettable Powder in a Water Soluble Packet

A SOIL FUNGICIDE FOR PREVENTION AND CONTROL OF SOIL BORNE DISEASES WHICH ATTACK TURFGRASS, NURSERY & GREENHOUSE PLANTS, FIELD GROWN ORNAMENTALS, BEDDING & FLOWERING PLANTS, HERBACEOUS & WOODY ORNAMENTALS.

ACTIVE INGREDIENT:

Pentachloronitrobenzene (PCNB) - - - 75.0%

INERT INGREDIENTS: - - - 25.0%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN - CAUTION

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 to 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists. **IF ON SKIN:** Wash affected area with soap and warm water. Get medical attention if irritation persists. **IF INHALED:** Remove victim to fresh air and give artificial respiration if indicated.

AMVAC CHEMICAL CORPORATION

4100 EAST WASHINGTON BLVD - LOS ANGELES, CA 90023 U.S.A.

GROUND SPRAY for ESTABLISHED AZALEAS and CAMELIAS

For **Azalea Petal Blight** (Oomyces azaleae) on **Azaleas** and **Camelia Flower Blight** on **Camellia** use 1 to 1.5 lbs. per 100 sq ft. Mix and apply in sufficient water (5 to 10 gallons per 100 sq ft.) to give uniform coverage to soil beneath bushes and the plant bed. Begin application prior to opening of buds and repeat every 3 to 4 weeks during bloom season.

DIP TREATMENT for BULBS and DORMANT ROSES

Use before planting or storing. It is important to maintain continuous circulation of dip solution to assure uniform application of fungicide. Add 1% "Sticker" to solution. Use wire mesh for dipping and drain boards to return excess solution to dip tank. Dispose of unused dip solution according to "Storage and Disposal" and observe use of protective clothing (gloves etc.) as listed under "Hazardous to Humans".

For control of **Crown Rot** and **Black Rot** on **Iris**, **Hyalanth**, **Narcissus** and **Tulips** dip bulbs for 5 minutes in a solution containing 2 lbs. PCNB 75 WSP in 3 1/2 gallons of water.

For **Root Rot** (Rhizoctonia) of **Easter Lilies** dip in a solution containing 4 to 6 lbs. PCNB 75 WSP in 100 gallons of water for a period of 5 to 15 minutes.

For **Botrytis Storage Rot** on **Roses**: Dip or spray dormant plants before packaging or storage, with a solution of 2 lbs. PCNB 75 WSP in 100 gals. of water.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. The Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow work to be done in treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, or covers, waterproof gloves and shoes plus socks.

7

BEST COPY AVAILABLE

3