

Ms. Brenda K. Balsz
AMVAC
4695 Mac Arthur Court, Suite 1250
Newport Beach, CA 92660

OCT 2 2002

Dear Ms. Balsz:

Subject: Label Amendment- final printed labels
Ambush 25W Insecticide Wettable Powder
EPA Registration Number 5481-502
Your submission dated September 20, 2002

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a) Update your Statement of Practical Treatment section according to PR Notice 2001-6.
3. Submit two (2) copies of your final printed labeling before you release

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

George T. LaRocca
Product Manager 13
Insecticide Branch
Registration Division (7505C)

Enclosure

RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and
only for those uses covered by the Certified Applicator's certification.



WETTABLE POWDER

ACTIVE INGREDIENT:

Permethrin

(3-Phenoxyphenyl)methyl(z)-cis, trans-3-(2,2-dichloroethyl)-

2,2-dimethylcyclopropanecarboxylate* 25.0%

INERT INGREDIENTS 75.0%

TOTAL 100.0%

*Cis/trans ratio: Min. 35% (z) cis and max. 65% (z) trans.

FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY:

Transportation: CHEMTREC 1-800-424-9300

Other: AMVAC 1-323-264-3910

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE.

EPA Reg. No. 5481-502



EPA Est. No. 37429-GA-2

4100 E. Washington Blvd.

Los Angeles, CA 90023

Made in U.S.A.

1-323-264-3910 • www.amvac-chemical.com

NET WEIGHT 2 lbs.

11483-1



APPROVED
by the California State Board
for Pesticide Control

OCT - 2 2002

Product Name: Permethrin
EPA Reg. No.: 5481-502
EPA Est. No.: 37429-GA-2
Net Weight: 2 lbs.
11483-1

5481-502

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with water for 15 minutes. Get medical attention if irritation persists.

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May cause eye irritation. Harmful if swallowed. Do not get in eyes, on skin or on clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Waterproof gloves.
- Shoes plus socks.
- Protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Waterproof gloves.
- Shoes plus socks.
- Protective eyewear.

GENERAL INFORMATION

Apply AMBUSH 25W as shown in "Directions for Application". Use sufficient water to obtain full coverage of foliage. Timing and frequency of applications should be based upon insect populations reaching economic thresholds. The higher rates may be required under conditions of heavy infestation. Do not apply more than 2.0 lbs. a.i. per acre per season. Crops in this label may be planted immediately after last treatment. Do not plant other crops within 60 days after last application. Any other crop may be planted if it is not harvested or grazed.

Resistance: Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your Local or State Agricultural Authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

CHEMIGATION**SPRINKLER IRRIGATION APPLICATION**

Apply AMBUSH 25W Insecticide at rates and timing described elsewhere in this label. Apply with center pivot or continuous-move equipment distributing 1/2 acre-inch or less during treatment. If stationary (solid sets, handlines or wheel lines other than continuous-move) are used, AMBUSH 25W insecticide should be injected into no more than the last 20-30 minutes of the set. Do not apply when winds are greater than 10-15 mph to avoid drift or wind skips. Thorough coverage of foliage is required for good control. Good agitation should be maintained during the entire application period.

USE PRECAUTIONS-SPRINKLER IRRIGATION APPLICATION

- A. Apply this product only through (sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.
- B. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
- C. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- D. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- E. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- F. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- G. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- H. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- I. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- J. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- K. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- L. Do not apply when wind speed favors drift beyond the area intended for treatment.
- M. Do not apply through chemigation systems connected to public water systems.

6/20

| DIRECTIONS FOR APPLICATION | | | |
|--|---|----------------------------|---|
| CROP | TARGET PESTS | RATE LBS AI/A (OZ/A) | REMARKS |
| Cauliflower | Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid* | 0.05-0.1 (3.2-6.4) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 days or as needed. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not apply within 1 day of harvest. |
| Chinese Broccoli [†] (gai lan, white flowering) | Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid* | 0.05-0.1 (3.2-6.4) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 days or as needed. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not apply within 1 day of harvest. |
| Chinese Cabbage (Tight-heading varieties only) | Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid* | 0.05-0.2 (3.2-12.8) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5-10 days or as needed. Do not apply more than 1.0 lb. a.i. per acre per season. Do not apply within 1 day of harvest. |
| Collards (AR, AZ, GA, IL, NC, OK, SC & TX only) | Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid* | 0.05-0.1 (3.2-6.4) | Apply by ground equipment only. Apply in a minimum of 10 gallons of water per acre. Spray using sufficient water to obtain uniform coverage. Apply as needed. Do not exceed 8 applications. Do not apply within 1 day of harvest. |

7/20

| DIRECTIONS FOR APPLICATION | | | |
|----------------------------|--|----------------------------|---|
| CROP | TARGET PESTS | RATE LBS AI/A (OZ/A) | REMARKS |
| CUCURBIT VEGETABLES | | | |
| Cucurbit Vegetables+ | Leafminers and Squash Bugs | 0.2 (12.8) | Apply by air or ground equip- ment. Apply a minimum of 4 gallons of finished spray per acre by air or 20 gallons of finished spray per acre with ground equipment. Do not apply more than 1.6 pounds active ingredient per acre per season. AMBUSH 25W may be applied up to harvest. |
| | Cabbage Looper Cucumber Beetle (adults) Cutworms Melon Worm Pickworm Plant Bugs (including Lygus and Stink Bugs) Rindworms and Squash Vine Borer | 0.1-0.2 (6.4-12.8) | |
| FRUITING VEGETABLES | | | |
| Eggplant | Colorado Potato Beetle | 0.2 (12.8) | Apply by air or ground equip- ment. By air, apply in a mini- mum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform cov- erage. Apply as needed. Do not apply more than 2.0 lbs. a.i. per acre per season. Do not apply within 3 days of harvest. |
| | Vegetable Leafminer Cabbage Looper Flea Beetle | 0.1-0.2 (6.4-12.8) | |
| Peppers (Bell) | Vegetable Leafminer Cabbage Looper Flea Beetle Pepper Weevil | 0.1-0.2 (6.4-12.8) | Apply by air or ground equip- ment. By air, apply in a mini- mum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform cov- erage. Apply as needed. Do not apply more than 1.6 lbs. a.i. per acre per season. Do not apply within 3 days of harvest. |
| | European Corn Borer | 0.2 (12.8) | |

| DIRECTIONS FOR APPLICATION | | | |
|--|--|----------------------------|--|
| CROP | TARGET PESTS | RATE LBS AI/A (OZ/A) | REMARKS |
| Tomatoes | Tomato Fruitworm Cabbage Looper Tomato Pinworm Vegetable Leafminer Beet Armyworm Southern Armyworm Colorado Potato Beetle Granulate Cutworm Hornworm | 0.05-0.2 (3.2-12.8) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Do not apply more than 1.2 lbs. a.i./acre per crop season. Can be applied up to the day of harvest. Do not apply as a band. Do not use on tomato varieties in which the mature tomatoes will be less than 1 inch in diameter (such as cherry tomatoes). |
| LEAFY VEGETABLES | | | |
| Leafy Vegetables* (except Brassica) | Vegetable Leafminer Cabbage Looper Cutworm Complex Fall Armyworm Beet Armyworm Corn Earworm Green Cloverworm Leafhoppers European Corn Borer | 0.1-0.2 (6.4-12.8) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 3-5 days or as needed. Do not apply more than 2.0 lbs. a.i. per acre per season. Can be applied up to 1 day of harvest. Spinach only-Do not apply more than 1.0 lb. a.i. per acre per season. |

9/20

| DIRECTIONS FOR APPLICATION | | | |
|--|--|---|--|
| CROP | TARGET PESTS | RATE LBS AI/A (OZ/A) | REMARKS |
| ROOT, TUBER AND BULB VEGETABLES | | | |
| Onions (Dry Bulb and Garlic) | Onion Thrips Armyworms Cutworms Leafminers Onion Maggots (adult) Stink bugs | 0.15-0.3 (9.6-19.2) 0.1-0.3 (6.4-19.2) | Apply by air or ground equip- ment. Apply in a minimum of 20 gallons of water per acre with ground equipment or in a minimum of 6 gallons per acre by air as foliar spray. Begin applications when pests appear and repeat as necessary to maintain con- trol. Use the higher label rates as onion thrips popula- tion increases and avoid res- cue situations. Do not apply more than 2.0 lbs. active per acre per season. Do not apply within 1 day of harvest. |
| Potatoes | Colorado Potato Beetle Potato Leafhopper Aster Leafhopper Potato Flea Beetle Potato Aphid Potato Tuberworm Potato Psyllid Cabbage Looper | 0.05-0.2 (3.2-12.8) | Apply by air or ground equip- ment. By air, apply in a mini- mum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply as needed. Do not apply more than 1.6 lbs. a.i. per acre per season. Can be applied up to 14 days prior to harvest. |
| Turnips (FL, GA, IL, IN, OK, SC, TX, WA only) (TX, OK, IL, IN only) (SC, GA, FL, WA only) | Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a | 0.05-0.1 (3.2-6.4) | Apply by ground equipment only. Apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply as needed. Do not apply within 1 day of harvest. Do not graze treated areas or feed crop refuse to livestock. Do not exceed 8 applications. Do not exceed 4 applications. |

| DIRECTIONS FOR APPLICATION | | | |
|---|---|----------------------------|---|
| CROP | TARGET PESTS | RATE LBS AI/A (OZ/A) | REMARKS |
| CEREAL GRAINS | | | |
| Field Corn/ Popcorn/Field Corn Grown for Seed (Preemergent Use)** | Armyworm Cutworms | 0.1-0.2 (6.4-12.8) | Apply by air or ground equip- ment. By air, apply in a mini- mum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Apply in the time period from 5 days prior to planting up to emergence of the crop. |
| Field Corn/ Popcorn/Field Corn Grown for Seed (Foliar Use)** | Cutworms European Corn Borer Armyworm Southwestern Corn Borer Flea Beetle Corn Earworm*** Corn Rootworm Beetle*** Tobacco Budworm*** Stalk Borer | 0.1-0.2 (6.4-12.8) | Apply by air or ground equip- ment. By air, apply in a mini- mum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Do not apply more than 0.6 lbs. a.i. per acre per season. Allow a minimum of 6 days between treatments. Do not apply within 30 days of har- vest of grain or fodder (slover). Forage may be harvested on the day of application. When treating for stalk borer, AMBUSH 25W must be applied when or shortly before the stalk borer larvae are moving into the corn from surrounding weeds and grasses. Mowing or burn- down herbicide is suggested to initiate movement.** |

11/20

| DIRECTIONS FOR APPLICATION | | | |
|------------------------------|---|----------------------------|---|
| CROP | TARGET PESTS | RATE LBS AI/A (OZ/A) | REMARKS |
| Sweet Corn | Corn Earworm European Corn Borer Fall Armyworm Corn Rootworm Adults Cutworms Aster Leafhopper Flea Beetle* | 0.1-0.2 (6.4-12.8) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Apply every 5 days or as needed. Do not apply more than 1.2 lbs. a.i. per acre per season. Do not apply within 1 day of harvest of ears or forage or livestock grazing. AMBUSH 25W may be mixed with LANNATE* (methomyl) in accordance with label limitations and precautions. Label dosage rates should not be exceeded. |
| Sweet Corn (Florida only) | Corn Earworm European Corn Borer Fall Armyworm Southern Armyworm Corn Rootworm Adults Cutworms Aster Leafhopper | 0.1-0.25 (6.4-16.0) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Apply every 5 days or as needed. Do not apply more than 2.0 lbs. a.i. per acre per crop season. AMBUSH 25W may be mixed with LANNATE* (methomyl) in accordance with label limitations and precautions. Label dosage rates should not be exceeded. |
| TREE CROPS | | | |
| Apples | Apple Aphid Redbanded Leafroller Oblique banded Leafroller Plum Curculio White Apple Leafhopper Spotted Tentiform Leafminer Tarnished Plant Bug | 0.1-0.4 (6.4-25.6) | Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Do not apply after petal fall. |

| DIRECTIONS FOR APPLICATION | | | |
|----------------------------|--|----------------------------|--|
| CROP | TARGET PESTS | RATE LBS AI/A (OZ/A) | REMARKS |
| Avocados | Thrips Leafhoppers Mirid bugs Lepidopterous larvae | 0.2 (12.8) | Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Apply when insects appear and repeat at 7-10 day intervals as needed to provide control. Do not make more than 6 applications per season. Do not apply within 7 days of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |
| Cherries | Lesser Peachtree Borer Rose Chafer Green Fruitworm Redbanded Leafroller Plum Curculio Tarnished Plant Bug | 0.1-0.2 (6.4-12.8) | Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Apply when insects appear. In the East, do not exceed 6 applications per season, with no more than 4 applications after petal fall. In the West, do not apply more than 4 applications per season, with no more than 3 applications after petal fall. Do not apply within 3 days of harvest. Do not graze livestock in treated orchards. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |
| Filberts | Filbertworm Oblique Banded Leafroller | 0.2-0.4 (12.8-25.6) | Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply when insects appear. Do not apply more than 1.6 lbs. a.i. per acre per season. Do not apply within 14 days of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |

| DIRECTIONS FOR APPLICATION | | | |
|----------------------------|---|----------------------------|---|
| CROP | TARGET PESTS | RATE LBS AI/A (OZ/A) | REMARKS |
| Papaya (Florida only) | Fruit Fly | 0.1 (6.4) | Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Apply when insects appear and repeat at 7-10 day intervals as needed to provide control. Do not make more than 6 applications per season. Do not apply within 7 days of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |
| Peaches | Peach Twig Borer Oriental Fruit Moth Green Fruitworm Lesser Peachtree Borer Plum Curculio Tarnished Plantbug Rose Chaler | 0.1-0.3 (6.4-19.2) | Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Can be applied up to 14 days before harvest. Do not apply more than 1.5 lbs. a.i. per acre per season. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |
| Pears (Dormant) | Pear Psylla | 0.2-0.4 (12.8-25.6) | Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply at a rate of 0.8 to 1.6 pints (0.2 to 0.4 lbs. active) per acre. May be combined with 2 to 8 gallons of spray oil per acre. Apply during the dormant through delayed dormant growth periods only. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |

| DIRECTIONS FOR APPLICATION | | | |
|----------------------------|---|----------------------------|--|
| CROP | TARGET PESTS | RATE LBS AI/A (OZ/A) | REMARKS |
| Pears (Pre-bloom) | Pear Psylla Codling Moth Green Fruitworm | 0.2-0.4 (12.8-25.6) | Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply two pre-bloom sprays. Pre-bloom sprays can be applied from dormant through bud burst stages. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |
| Pistachios | Peach Twig Borer Naval Orangeworm Chinch Bug | 0.2-0.4 (12.8-25.6) | Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply at a rate of 0.8 to 1.6 pints (0.2-0.4 lbs. active) per acre. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not apply after 10 percent hull split. Nuts may be harvested on the day of application. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |
| Walnuts | Naval Orangeworm Codling Moth Walnut Husk Fly | 0.2-0.4 (12.8-25.6) | Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply when insects appear. Do not apply more than 1.6 lbs. a.i. per acre per season. Do not apply within 1 day of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |

| DIRECTIONS FOR APPLICATION | | | |
|---|---|---|--|
| CROP | TARGET PESTS | RATE LBS A/A (OZ/A) | REMARKS |
| Mushrooms (Mushroom houses and adjacent premise areas) | Mushroom Flies (Sciand and Phorid Adults) | Apply 6 oz. ¹ to 11 gal. water or 8 level table- spoons to 3 gal. water. As a guide, use 1 gal. spray per 750 sq. ft. (400-1,400 sq. ft.) | Spray directly to walls and ceilings as residual surface treatment only. Spray to point of runoff. Use AMBUSH 25W prior to filling house, during cooldown, during spawning, up to pinning and between breaks. Do not use when mushrooms are present. Treat as needed when flies appear. Do not make more than 20 applications prior to pinning of first break; apply no more than two applications between each break. Do not apply more than 30 applications total per crop of 5 breaks. Do not apply within 3 days of harvest. |
| Soybeans | Green Cloverworm Mexican Bean Beetle Bean Leaf Beetle Cabbage Looper Potato Leafhopper Velvetbean Caterpillar Saltmarsh Caterpillar | 0.05-0.1 (3.2-6.4) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Do not apply more than 0.4 lbs. a.i. per acre per season. Can be applied up to 60 days before harvest. Do not graze treated areas or harvest for forage or hay. |
| | Corn Earworm Beet Armyworm Japanese Beetle*** Soybean Looper* | 0.1-0.2 (6.4-12.8) | |
| ORNAMENTALS | | | |
| Asparagus Fern | Japanese Beetle Asparagus Beetle Lygus Bugs | 0.1 (6.4) | Apply to the fern stage of the asparagus plant after spear harvest when larval and adult stage of the asparagus beetle, lygus bugs and the adult stage of the Japanese beetle are present. |
| Christmas Trees* (container and field grown) | Nantucket Pine Tip Moth | 0.1-0.2 (6.4-12.8) | Apply first application when adults appear and repeat at 5-7 day intervals or as needed throughout the season. Use sufficient water to obtain full coverage of foliage. |

| DIRECTIONS FOR APPLICATION | | | |
|--|--|--------------------------------|---|
| CROP | TARGET PESTS | RATE LBS AI/A (OZ/A) | REMARKS |
| Chrysanthemums (Florida only) | Vegetable Leafminer | 0.1-0.2 (6.4-12.8) | Apply by ground. |
| Conifers ^a (Container and Field Grown) | Nantucket Pine Tip Moth | 0.1-0.2 (6.4-12.8) | Apply first application when adults appear and repeat at 5-7 day intervals or as need- ed throughout the season. Use sufficient water to obtain full coverage of foliage. |
| Ornamentals Nursery Stock ^a (field-grown) | Beet Armyworm Cabbage Looper Citrus Thrip Heliothis Spp. Leafhoppers Leafminers Whiteflies Cutworms | 0.1-0.2 (lbs ai/100 gal) | Marginal leaf burn may occur on Salvia, Dieffenbachia and Pteris Fern. Application to blooming plants may cause browning of petals. Not all species and varieties have been tested. Before treating large numbers of plants of a particular variety, treat a few plants and observe prior to full scale application. Apply to nonbearing plants of fruit- ing species. |
| Roses ^a (field-grown) | Heliothis Spp. | 0.1-0.2 (6.4-12.8) | Apply by air or ground equip- ment. Apply 2 gallons/acre by air and 10 gallons of fin- ished spray per acre with ground equipment. |
| Roses ^a (greenhouse) | Beet Armyworm Cabbage Looper Omnivorous Leafroller | 8 fl oz/100 gals water | AMBUSH 25W is not phyto- toxic to the following varieties of greenhouse roses: Ballena, Bettina, Cara Mia, Coquette, Exclamation, Forever Yours, G. Wave, Jack Frost, Jr. Bridesmaid, Matador, Paul's Pink, Samantha, Seventeen, Town Crier, Tropicana and Visa. Other varieties may vary in their sensitivity to AMBUSH 25W, and a small number of plants should be treated under local conditions to determine plant safety prior to commercial use. |

NOTE: Do not tankmix AMBUSH 25W Insecticide with BICEP®, GRAMOXONE® SUPER/GRAMOXONE® EXTRA.

- **LEAFY VEGETABLES** (Except Brassica): Amaranth (leafy amaranth, Chinese spinach, tam-pala); arruglia (Roquette); Cardoon; celery; Chinese celery; celtuce; chervil; corn salad; chrysanthemum, edible-leaved; chrysanthemum, garland; cress, garden; cress upland (yellow rocket, winter cress); dandelion; dock (sorrel); endive (escarole); fenmel, Florence; lettuce (head and leaf); orach; parsley; purslane, garden; purslane, winter; radicchio (red chicory); rhubarb; spinach; spinach, vine (Malabar spinach, Indian spinach); spinach, new Zealand; Swiss chard.
- **CUCURBIT VEGETABLES:** Chayote (fruit); Chinese Waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourds, edible (includes hyotan, cucurza, hechima, Chinese okra); Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); muskmelons, including hybrids such as true cantaloupe, cantaloupe, casaba, crenshaw, golden pershaw melon, honeydew melons, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon; pumpkin; squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini, and winter (butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon, including hybrids.
- **AMBUSH 25W** may be applied in tank mixtures with the following herbicides in accordance with label limitations and precautions. Label dosage rates should not be exceeded. LASSO®/Atrazine Premix; LASSO® 4E & ME; DUAL® RE; Atrazine 4L & BOW; BLADEX® BOW & 4L; PRINCEP®; PRINCEP® CALIBER® 90; 2,4-D; ROUNDUP®, BRONCO®, BANVEL®, EXTRAZINE®.

*** Not for control of this insect in California.

• Aids in suppression.

• Currently not registered for use in California.

• 1 oz. of this powder equals 5 level teaspoons.

• Maximum use rate in California is 0.1 lb. a.i./Acre.

• Under heavy insect pressure use 0.2 lbs. a.i./Acre. A second application of alternate chemistry may be necessary to reduce populations below economic thresholds.

| RATE CONVERSION CHART | | |
|-----------------------|----------------------------|-------------------------|
| POUNDS ACTIVE/ACRE | FORMULATION OUNCES/ACRE | FORMULATION LBS/ACRE |
| 0.05 | 3.2 | 0.2 |
| 0.10 | 6.4 | 0.4 |
| 0.125 | 8.0 | 0.5 |
| 0.15 | 9.6 | 0.6 |
| 0.175 | 11.2 | 0.7 |
| 0.20 | 12.8 | 0.8 |

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container.

STORAGE: Mix as needed. Do not store diluted material. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, sweep up and remove to chemical waste area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use, subject to the inherent risks referred to herein, when it is used in accordance with such directions; and (c) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions.

THERE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT, SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

AMVAC offers this product, and Buyer accepts it, subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC.

AMBUSH®, GRAMOXONE®, BICEP®, CALIBER®, DUAL®, and PRINCEP® are trademarks of a Syngenta Group Company.

Ammo®, Capture®, Mustang® and Pounce® are trademarks of FMC Corporation.

Asana®, Bladex®, Extradine® and Lannate® are trademarks of E. I. duPont de Nemours & Co.

Bronco®, Lasso®, and Roundup® are trademarks of Monsanto Company.

Barvel® is a trademark of Sandoz Crop Protection Corp.

Baythroid® is a trademark of Mobay Corporation.

Danitol® is a trademark of Sumitomo Chemical Company, Ltd.

Amvac Chemical Corporation

4100 E. Washington Boulevard

Los Angeles, CA 90023 U.S.A.

1-323-264-3910

www.amvac-chemical.com