10911



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

September 17, 2007

Ms. Kaila Moran Product Manager AMVAC CHEMICAL COMPANY 4695 MacArthur Court Suite 1250 Newport Beach, CA 92660

Subject: Label Notification

Dear Ms. Moran:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated <u>8/17/07</u> for **EPA Registration 5481-521**. The Registration Division (RD) has conducted a review of this request for applicability under PRN 98-10 and finds that the label change(s) requested falls within the scope of PRN 98-10. The label has been date-stamped "Notification" and will be placed in our records.

If you have any questions, please contact me directly at 703-305-6249 or Sherada Hobgood of my staff at 703-308-8893.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader

Registration Division (7505P)

Office of Pesticide Programs

Please read instructions on reve	rse before completing t	form.			·	Form	Approved, Of	/IB No. 2070	201 0-0060, Approval expires 2-28-
\$ EPA	United States Environmental Protection A Washington, DC 20460			gency	☐ Registra		tion	OPP Identifier Number	
	A	pplication	n for F	estic	ide - Sec	tion)		
1. Company/Product Number			2. EPA Product Manager			•	3. Propo	sed Classification	
	5481-521			George La Rocca			ca		
4. Company/Product (Name)				PM#				X None	e Restricted
Wisdom L	awn Granular Insection	cide			13	3			
5. Name and Address of App Amvac Chemical Corporat 4695 MacArthur Court, Su	tion ite 1250	Code)		proc		ar or ic	dentical in co	ompositio DTIFIC	RA Section 3(c)(3)(b)(i), monand labeling to: ATION
Newport Beach, CA 92660 Check if this is a new address			٠.	Product Name			(SEP 17 2007	
			Spet	ion -	11				
Amendment - Explain b	elow		0601			labels	in response	to Agend	cy letter dated
Resubmission in respo		r dated		ப	•			to rigoni	
Notification - Explain be		uateu	-	" Me Too" Application Other - Explain below					
Explanation: Use additional	page(s) if necessary	. (For Section	n I and Se	ction II.)	<u> </u>				
and models on labeling This notification is consistent wit or the confidential statement of further understand that if this not be subject to enforcement action	th the provisions of PR formula of this product tificaiton is not consiste	Notice 98-10 t. I understan ent with the te	and EPA re d that it is a rms of PR N	gulaitons violation otice 98-	s at 40 CFR 1 of 18 U.S.C.	Sec. 1	1001 to willfully	y make any	y false statement to the EPA.
			Sect	ion -	<u> </u>		·	_	,
1. Material This Product Will	T		T,						
Child-Resistant Packaging	Unit Packaging		Water		Packaging		2. Type of		•
☐ Yes* ☑ No * Certification must	☐ Yes ☐ No If "Yes" Unit Packaging	No. per	☐ ☑ If "Yes	Yes No s" ge wt.	No. per container			Metal Plastic Blass Paper	•
be submitted	wt.						C	Other (Spe	ecify)
3. Location of Net Contents	Information	4. Size(s) F	Retail Cont	ainer		5. Lo	cation of Lal	pel Direct	ions
☑ Label ☐ Container			☐ On Label☐ On Labeling accompanying product			ying product			
6. Manner in Which Label is	Affixed to Product		graphed glued				tenciled ther		
			Secti	on –	IV				
1. Contact Point (Complete it	tems directly below	for identifica	ation of inc	lividual	to be contac	ted, if	necessary,	to proces	s thisှ ^c a်ρ်ပျံငation.)
Name Kaila Moran			Title	Product Manager			Telephone No. (ใก้เป็นde Area Code) 949 ู้-221 6109		
I certify that the statem	ents I have made o	Certification this form		ttachme	nts thereto	are tr	ue, accurate	e and	8. Date Application o ç c

complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or

3. Title

5. Date

Kaila Moran EPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete.

imprisonment or both under applicable law.

2. Signature

4. Typed Name

Product Manager

17- aug - 07

င်္ဂ (Ståmped)ေ့

CCCCC 0 0 0 0 0 0 0 0



17 August 2007

9/17

Mr. George LaRocca
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
One Potomac Yard
2777 S. Crystal City
Arlington, VA 22202

Subject:

Notification of Optional or Alternate Language Wisdom Lawn Granular (EPA Reg. No.: 5481-521)

Dear George,

This is being sent to notify the Agency of a minor label addition to our Wisdom Lawn Granular Insecticide. We would like to add an optional or alternate recommended spreader setting table to allow for a wider variety of spreader makes and models. Please note that all the spreaders listed are calibrated to be consistent with our label rates and use directions.

In support of this submission the following documents have been enclosed:

- Registration Application (Form 8570-1)
- One copy of the updated Wisdom Lawn Granular labeling. (Ref. No. 521-0700817r5)

If you have questions or require additional information, please do not hesitate to contact me at 949-221-6108 or kailam@amvac-chemical.com. Thank you for your time and attention with this submission.

Best regards,

Kaila Moran

Product Manager

4061 notification

WISDOMTM LAWN GRANULAR INSECTICIDE

SEP 17 2007

Only for sale to, use, and storage by commercial applicators for use to control Ants (including Imported Fire Ants), Mole Crickets and other insect pests on lawns in landscaped areas and perimeters around residential, institutional, public, commercial and industrial buildings, parks, recreational areas, and athletic fields.

Optional or Alternate Wording: Treats up to 5,000 sq. ft.

ACTIVE INGREDIENT:	By Weight
Bifenthrin*	
INERT INGREDIENTS	99.8%
TOTAL	100.0%

^{*}Cis isomers 97% minimum, trans isomers 3% maximum.

KEEP OUT OF REACH OF CHILDREN

CAUTION

•	FIRST AID						
If on skin or	Take off contaminated clothing.						
clothing:	lothing: Rinse skin immediately with plenty of water for 15-20 minutes.						
	• Call a poison control center or doctor for treatment advice.						
	EMERGENCY INFORMATION						
Have the product	container or label with you when calling a poison control center or doctor, or going for						
treatment. FOR T	HE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY:						
Transportation: C	HEMTREC1-800-424-9300						
Other: AMVAC							
	NOTE TO PHYSICIAN						
-	byrethroid. If large amounts have been ingested, milk, cream and other digestible fats and orption and so should be avoided.	doils					

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE.

EPA Reg. No. 5481-521 EPA Est. No. 39578-TX-1 **Net Contents:**

As Marked on Container

Made in U.S.A.



1-323-264-3910 • www.amvac-chemical.com

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climate modification and being grown in interior plantscapes, ornamental gardens or parks, or lawns or grounds.

Do not apply more than 0.2 lb. active ingredient (a.i.) per acre per application on residential use sites (i.e., around private homes, apartment buildings, condominiums, non-agricultural outbuildings, non-commercial greenhouses, preschools or day care facilities). May be applied at up to 0.4 lb. a.i. per acre per application on non-residential use sites (i.e., around institutional, public, commercial or industrial buildings; parks; recreational areas or athletic fields).

Do not broadcast apply more than 0.4 lbs. a.i. (200 lbs. WISDOM LAWN GRANULAR INSECTICIDE) per acre per year.

Not for use on golf courses, sod farms, nurseries, in commercial greenhouses or on grass grown for seed.

LAWN AND PERIMETER APPLICATIONS

LAWN APPLICATION: Broadcast WISDOM LAWN GRANULAR INSECTICIDE with suitable application equipment to ensure uniform coverage over the treatment area.

APPLICATION RATES

The application rates listed in the following table will provide excellent control of the respective pests under typical conditions. However, at the discretion of the applicator, WISDOM LAWN GRANULAR INSECTICIDE may be applied at up to 0.4 lb a.i./Acre to control each of the pests listed in this Table. For residential lawn use, do not apply more than 0.2 lbs. a.i. per acre per application.

PEST ACTIVE INGREDIENT LBS. PER ACRE		APPLICATION RATE WISDOM LAWN GRANULAR INSECTICIDE				
Armyworms ¹ Cutworms ¹ Sod Webworm ¹	0.1 lbs. a.i./A	50 lbs. per acre	1.15 lbs. per 1000 sq. ft.			
Annual Bluegrass Weevil (Hyperodes) (Adult) ² Billbugs (Adult) ³ Black Turfgrass Ataenius (Adult) ⁴ Leafhoppers Mealybugs	0.1 – 0.2 lbs. a.i./A	50 - 100 lbs. per acre	1.15 – 2.3 lbs. per 1000 sq. ft.			
European Crane Fly	0.15 – 0.3 lbs. a.i./A	75 – 150 lbs. per acre	1.75 – 3.50 lbs. per 1000 sq. ft.			
Ants Centipedes Chinch Bugs ⁵ Fleas (Larvae) ⁶ Fleas (Adults) Imported Fire Ants (Adult) ⁷ Imported Fire Ants (Mounds) ⁷ Millipedes Mole Cricket (Adult) ⁸ Mole Cricket (Nymph) ⁹	0.2 – 0.4 lbs. a.i./A	100 – 200 lbs. per acre	2.3 – 4.6 lbs. per 1000 sq. ft.			
Pillbugs Sowbugs	0.4 lbs. a.i./A	200 lbs. per acre	4.6 lbs. per 1000 sq. ft.			

In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of Wisdom Lawn Granular Insecticide if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

COMMENTS

¹Armyworms, Cutworms, and Sod Webworms: To ensure optimum control, irrigate the treated area with up to 0.1 inches of water immediately after application to activate (release from the granule) the insecticide.

²Annual Bluegrass Weevil (*Hyperodes*) adults: Applications should be timed to control adult weevils as they leave their overwintering sites and move into grass areas. This movement generally begins when *Forsythia* is in full bloom and concludes when flowering dogwood is in full bloom. Consult your State Cooperative Extension Service for more specific information regarding application timing.

³Billbugs: Applications should be made when adult billbugs are first observed during April and May. Degree day models have been developed to optimize application timing. Consult your State Cooperative Extension Service for information specific to your region. In temperate regions, spring applications targeting billbug adults will also provide control of over-wintered chinch bugs.

⁴Black Turfgrass Ataenius adults: Applications should be made during May and July to control the first and second generation of black turfgrass ataenius adults, respectively. The May application should be timed to coincide with the full bloom stage of Vanhoutte spiraea (*Spiraea vanhouttei*) and horse chestnut (*Aesculus hippocastanum*). The July application should be timed to coincide with the blooming of Rose of Sharon (*Hibiscus syriacus*).

⁵Chinch Bugs: Chinch bugs infest the base of grass plants and are often found in the thatch layer. Irrigate the treated area with up to 0.25 inches of water immediately after application to activate (release from the granule) the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher applications rates may be required to control populations that contain both nymphs and adults during the summer.

⁶Flea larvae: Flea larvae develop in the soil of shaded areas that are accessible to pets or other animals. Irrigate the treated area with up to 0.5 inches of water immediately after application to activate (release from the granule) the insecticide.

⁷Imported Fire ants: Control will be optimized by combining broadcast applications that will control foraging workers and newly mated fly-in queens with mound drenches that will control existing colonies. If the soil is not moist, then it is important to irrigate before application. Broadcast treatments should apply up to 0.4 lb a.i./A. Mounds should be treated by diluting 1 teaspoon of Wisdom TC Flowable (EPA Reg. No. 5481-520) formulation per gallon of water and applying 1 to 2 gallons of finished spray per mound. The mounds should be treated with sufficient force to break their apex and allow the insecticide solution to flow into the ant tunnels. A four-foot diameter circle around the mound should also be treated. For best results, apply in cool weather $(65 - 80^{\circ} \text{ F})$ or in early morning or late evening hours.

Note: A spray rig that is calibrated to apply 0.1 lb a.i./A (20 ozs.) of Wisdom TC Flowable (EPA Reg. No. 5481-520) in 2.5 gallons per 1,000 square feet contains the approximate dilution (1 teaspoon per gallon) that is required for fire ant mound drenches in the spray tank. *Application of 125 lbs/acre of WISDOM LAWN GRANULAR INSECTICIDE will provide six months residual activity for control of foraging imported Fire Ants and newly mated fly-in queens. *Not applicable in California.

⁸Mole Cricket adults: Achieving acceptable control of adult mole crickets is difficult because preferred grass areas are subject to continuous invasion during the early spring by this extremely active stage. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be maximized. Grass areas that receive pressure from adult mole crickets should be treated at peak egg hatch to ensure optimum control of subsequent nymph populations (see below).

⁹Mole Cricket nymphs: Grass areas that received intense adult mole cricket pressure in the spring should be treated immediately prior to peak egg hatch. Optimal control is achieved at this time because young nymphs are more susceptible to insecticides and they are located near the soil surface where the insecticide is most concentrated. Control of larger, more damaging, nymphs later in the year may require both higher application rates and more frequent applications to maintain acceptable control Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be maximized.

PERIMETER APPLICATION

Apply to ground area 5 to 10 feet wide around and adjacent to the structure.

Boundary Applications: Apply to soil areas, ornamental gardens, pavement crevices, fencelines, planter areas, vegetation areas and mulch areas.

80911

Appl. Rate: Pounds Per 1,000	Band Width In Feet	Pounds of WISDOM LAWN GRANULAR INSECTICIDE for Perimeter Applications Linear Distance (Feet) Around Structure							
sq. ft.		100	150	200	250	300	400	500	1,000
2.3	5	1.3	1.9	2.5	3.0	3.6	4.8	5.9	11.7
2.3	10	3.0	4.2	5.3	6.5	7.6	9.9	12.2	23.7
3.0	5	1.7	2.5	3.3	3.9	4.7	6.3	7.7	15.2
3.0	10 .	3.9	5.4	6.9	8.5	9.9	12.9	15.9	30.9
3.5	5	2.0	2.9	3.8	4.5	5.4	7.2	8.9	17.5
3.5	10	4.5	6.3	8.0	9.8	11.4	14.9	18.3	35.5
4.6	5	2.6	3.8	5.0	6.0	7.2	9.6	11.8	23.4
4.6	10	6.0	8.4	10.6	13.0	15.2	19.8	24.4	47.4

NOTE: The amounts of WISDOM LAWN GRANULAR INSECTICIDE listed in the above table are based on the total area being treated. This area is determined by multiplying the linear distance by the band width and then adding to that value the area of a circle whose radius is equal to the band width (to account for the area treated at the four corners of a rectangular structure).

FIRE ANT MOUND TREATMENT: Apply 1/2 cup of WISDOM LAWN GRANULAR INSECTICIDE per mound and then drench the mound with 1 to 2 gallons of water. The mound should be drenched with sufficient force to break its apex and allow the insecticide to flow into the ant tunnels. For best results, apply in cool weather (65 to 80° F) or in early morning or late evening hours. Treat new mounds as they appear. Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with insecticide.

INDUSTRIAL NON-AGRICULTURAL APPLICATIONS

(Including Industrial Sites, Parks, Ball Fields and Cemeteries)

BROADCAST SPOT APPLICATIONS: For the control of actively foraging ants (including Imported Fire Ants) broadcast uniformly 2.3 to 4.6 lbs. WISDOM LAWN GRANULAR INSECTICIDE per 1000 sq. ft. (0.2 to 0.4 lbs. a.i./Acre). See comments section under Lawn & Perimeter Applications for additional rate recommendations.

PAD MOUNTED ELECTRICAL SITES: To reduce or prevent ant mound building activity on electrical pads, uniformly distribute 1.0 to 1.5 Tablespoons (0.06 to 0.09 lbs.) of WISDOM LAWN GRANULAR INSECTICIDE per 4 sq. ft. of mounting pad. Where existing transformers are completely or partially filled with soil excavated by resident Ants, remove any excess fill and level the remaining soil prior to application.

NOTE: Electrical transformers appear to be more attractive to ants during the cooler months. For best results, treat in the Fall or early Spring.

WISDOM LAWN GRANULAR INSECTICIDE - Recommended Spreader Settings for Broadcast Applications

EQUIPMENT	GROUND	SPREADER SETTING						
	SPEED (MPH)	100 POUND	S PER ACRE	200 POUNDS PER ACRE				
	,	ONCE OVER	TWICE OVER	ONCE OVER	TWICE OVER			
Hand-Pushed								
LESCO Rotary	3	B + 1/4	A + 1/2	C + 1/2	B + 1/4			
Scotts R-8A-1 Rotary	3	G	E	Not Rec.	G			
Gandy Models (42, 36, 30, 24)	3	12	10.5	16	12			
Spyker (all Models)	3	2.6	2	3.1	2.6			
Equipment Drawn	(PTO=540 RP	PM)						
Gandy Models (42, 36, 30, 24)	4.5	14	11.5	19	14			
Gandy 6500 Series	4.5	15	11	20	15			
Gandy 10T Series	5.0	16.5	12	24	16.5			
Herd GT-77	4.0	-		8-H	-			
Herd GT-77	8.0	8-H		_	8-H			
Lely 2.3202.000.1	4.5	2	1	3.5	2			
Lely Model L- 400	4.5	Not Rec.	Not Rec.	2	Not Rec.			
Vicon PS402	4.5	10	7	16	10			
Vicon PS1002T	4.5	13	9	21	13			

Optional or Alternate Spreader Settings

Recommended Spreader Settings for Broadcast Applications							
Sangalar Maka and Madal	Setting To Deliver						
Spreader Make and Model	2.3 lbs. Per 1,000 sq. ft.	3.5 lbs. Per 1,000 sq. ft.					
Spyker (all modėls)	4.7	5					
Earthway Drop 7300B	16	17					
Republic/Scotts Broadcast	3 -	4					
Republic/ Scotts Drop	. 6	7					
Precision Drop FH-22D	9	10					
Precision Broadcast/Red Devil	12	13					
Scott Speedy Green	3	4.5					
Broadcast/Earthway 2850	14	15					
Broadcast/Earthway 2800A	14	15					
Broadcast/Earthway 2050 S.U.	14	15					

NOTE: These recommended settings should be used as starting points when calibrating a spreader to achieve the desired application rate of WISDOM LAWN GRANULAR INSECTICIDE.

After calibration, the technician should regularly observe the amount of WISDOM LAWN GRANULAR INSECTICIDE that is applied to a known area and make further setting adjustments, as necessary, to maintain the appropriate application rate. Spreaders should be calibrated periodically to ensure that equipment wear or other

100f11

factors have not altered the flow rate of WISDOM LAWN GRANULAR INSECTICIDE. Calibration instructions are included in the Owner's Manual that was provided by the equipment manufacturer at the time of purchase.

RESISTANCE: Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state pest management authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and suspect that resistance is a reasonable cause, immediately consult your local company representative or pest management advisor for the best alternative method of control for your area.

Dealers Should Sell in Original Packages Only.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use close tightly.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call Chemtrec 1-800-424-9300.

To confine spill: Cover to prevent dispersal. Place damaged package in a holding container. Identify contents. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Paper and plastic bags: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use, subject to the inherent risks referred to herein, when it is used in accordance with such directions; and (c) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions.

THERE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE FULLEST EXTENT PERMITTED BY LAW MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

AMVAC offers this product, and Buyer accepts it, subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC.

Wisdom is a trademark of AMVAC Chemical Corporation.

AMVAC Chemical Corporation 4100 E. Washington Boulevard Los Angeles, CA 90023 U.S.A.

110011

1-323-264-3910 www.amvac-chemical.com

NOTIFICATION
SEP 17 2007