



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

September 24, 2025

Jordan Moseley
Regulatory Affairs Manager
AMVAC Chemical Corporation
4695 MacArthur Court, Suite 1200
Newport Beach, CA 92660

Subject: Label Amendment - Registration Review Mitigation for Bifenthrin
Product Name: SmartChoice 5G
EPA Registration Number: 5481-561
Case Number: N/A
Application Date: March 24, 2021

Dear Jordan Moseley:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Bifenthrin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for

shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Meaghan Robertson by phone at 202-566-0169, or via email at robertson.meaghan@epa.gov.

Sincerely,

A handwritten signature in blue ink that reads "Cathryn Britton". The signature is fluid and cursive, with the first name and last name clearly distinguishable.

Cathryn Britton
Chief, Risk Management and Implementation
Branch V
Pesticide Re-evaluation Division (7508M)
Office of Pesticide Programs

ENCLOSURE: Stamped label

RESTRICTED USE PESTICIDE

DUE TO ACUTE HUMAN, AVIAN, AND AQUATIC INVERTEBRATE TOXICITY

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification

ACCEPTED**09/24/2025**Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under

EPA Reg. No. 5481-561

Chlorethoxyfos	GROUP	1B	INSECTICIDE
Bifenthrin	GROUP	3A	INSECTICIDE

SMARTCHOICE™ 5G**granular insecticide***For Use On Corn Only*Alternate Wording when packaged in SMARTBOX: Available in the SMARTBOX®, a Closed Handling and Application System.Alternate Wording when packaged in LOCK 'N LOAD: Available in LOCK 'N LOAD®, a Closed Loading System.TO CONTROL CORN ROOTWORM LARVAE, WIREWORMS, CUTWORMS, SEEDCORN MAGGOTS, WHITE GRUBS,
GRAPE COLASPIS AND SYMPHYLANS ON FIELD CORN, SEED CORN, SWEET CORN, AND POPCORN.**ACTIVE INGREDIENT: (% by weight)**

Chlorethoxyfos:

Phosphorothioic acid, O,O-diethyl O-(1,2,2,2-tetrachloroethyl)ester.....4.3%

Bifenthrin:

(2-methyl [1,1'-biphenyl]-3-yl)methyl 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-
2,2-dimethylcyclo-propanecarboxylate.....0.7%**INERT INGREDIENTS:** 95.0%**TOTAL:** 100.0%**STOP! READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN****DANGER  PELIGRO
POISON**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)**FIRST AID**

If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

EMERGENCY INFORMATION

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY:

For Medical Emergencies phone:.....1-888-681-4261

For Transportation Emergencies, including spill, leak or fire, phone: CHEMTREC.....1-800-424-9300

For Product Use Information phone : AMVAC.....1-888-462-6822

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE.

EPA Reg. No. 5481-561

NET WEIGHT: 50 LBS.

EPA Est. No. _____

AMVAC Chemical Corporation
4695 MacArthur Court, Suite 1200
Newport Beach, CA 92660 U.S.A.

NOTE TO PHYSICIAN

Organophosphate and Pyrethroid

ATROPINE IS THE EMERGENCY ANTIDOTE FOR SMARTCHOICE INSECTICIDE POISONING. 2-PAM IS ALSO ANTIDOTAL AND MAY BE USED IN CONJUNCTION WITH ATROPINE.

Warning Symptoms

SMARTCHOICE is an organophosphate insecticide containing a pyrethroid and is a cholinesterase inhibitor. Overexposure to this product may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system.

Symptoms of poisoning may include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, muscle tremors and abdominal cramps. In severe cases, convulsions, unconsciousness and respiratory failure may occur.

Treatment

Atropine is the specific therapeutic antagonist of choice against this type of parasympathetic nervous stimulation. If warning symptoms appear, treatment should consist of parenteral atropine sulfate. This can be administered intramuscularly every 10 minutes until full atropinization (as indicated by dilated pupils, dry flushed skin and tachycardia) has been achieved and repeated thereafter whenever symptoms reappear. For infants and children, the dose of atropine sulfate is 0.05 mg/kg. For adults, 1 to 2 milligrams will be required for mild cases, but up to 4 mg may be needed for severe intoxication. 20 to 30 milligrams or more may be required during the first 24 hours.

Pralidoxime chloride (2-PAM chloride) may also be used as an effective antidote in addition to and while maintaining full atropinization. In adults, an initial dose of 1 gram of 2-PAM should be injected, preferably as an infusion in 250 cc of saline over a 15- to 30-minute period. If this is not practical, 2-PAM may be administered slowly by intravenous injection as a 5 percent solution in water over not less than two minutes. After about an hour, a second dose of 1 gram of 2-PAM will be indicated if muscle weakness has not been relieved. For infants and children, the dose of 2-PAM is 20-50 mg/kg. Morphine is an improper treatment.

Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase regeneration has been attained as determined by blood test. Reversible skin sensations (parasthesia) may occur and ordinary skin salves have been found useful in reducing discomfort.

For medical emergencies involving this product, call 1-888-681-4261.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Fatal if swallowed. May be fatal if inhaled. Causes moderate eye irritation. Harmful if absorbed through the skin. Do not breathe dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

Repeated exposure can lead to an asymptomatic increase in the susceptibility to poisoning. Avoid prolonged exposure to the product.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves (except for applicators) such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, Polyvinyl Chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. All washable PPE should be washed in detergent and hot water prior to reuse. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

IMPORTANT: The SMARTBOX (*Alternate Wording when packaged in LOCK 'N LOAD*: LOCK 'N LOAD) system when used correctly qualifies as a closed loading system under the WPS. Loaders using the SMARTBOX (*Alternate Wording when packaged in LOCK 'N LOAD*: LOCK 'N LOAD) system must:

- Wear the PPE specified above for loaders
- In addition to wearing the required PPE, have immediately available for use in case of an emergency
- Chemical-resistant apron
- Chemical-resistant footwear, and

Wear a minimum of a NIOSH-approved particulate filtering faceplate respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters. Applicators must be in enclosed cabs and must:

- Wear the PPE specified above for applicators
- In addition to wearing the required PPE, have available for use coveralls, chemical-resistant gloves, and protective eyewear if it is necessary to exit the cab and contact pesticide-treated surfaces in the treated area
- Remove PPE that was worn in the treated area before reentering the cab, and
- Store all PPE in a chemical-resistant container, such as a plastic bag, to prevent contamination of the inside of the cab.

USER SAFETY RECOMMENDATIONS

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to wild mammals, birds, fish, and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label

about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours. The REI is 72 hours where average rainfall is less than 25 inches per year. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Double Notification: Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, Polyvinyl Chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils
- Shoes plus socks.

IMPORTANT: Apply this product only as specified on this label. Do not apply this product through any type of irrigation system.

GENERAL INFORMATION

SMARTCHOICE 5G is a granular soil insecticide for use on corn (field, sweet, popcorn and seed) at planting for the control of northern, western and southern corn rootworm larvae, wireworms, cutworms, seedcorn maggots, white grubs, grape colaspis and symphylans.

Herbicide Compatibility Information

- All registered corn herbicides may be applied to SMARTCHOICE-treated corn.

Application Instructions

Crop	Pest	Dosage and Comments
Field Corn Seed Corn Sweet Corn Popcorn	Cutworms Wireworms White Grubs Seedcorn Maggots Grape Colaspis Symphylans	3.0 to 3.5 ounces per 1,000 feet of row
	Corn Rootworm Larvae* Northern Western Southern	*Aids in the control of corn rootworm larvae when used in combination with corn hybrids containing in-plant protection against corn rootworm larvae
	Corn Rootworm Larvae Northern Western Southern Cutworms Wireworms White Grubs Seedcorn Maggots Grape Colaspis Symphylans	4.5 to 5.0 ounces per 1,000 feet of row
Restrictions:		

Do not make more than one application per year.

Crop Rotational Intervals

Corn.....Anytime
Other crops.....30 days

Application

SMARTCHOICE 5G should be applied in-furrow at planting for optimum control of labeled pests. DO NOT apply as a surface band behind the press wheel.

In-Furrow Application at Planting:

Apply SMARTCHOICE 5G in conventional, conservation or no-till corn as an in-furrow treatment. Place the applicator tube either directly into or behind the planter shoe, so the granules drop into the open seed furrow and are completely covered with soil. In-furrow application provides optimum control of corn rootworms, wireworms, white grubs, seedcorn maggots, and grape colaspis.

Further Application Information:

- Under windy conditions the use of wind shields is recommended to ensure proper placement of granules.
- The SMARTBOX Alternate Wording when packaged in LOCK 'N LOAD: LOCK 'N LOAD system must be properly calibrated to assure an accurate application rate of SMARTCHOICE 5G.
- SMARTCHOICE 5G is non-corrosive to equipment and is nonflammable.

Determining the Amount of SMARTCHOICE 5G per Acre

To determine the approximate amount of SMARTCHOICE 5G needed per acre, select the use rate per 1,000 row feed and the row spacing as shown in the following chart:

Amount of SMARTCHOICE per 1,000 feet of row	Pounds of SMARTCHOICE 5G Required per Acre at Various Row Spacings						
	15 ' or twin row	20 in	22 in	30 in	34 in	36 in	40 in
3.0 oz.	6.5	4.9	4.5	3.3	2.9	2.8	2.6
3.5 oz.	7.6	5.7	5.2	3.8	3.4	3.2	3.0
4.5 oz.	9.8	7.3	6.7	4.9	4.3	4.1	3.9
5.0 oz.	10.9	8.2	7.4	5.4	4.8	4.5	4.3

Alternate Wording when packaged in LOCK 'N LOAD:

SMARTCHOICE 5G CALIBRATION GUIDELINES

The following **equipment calibration procedures** must be performed to ensure that the proper amount of SMARTCHOICE 5G will be applied:

1. Fill insecticide hoppers with SMARTCHOICE 5G using Lock 'N Load containers.
2. Attach a collection container to each tube opening to catch the insecticide.
3. Set your planter to the initial gauge settings shown in the chart.
4. Measure off 250 ft of row and drive planter, while metering the insecticide, the premeasured distance at the speed at which you will be planting.
5. The target catch weight for each container is 0.88 and 1.25 oz of granules for the rates of 3.5 oz and 5.0 oz per 1000 ft, respectively, when collected for 250 ft of row. Determine the amount caught in each row either by weight or by volumetric approximation of weight.
6. Repeat the calibration procedure above, as needed, until the insecticide caught in each container is as close as possible to the target amount.

Planting Speed MPH	4		5		6		7		8	
Application Rate (oz per 1000 row ft)	3.5	5.0	3.5	5.0	3.5	5.0	3.5	5.0	3.5	5.0
Planter / Metering System										
Case International Harvester (One guage per hopper; Gate/Dial) - Older	1/5	1/7	1/7	1/9	1/8	2/1	1/9	2/3	2/1	2/5
Case International Harvester (Two guages per hopper; (Gate/Dial) - Newer	1/1	1/3	1/3	1/5	1/4	1/6	1/5	1/7	1/6	1/9
John Deere MaxEmerge® 2 (Models 7200/7300) & 1760 Conservation Planter	12	15	14	17	16	20	18	24	20	27
John Deere 7000 MaxEmerge (Even) (0-10-20-30-40)	8	11	10	13	11	14	13	16	15	18
John Deere 7000 MaxEmerge (Odd) (5-15-25-35-45)	7	9	9	11	11	13	13	15	14	16
John Deere MaxEmerge Plus	11	14	13	16	15	19	17	23	19	26
Kinze (2600 Series & Newer)	7	10	9	12	10	13	12	15	14	17
White (5100 Series)	7	8	8	9	9	11	10	12	12	14
White (6100 Series)	8	10	10	12	12	14	13	15	15	17

NOTE: These settings should be used as **starting points only**. Follow manufacturers recommendations for calibration of the specific equipment you are using or the calibration guidelines above to make final adjustments. After initial calibration, recheck flow rate occasionally to make sure equipment wear, changing conditions, etc. have not caused a change in flow rate.

Alternate Wording when packaged in SMARTBOX: SMARTBOX INFORMATION

SMARTCHOICE 5G must be applied with the SMARTBOX system. Please refer to the SMARTBOX operator's manual for detailed information on system operation and calibration procedures.

Alternate Wording when packaged in SMARTBOX: DO NOT ATTEMPT TO OPEN OR TAMPER WITH THE CONTAINER. DO NOT ATTEMPT TO REPAIR THE SMARTBOX APPLICATION SYSTEM. IF REPAIRS ARE NECESSARY, CONTACT YOUR AMVAC SALES REPRESENTATIVE OR THE MANUFACTURER OF THE APPLICATION SYSTEM.

VEGETATIVE FILTER STRIPS

Construct and maintain a vegetative filter strip, according to the width specified below, of grass or other permanent vegetation between the field edge and nearby down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing bifenthrin onto fields where a maintained vegetative filter strip of at least 25 feet exists between the field edge and where a down gradient aquatic habitat exists. This minimum required width of 25 feet may be reduced or removed under the following conditions:

- For Western irrigated agriculture, a maintained vegetative filter strip of at least 10 feet wide is required. Western irrigated agriculture is defined as irrigated farmland in the following states: WA, OR, CA, ID, NV, UT, AZ, MT, WY, CO, NM, and TX (west of I-35).
 - o For Western irrigated agriculture, if a sediment control basin is present, a vegetative filter strip is not required.
- In all other areas, a vegetative filter strip with a minimum width of 25 feet is required, unless the following conditions are met. The vegetative filter strip requirement may be reduced from 25 feet to 15 feet if at least one of the following applies:

- o The area of application is considered prime farmland (as defined in 7 CFR § 657.5).
- o Conservation tillage is being implemented on the area of application. Conservation tillage is defined as any system that leaves at least 30% of the soil surface covered by residue after planting. Conservation tillage practices can include mulch-till, no-till, or strip-till.
- o A functional terrace system is maintained on the area of application.
- o Water and sediment control basins for the area of application are functional and maintained.
- o The area of application is less than or equal to 10 acres.

For further guidance on vegetative filter strips, refer to the following publication for information on constructing and maintaining effective buffers: Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. <https://www.regulations.gov/document?D=EPA-HQ-OPP-2008-0331-0175>

Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marches, ponds, estuaries, and commercial fish ponds).

Resistance Management

For resistance management, please note that SMARTCHOICE 5G contains both Group 1B (chlorothoxyfos) and Group 3A (bifenthrin) insecticides. Any insect population may contain individuals naturally resistant to SMARTCHOICE 5G and other Group 1 or Group 3 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same locations. Appropriate resistance management strategies should be followed.

To delay insecticide resistance, take one or more of the following steps:

- Rotate the use of SMARTCHOICE 5G or other Group 1 or Group 3 insecticides within the year with different groups that control the same pests.
- Use combinations with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same combination repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of the mixture. In addition, consider the following recommendations from the Insect resistance Action Committee (IRAC):
 - o Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - o Combinations with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - o When using combinations, consider any known cross-resistance issues between the individual components for the targeted pests.
 - o Combinations become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - o The insect resistance management benefits of an insecticide combination are greater if the two components have similar periods of residual insecticidal activity. Combinations of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated insect management program for insecticide use that includes scouting, uses historical information related to pesticide use record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or pest control advisor for any additional pesticide resistance-management and/or IPM recommendations for specific site and pest problems in your area.

For further information or to report suspected resistance contact AMVAC at (1-888-462-6822). You can also contact your pesticide distributor or university extension specialist to report resistance.

STORAGE AND DISPOSAL

STORAGE: Store in a secure, locked, dry and temperate area where children, unauthorized persons or animals cannot enter. Keep the container capped when not in use. Do not transport or store near food or feed intended for human or animal consumption. Do not use or store in or around the home. Do not contaminate water, other pesticides, fertilizer, food or feed by storage or disposal.

PRODUCT DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide or container is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions contact our State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

Alternate Wording when packaged in SMARTBOX:

DO NOT DISCARD THE CONTAINER. Do not attempt to open or tamper with the container. Completely empty container into application equipment. All SMARTBOX containers must be returned per instructions provided or by calling 1-888-SMARTBOX. Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Alternate Wording when packaged in LOCK 'N LOAD:

DO NOT DISCARD THE CONTAINER. Do not attempt to open or tamper with the container. Completely empty container into application equipment. Return empty container to point of purchase or call 1-888-GO-AMVAC for instructions. Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Alternate Wording when packaged in Super Sacks:

Nonrefillable container. Do not reuse or refill this container. Completely empty bag into packaging equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use, subject to the inherent risks referred to herein, when it is used in accordance with such directions; and (c) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions.

THERE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

AMVAC offers this product, and Buyer accepts it, subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC.

SMARTCHOICE is a trademark and SMARTBOX and LOCK 'N LOAD are registered trademarks of Amvac Chemical Corporation. MaxEmerge is a registered trademark of Deere and Company.

AMVAC Chemical Corporation
4695 MacArthur Court, Suite 1200
Newport Beach, CA 92660 U.S.A.
1-888-462-6822