



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

June 1, 2017

Jon C. Wood
Sr. Regulatory Manager
Amvac Chemical Corporation
4695 MacArthur Court, Suite 1200
Newport Beach, CA 92660

Subject: Label Amendment – Revising directions for use and updating the trademark
Product Name: Scepter 70 DG Herbicide
EPA Registration Number: 5481-610
Application Date: 3/31/2017
Decision Number: 528011

Dear Jon C. Wood:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 5481-610
Decision No. 528011

with FIFRA section 6. If you have any questions, please contact Nathan Mellor by phone at 703-347-8562, or via email at mellor.nathan@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Shaja Joyner".

Shaja Joyner, Product Manager 20
Fungicide and Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

Group	2	Herbicide
-------	---	-----------



ACCEPTED
06/01/2017
Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 5481-610

FOR USE IN SOYBEANS

Active Ingredient:

Imazaquin (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinoline-carboxylic acid)..... 70.0%
Other Ingredients..... 30.0%
TOTAL100.0%

EPA Reg. No. 5481-610 EPA Est. No. 39578-TX-1

**KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. In case of an emergency endangering life or property involving this product, call AMVAC 1-888-462-6822 or Chemtrec 1-800-424-9300.	

See inside for complete **Precautionary Statements, Directions For Use, and Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

In case of an emergency endangering life or property involving this product, call AMVAC® 1-888-462-6822 or Chemtrec® 1-800-424-9300.

Product of U.S.A.

Net Contents: 8.75

pounds



AMVAC Chemical Corporation
4100 E. Washington Blvd.
Los Angeles, CA 90023 U.S.A
1-888-462-6822

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION!

Causes eye irritation. Avoid contact with skin, eyes, or clothing.
Harmful if absorbed through skin.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

For terrestrial uses: DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Observe all cautions and limitations on this label and on the labels of products used in combination with **Scepter® 70 DG herbicide**. **DO NOT** use **Scepter 70 DG** other than in accordance with the instructions set forth on this label. The use of **Scepter 70 DG** not consistent with this label may result in injury to crops. Keep containers closed to avoid spills and contamination.

DO NOT apply this product through any type of irrigation system.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves made of any waterproof material
- Shoes plus socks

PRODUCT INFORMATION

Scepter® 70 DG herbicide is a flexible broad-spectrum herbicide which may be applied preplant incorporated, preemergence or postemergence in soybeans.

When **Scepter 70 DG** is applied to the soil, susceptible weeds may emerge but, growth stops, and the weeds either die or are not competitive with the crop.

When **Scepter 70 DG** is applied postemergence, absorption occurs through both the foliage and roots. Susceptible weeds stop growing and either die or are not competitive with the crop.

A timely cultivation may aid in the control of certain weeds or improve weed control when adequate moisture is not received after application. Cultivation should be shallow. After postemergence treatments, wait at least 10 days before cultivating.

Scepter 70 DG is absorbed by weed roots and/or foliage and is rapidly translocated to growing points. Therefore, adequate soil moisture is necessary for optimum **Scepter 70 DG** activity. The amount of rainfall or irrigation required following application depends on existing soil moisture, soil texture and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is normally adequate. If adequate moisture is not received within 7 days after treatment, then a cultivation or postemergence herbicide application can improve weed control. When adequate moisture is received after dry conditions, **Scepter 70 DG** will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and weed size.

Occasionally, internode shortening of soybean plants may occur from **Scepter 70 DG** applications. The internode shortening does not impact soybean yields.

Some of the weeds listed on this label have populations with evolved resistance to HG 2 (ALS/AHAS inhibiting) herbicides. If HG 2 resistant biotypes are present in a field, tank-mix or apply **Scepter 70 DG** sequentially with an alternate registered herbicide having an effective mode of action to ensure control. Consider including cultural and mechanical tactics to supplement herbicides when herbicide-resistant weed biotypes are present in fields.

See your AMVAC representative for additional information.

PRECAUTIONS

Crops other than soybeans, such as cotton, corn and vegetables, may be injured by spray drift or other contact with **Scepter 70 DG**.

To avoid injury to sensitive crops from spray drift, follow all use directions and precautions in **SPRAYING INSTRUCTIONS** section.

To avoid injury to sensitive crops, spray equipment used for **Scepter 70 DG** applications must be drained and thoroughly cleaned with water before being used to apply other products to these crops.

Use of **Scepter 70 DG** herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is possible. See **ROTATIONAL CROP RESTRICTIONS** section.

Replanting: If replanting is necessary in a field previously treated with **Scepter 70 DG**, the field may be replanted to soybeans at any time. Rework the soil no deeper than the treated zone. **DO NOT** apply a second treatment of **Scepter 70 DG**.

RESTRICTIONS

DO NOT use **Scepter 70 DG** other than in accordance with the instructions set forth on this label or approved supplemental labels.

DO NOT apply this product more than once per year to soybeans.


DO NOT apply more than 2.8 ounces of **Scepter 70 DG** per acre per year.

There must be an interval of at least 90 days between **Scepter 70 DG** application and soybean harvest.

DO NOT graze or feed treated soybean forage, hay or straw to livestock.

USE AREA

For use in soybeans, **Scepter® 70 DG herbicide** can be applied only in the states or parts of states shaded in the following map:



USE REGION 1 includes eastern Oklahoma (east of I-35), Arkansas, the Missouri bootheel, Tennessee, North Carolina, South Carolina, Georgia, Florida, Alabama, Mississippi, Louisiana, and eastern Texas (east of I-35 north of San Antonio, east of I-37 south of San Antonio).

USE REGION 2 includes eastern Kansas (east of U.S. 81; the counties of Cloud, Ellsworth, Harvey, Jewell, Lincoln, Mitchell, McPherson, Ottawa, Republic, Saline, Sedgewick, and Sumner), southeastern Nebraska (east of U.S. 81, south of U.S. 34), Missouri, Illinois (south of S.R. 116 west of Peoria, south of U.S. 24 east of Peoria), Indiana, Ohio, Kentucky, Virginia, West Virginia, Pennsylvania, Maryland, Delaware, and New Jersey.

USE REGION 3 includes Nebraska (east of U.S. 81, north of U.S. 34 and also that area east of U.S. 283, south of U.S. 30, and west of U.S. 81), South Dakota (east of U.S. 81), Illinois (north of S.R. 116 west of Peoria; north of U.S. 24 east of Peoria), Wisconsin, Iowa, Minnesota (south of S.R. 210), and Michigan.

NOTE: See the **ROTATIONAL CROP RESTRICTIONS** section for directions applying to each USE REGION.

WEEDS CONTROLLED

WEEDS CONTROLLED BY SOIL APPLICATIONS

When applied as directed, soil-applied treatments of **Scepter® 70 DG herbicide** will control or suppress the following weeds:

WEED		LEVEL OF CONTROL
BROADLEAF WEEDS		
Alligatorweed	(<i>Alternanthera philoxeroides</i>)	Control
Beggarweed, Florida	(<i>Desmodium tortuosum</i>)	Suppression
Bristly Starbur	(<i>Acanthospermum hispidum</i>)	Control
Burcucumber	(<i>Sicyos angulatus</i>)	Control
Cocklebur*, Common	(<i>Xanthium strumarium</i>)	Control
Copperleaf, Hophornbeam	(<i>Acalypha ostryaefolia</i>)	Suppression
Jimsonweed*	(<i>Datura stramonium</i>)	Control

Lambsquarters*, Common	(<i>Chenopodium album</i>)	Control
Mallow, Venice	(<i>Hibiscus trionum</i>)	Control
Mexicanweed	(<i>Caperonia castanifolia</i>)	Suppression
Morningglory,		
Entireleaf	(<i>Ipomoea hederacea</i> var. <i>intergriuscula</i>)	Suppression
Ivyleaf	(<i>Ipomoea hederacea</i>)	Suppression
Palmleaf	(<i>Ipomoea wrightii</i>)	Control
Pitted	(<i>Ipomoea lacunose</i>)	Control
Smallflower	(<i>Jacquemontia tamnifolia</i>)	Control
Tall	(<i>Ipomoea purpurea</i>)	Suppression
Mustard species*	(<i>Brassica</i> species)	Control
Nightshade*, Eastern Black	(<i>Solanum ptycanthum</i>)	Control
Pigweed,		
Palmer*	(<i>Amaranthus palmeri</i>)	Control
Redroot*	(<i>Amaranthus retroflexus</i>)	Control
Smooth*	(<i>Amaranthus hybridus</i>)	Control
Spiny	(<i>Amaranthus spinose</i>)	Control
Poinsettia*, Wild	(<i>Euphorbia heterophylla</i>)	Control
Puncturevine	(<i>Tribulus terrestris</i>)	Control
Pusley, Florida	(<i>Richardia scabra</i>)	Control
Ragweed,		
Common*	(<i>Ambrosia artemisiifolia</i>)	Control
Giant*	(<i>Ambrosia trifida</i>)	Control
Redweed	(<i>Melochia corchorifolia</i>)	Control
Sesbania, Hemp	(<i>Sesbania exaltata</i>)	Suppression
Sicklepod (<i>Cassia obtusifolia</i>)		Control
Sida, Prickly (teaweed)*	(<i>Sida spinosa</i>)	Control
Smartweed,		
Ladysthumb*	(<i>Polygonum persicaria</i>)	Control
Pennsylvania	(<i>Polygonum pennsylvanica</i>)	Control
Spurge, Spotted	(<i>Euphorbia maculata</i>)	Suppression
Sunflower*, Common	(<i>Helianthus annuus</i>)	Control
Texasweed	(<i>Caperonia palustris</i>)	Suppression
Velvetleaf	(<i>Abutilon theophrasti</i>)	Suppression
Waterhemp*, Common/ Tall	(<i>Amaranthus tuberculatus</i> syn. <i>rudis</i>)	Control

GRASSES

Barnyardgrass*	(<i>Echinochloa crus-galli</i>)	Suppression
Corn, Volunteer	(<i>Zea mays</i>)	Suppression
Foxtail,		
Giant*	(<i>Setaria faberi</i>)	Control
Green*	(<i>Setaria viridis</i>)	Control
Yellow*	(<i>Setaria pumila/glauca</i>)	Control
Goosegrass*	(<i>Eleusine indica</i>)	Suppression
Johnsongrass*, seedling	(<i>Sorghum halepense</i>)	Control
Shattercane*	(<i>Sorghum bicolor</i>)	Suppression
Signalgrass, Broadleaf	(<i>Brachiaria platyphylla</i>)	Suppression

SEDGES

Nutsedge*, Yellow (*Cyperus esculentus*) Suppression

*Weeds that have HG 2 resistant populations based on reports to the International Survey of Herbicide Resistant Weeds (<http://weedsience.org/>)

WEEDS CONTROLLED BY POSTEMERGENCE APPLICATIONS

When applied as a postemergence treatment, **Scepter® 70 DG herbicide** will control the following weeds:

WEEDS CONTROLLED	MAXIMUM WEED HEIGHT CONTROLLED (inches)	
	1.4 ounces/acre	2.8 ounces/acre
Cocklebur*, Common (<i>Xanthium strumarium</i>)	8	12
Pigweed,		
Palmer* (<i>Amaranthus palmeri</i>)	N	6
Redroot* (<i>Amaranthus retroflexus</i>)	4	12
Smooth* (<i>Amaranthus hybridus</i>)	4	12
Spiny (<i>Amaranthus spinosus</i>)	4	12
Poinsettia*, Wild (<i>Euphorbia heterophylla</i>)	N	6
Sunflower*, Common (<i>Helianthus annuus</i>)	4	8
Volunteer Corn (<i>Zea mays</i>)	8	12

N = Not Controlled

Apply when weeds are actively growing. **DO NOT** apply **Scepter 70 DG** postemergence when soybeans and weeds have been subjected to stress conditions such as temperature and moisture extremes.

*Weeds that have HG 2 resistant populations based on reports to the International Survey of Herbicide Resistant Weeds (<http://weedsience.org/>)

DIRECTIONS FOR CONVENTIONAL, MINIMUM, AND NO-TILL APPLICATIONS

APPLICATION RATE

Apply **Scepter 70 DG** at a maximum broadcast rate of 2.8 ounces per acre for preplant incorporated and preemergence (including minimum and no-till systems) applications. Apply **Scepter 70 DG** at a broadcast rate of 1.4 to 2.8 ounces per acre for postemergence applications.

NOTE: In USE REGION 3 as defined by the **USE AREA** section of this label, apply **Scepter 70 DG** prior to July 1.

APPLICATION INSTRUCTIONS

SOIL APPLICATIONS

Scepter® 70 DG herbicide may be applied in conventional, minimum, or no-till as a preplant incorporated or preemergence application up to 45 days (30 days in USE REGION 1) before planting. Preplant incorporated applications may be applied immediately before planting or up to 30 days prior to planting in USE REGION 1. For USE REGION 2 and USE REGION 3, **Scepter 70 DG** may be applied up to 45 days prior to planting. Incorporate uniformly into the top 1 to 2 inches of soil within 7 days after application. If soybeans are planted on beds, apply and incorporate after bed formation using PTO-driven equipment or rolling cultivator.

Preemergence applications may also be applied during or after planting but before the crop emerges.

Adequate moisture is required for activation of **Scepter 70 DG**.

Tank mixes: **Scepter 70 DG** herbicide may be applied preplant incorporated or preemergence in a tank-mixture with the following products to improved residual weed control spectrum in soybeans and to better manage herbicide-resistant weed populations. Products listed for tank-mixture with **Scepter 70 DG** herbicide must be registered for preplant incorporated or preemergence application in soybean. Read and follow the label directions of all products in the tank-mixture. Tank-mixtures with **Scepter 70 DG** are not limited to the tank-mix partners

listed below. It is the pesticide user’s responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture:

Authority/Spartan (sulfentrazone), Cinch/Dual II MAGNUM/Medal/Charger Max (S-metolachlor/metolachlor), Frontier (dimethenamid-P), Warrant (acetochlor), Zidua (pyroxasulfone), Linex/Lorox (linuron), Metribuzin/Sencor/TriCor (metribuzin), Pendimax/Prowl (pendimethalin), Treflan/Trust (trifluralin), Sharpen (Saflufenacil), Valor SX/Rowel (Flumioxazin), Cadet (Fluthiacet), Aim (Carfentrazone), Resource (flumiclorac).

Glyphosate or glufosinate may be mixed with Scepter 70 DG when applied before soybean emergence.

POSTEMERGENCE APPLICATIONS

Scepter 70 DG applied postemergence controls the emerged weed species listed previously, but can also provide residual control of susceptible weeds that may emerge after application. Weeds should be small and actively growing. Weeds stressed by temperature and moisture extremes may not be effectively controlled.

To maximize weed control following a postemergence **Scepter 70 DG** application, wait at least 10 days before cultivating. This timely cultivation will enhance residual weed control, especially under dry conditions.

There must be an interval of at least 90 days between the **Scepter 70 DG** application and soybean harvest.

Tank mixes: **Scepter 70 DG** herbicide may be applied postemergence in a tank-mixture with the following products to improve weed control spectrum in soybeans and to better manage herbicide-resistant weed populations. Products listed for tank-mixture with **Scepter 70 DG** herbicide must be registered for postemergence application in soybean. It is the pesticide user’s responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture:

Assure II (quizalofop-P-ethyl), Fusilade DX (fluazifop-P-butyl), Fusion (fluazifop-P-butyl + fenoxaprop-P-ethyl), Hoelon (diclofop), Poast/Poast Plus (sethoxydim), Select/Select Max/Section/Arrow (clethodim), Blazer/Ultra Blazer (acifluorfen), Cobra/Phoenix (lactofen), Reflex/Flexstar/Rumble/Dawn/Rhythm (fomesafen).

Glyphosate or glufosinate may be mixed with Scepter 70 DG when applied to genetically modified soybean cultivars where glyphosate or glufosinate is labeled for use.

SOIL TEXTURE

Although **Scepter 70 DG application rate** does not vary with soil texture, the directed rates of other herbicides used with this product generally do vary with soil texture. Combination rate tables in this label refer to three soil texture groups: COARSE, MEDIUM, and FINE. Follow the rate instructions for soil texture included in the companion herbicide label. The following table lists soil textures included in each of these three groups:

COARSE	MEDIUM	FINE
sands	sandy clay loams*	silty clay loams*
loamy sands	sandy clays	silty clays
sandy loams	loams	clay loams
	silt loams	clays
	silts	

* Sometimes considered transitional soils

SPRAY ADDITIVES

Use a nonionic surfactant at 1-2 qt./100 gallons of water (0.25-0.50% v/v) or oil-based adjuvant such as methylated seed oil (MSO) at 0.5 to 1.5 qt./100 gallons of water (0.5-1% v/v) or crop oil concentrate (COC) at 1 to 2 qt./100 gallons of water (1-2% v/v) containing at least 14% emulsifiers and 80% oil active ingredient. Refer to the specific surfactant label for more specific use instructions. Apply the adjuvant at rate stated on manufacturer use label. When **Scepter 70 DG** is applied postemergence in a mixture with other herbicides, refer to the companion herbicide label for adjuvant recommendations.

A nitrogen fertilizer source can be added including urea ammonium nitrate (UAN; 28-34%) or ammonium phosphate (10-34-0) at 1.25 to 2.5 gallons per hundred gallons of water (1.25% to 2.5% v/v). Instead of a liquid fertilizer, spray grade ammonium sulfate (AMS) at 8.5 to 17 pounds per 100 gallons of water (or an equivalent liquid AMS product) may be used. Use the higher rate when making application during periods of hot dry weather and to larger weeds.

MIXING INSTRUCTIONS

Fill the spray tank one-fourth to one-half full with clean water. While agitating add the required amount of **Scepter 70 DG**, and then fill the remainder of the tank with water.

For postemergence applications, add the NIS, MSO, or COC as last ingredient in the tank. An antifoaming agent may be added to the tank if needed. Maintain agitation while spraying to ensure a uniform spray mixture.

When tank mixing **Scepter 70 DG** with directed herbicides, add **Scepter 70 DG** to the spray tank first and make sure it is thoroughly mixed before adding the other herbicide. Mixing instructions on the tank mix component herbicides need to be followed.

SOIL APPLICATIONS IN LIQUID FERTILIZERS

Scepter 70 DG can be applied to the soil in liquid fertilizers.

If **Scepter 70 DG** is applied in liquid fertilizer as a carrier, a pre-slurry must first be made. **Scepter 70 DG** must be completely dispersed in at least one gallon of water. Always test the compatibility of the **Scepter 70 DG** slurry with the liquid fertilizer before mixing in the spray tank.

Follow all **Scepter 70 DG** label directions regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment.

All individual state regulations relating to fluid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company selling the **Scepter 70 DG**/liquid fertilizer mixture.

SPRAYING INSTRUCTIONS

NOTE: DO NOT apply if wind conditions, temperature inversion conditions, or other conditions may cause drift onto adjacent areas or sensitive crops. Sensitive crops include but are not limited to leafy vegetables, sugarbeets, and cotton.

Avoid overlaps when spraying.

GROUND APPLICATIONS:

Uniformly apply with properly calibrated ground equipment in 10 or more gallons of water per acre. Use sprayers equipped with nozzles that provide accurate and uniform application. To minimize drift, use a maximum spray pressure of 40 psi.

To ensure thorough coverage, use a minimum of 20 gallons of water per acre when applying **Scepter® 70 DG herbicide** as a postemergence treatment. Use higher gallonage for fields with dense vegetation or heavy crop residues.

For band treatment, apply the broadcast equivalent rate and volume per acre. To determine these:

$$\begin{array}{ccccccc}
 & & & & \text{Broadcast} & & \text{Band} \\
 & & & & \text{RATE} & = & \text{RATE} \\
 \hline
 \text{Band width in inches} & \times & & & & & \\
 \text{Row width in inches} & & & & \text{per acre} & & \text{per acre}
 \end{array}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast VOLUME per acre} = \text{Band VOLUME per acre}$$

AERIAL APPLICATIONS:

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre.

To avoid injury to sensitive crops from drift, aerial applicators must adhere to the following SPECIAL AERIAL USE DIRECTIONS AND PRECAUTIONS:

- Use nozzles which produce a coarse spray.
- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed toward the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees.
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than three-fourths the distance from the center of the aircraft.
- Use a maximum spray pressure of 40 psi.
- A buffer zone must be established between the area to be sprayed and sensitive crops.

To the extent consistent with applicable law applicator is responsible for any loss or damage which results from spraying **Scepter 70 DG** in a manner other than directed in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted after applying **Scepter® 70 DG herbicide** at specified rates in soybeans:

CROP	USE REGION 1 and 2	USE REGION 3
Soybeans	Immediate	Immediate
Wheat	3 months ¹	18 months ²
Rice	Spring following Scepter 70 DG application	---
Barley	11 months	18 months
Field Corn (non IMI-CORN seed hybrids)	9.5 months	18 months ³
Edible Beans	11 months	11 months
Grain Sorghum	11 months	11 months
Oats	11 months	18 months
Peanuts	11 months	---
Tobacco	9.5 months	9.5 months
Sugar Beets & Red Table Beets	40 months	40 months
Other Crops	18 months	See FOOTNOTE 4

¹ In USE REGIONS 1 and 2 east of Interstate Highway 35, wheat may be planted 3 months following a soil application of **Scepter 70 DG** at a rate of 2.8 ounces per acre or a postemergence application up to 1.4 ounces per acre.

² In Nebraska, east of U.S. 283, south of U.S. 30, and west of U.S. 81, wheat may be planted 4 months after a **Scepter® 70 DG herbicide** application.

³ In USE REGION 3 corn may be planted in the spring of the following year following a postemergence application of Scepter 70 DG if the total amount applied in one season does not exceed 1.4 ounces per acre.

⁴ For USE REGION 3 as defined in the USE AREA section of this label; canola, strawberries, cabbage, tomatoes,

potatoes, carrots, celery, cole crops, garlic, onions, spinach, asparagus, cauliflower, and broccoli may be planted 26 months after a **Scepter 70 DG** application. Other crops may be planted 18 months after a **Scepter 70 DG** application.

USES WITH OTHER PRODUCTS (TANK-MIXES)

If this product is used in combination with any other product except as specifically directed in writing by AMVAC then to the extent consistent with applicable law AMVAC shall have no liability for any loss, damage, or injury arising out of its use in any such combination not so specified. If used in a combination directed by AMVAC the liability of AMVAC shall in no manner to the extent consistent with applicable law extend to any damage, loss or injury not directly caused by the inclusion of the AMVAC product in such combination use, and in any event shall be limited to return of the amount of the purchase price of the product.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Keep from freezing. **DO NOT** store below 32° F. **DO NOT** contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved disposal facility.

CONTAINER DISPOSAL

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity ≤ 50 pounds) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use, subject to the inherent risks referred to herein, when it is used in accordance with such directions; and (c) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions.

THERE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM

THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

AMVAC offers this product, and Buyer accepts it, subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC.

© 2016 AMVAC Chemical Corporation. A wholly owned subsidiary of American Vanguard Corporation. All Rights Reserved. American Vanguard, AMVAC, and the AMVAC logo are trademarks owned by AMVAC Chemical Corporation. Scepter is a trademark owned by BASF and used under license.

CHEMTREC is a service mark owned by the American Chemistry Council, Inc.

Scepter O.T., Basagran, Detail, Frontier, Galaxy, Prowl, Pursuit, Squadron, Steel, Storm and Tri-Scepter are trademarks of BASF.

Broadstrike, Sonalan and Strongarm are trademarks of Dow AgroSciences.

Canopy, Classic, Lexone, Staple, and Synchrony are trademarks of E.I. du Pont de Nemours and Company. Roundup, Roundup Ultra, and Lasso are trademarks of Monsanto Technology.

Cobra is a trademark of Valent.

Touchdown and Gramoxone are trademarks of Syngenta.

Command is a trademark of FMC Corporation.

Sencor is a trademark of Bayer AG.

CONCERT is a trademark of Syngenta Participations AG

GEMINI is a trademark of Everris International B.V.

PINNACLE is a trademark of Daniels Agrosiences, LLC.

PREMIER is a trademark of Premier Magnesia, LLC.

AMVAC Chemical Corporation
4100 E. Washington Blvd.
Los Angeles, CA 90023
1-888-462-6822