

50 LBS. NET WEIGHT

Fly Control Mineral No. 16-678 (Low Fluorine) (Contains RABON Oral Larvicide)

To prevent the development of Horn Flies, Face Flies, House Flies, and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENTS	By Weight
2-chloro-1-(2,4,6-trichlorophenyl) ethyl dimethyl phosphate	1.79%*
INERT INGREDIENTS	98.21%**

*Each pound contains 8.1 g. RABON.
**Refers only to ingredients which are not larvicidal.

GUARANTEED ANALYSIS

Calcium (Ca), Maximum	20.0%
Calcium (Ca), Minimum	17.0%
Phosphorus (P), Minimum	16.0%
Iodine (I), Minimum	0.001%
Salt (NaCl), Maximum	6.0%
Salt (NaCl), Minimum	4.0%
Magnesium (Mg), Minimum	2.0%

FEED INGREDIENTS

Plant Protein Products, Processed Grain By-Products, Mineral Oil, Anise Oil, D-Activated Animal Sterol (source of Vitamin D3), Salt, Calcium Carbonate, Dicalcium Phosphate, Iron Oxide, Manganous Oxide, Copper Oxide, Cobalt Carbonate, Calcium Iodate, Zinc Oxide, Magnesium Oxide.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See reverse side of tag for use directions and other precautions.



HONEGGER & CO., INC.
GENERAL OFFICES
FAIRBURY, ILLINOIS 61724

FAA REG. NO. 14804
FAA EX. NO. 14804
U.S. PAT. NO. 2,122,144 MAR. 1938
RABON® REG. U.S. PAT. & TM. OFF. © 1954

BEST DOCUMENT AVAILABLE

678
7D

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent to its labeling.

Feed Fly Control Mineral No. 16 678 at the rate of one ounce per 750 pounds of body weight per head per day to beef and dairy animals by mixing it into the ground grain or complete daily ration.

Start feeding Fly Control Mineral No. 16 678 early in the spring before flies begin to appear and continue feeding throughout the summer into the fall until cold weather restricts fly activity.

Fly Control Mineral No. 16 678 may be fed up to slaughter and to lactating dairy cows.

Fly Control Mineral No. 16 678 prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program during the fly season, it is desirable to use other control measures to reduce the population of existing adult flies.

In some cases, supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Fly Control Mineral No. 16 678 should be used in conjunction with other good management and sanitation practices.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE Store in a dry place.

CONTAINER DISPOSAL Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

BEST BUY