

poisonous bite when it bites. A person bitten by it  
 should wash the bite with soap and water. The legs  
 of the tick are long and slender. The body is oval  
 and back. The legs are long and slender. The body is oval  
 shaped like an hourglass. In the undersides of the  
 this marking is of a different shape. Black  
 Widow spiders seldom appear in living quarters of homes,  
 but often are found in basements, window wells, beneath  
 lawn benches or porches, and in garages, tool sheds, and  
 lumber piles, trash piles, sand boxes, and other outdoor  
 buildings. Spray suspected areas with 0.30% to 0.50% solu-  
 tion of F-T-C-SPRAY frequently. This same treatment  
 may be used to control all spiders around the home.

**SCORPIONS** Occasionally they enter the home. The best  
 way to keep scorpions out of the house is to use a surface  
 spray that will control scorpions. Apply F-T-C-SPRAY at  
 the 0.50% rate to the lower part of tree trunks, stumps,  
 piles of lumber, and rock walls. Also under structure that  
 is set on piers.

| DILUTION TABLE - Use 1 Gallon |          |          |          |
|-------------------------------|----------|----------|----------|
| 1.00%                         | 1.00 Oz. | 1.00 Oz. | 1.00 Oz. |
| 0.50%                         | 0.50 Oz. | 0.50 Oz. | 0.50 Oz. |
| 0.30%                         | 0.30 Oz. | 0.30 Oz. | 0.30 Oz. |
| 0.25%                         | 0.25 Oz. | 0.25 Oz. | 0.25 Oz. |

**CAUTION:** May be absorbed through skin. Avoid  
 inhalation and skin contact. In case of contact  
 wash immediately with soap and water. Avoid con-  
 tamination of feed and foodstuff. Do not use on  
 household pets or humans. Toxic to fish and wild  
 life. Do not contaminate any body of water (or ap-  
 ply to any area not specified on the label). **HARM-  
 FUL IF SWALLOWED. KEEP OUT OF THE  
 REACH OF CHILDREN.** Birds feeding on treated  
 areas may be killed. Do not contaminate water by  
 cleaning of equipment or disposal of wastes. **DO  
 NOT USE, POUR, SPILL OR STORE NEAR HEAT  
 OR OPEN FLAME.** Dispose of empty container  
 by burying.

JAMIESON CHEMICAL COMPANY  
 4011 Florida Ave., Tampa, Fla. 33603, Ph. 232-0111

"Products that make Life worth Living"



# F-T-C SPRAY

**AN EMULSIFIABLE CONCENTRATE  
 DILUTE WITH WATER  
 FOR OUTDOOR USE**

**CONTROL FLEAS - TICKS - CHIGGERS**  
 Also controls Spiders, Scorpions, Centipedes and  
 Millipeds

**ACTIVE INGREDIENTS** ..... 95.6%  
 Gamma Isomer of Benzene Hexachloride  
 From Lindane ..... 12%  
 Xylene ..... 83.6%  
**INERT INGREDIENTS** ..... 4.4%

**CAUTION: KEEP OUT OF THE REACH OF  
 CHILDREN. See side panel for additional precautions.**  
 U.S. DA REG. NO. 5490-7

5490-7

Jul 30 1971  
 UNDER THE FEDERAL INSECTICIDE  
 ACT AND REGULATIONS THEREUNDER  
 THIS PRODUCT IS REGISTERED FOR  
 SALE AND DISTRIBUTION IN THE  
 UNITED STATES OF AMERICA  
 BY JAMIESON CHEMICAL COMPANY

**DIRECTIONS FOR USE**

**FLEAS** To destroy fleas in yards and lawns, under  
 houses, spray with F-T-C-SPRAY using the 0.25% solu-  
 tion. Apply the spray at the rate of 2 quarts per 1,000  
 sq. ft. Apply with tank sprayer, garden hose sprayer or  
 other similar means. To control fleas inside the home use  
 our A-M-P-EX SPRAY.

**FLEAS ON DOGS** Wet the animal thoroughly with an ap-  
 plication of 0.25% solution of F-T-C-SPRAY. **CAUTION:**  
 Do not apply to puppies under two months old or to any  
 cats. Do not use more than once a week.

**TICKS** Control ticks on trees obtained by the use  
 of 0.50% solution of F-T-C Spray applied as a spot  
 treatment where ticks are found. Adequate coverage  
 should be made to obtain complete control. Treat-  
 ment will control infestation for about thirty days or  
 more. Warning to not spray in streams, ponds or  
 other water courses or their margins. Do not use  
 on food plants.

**CHIGGER CONTROL (Mites)** Control of chiggers can be  
 helped by clearing and removing underbrush and keeping  
 weeds and grass cut which will reduce infestation. Treat  
 the areas the same as for ticks outlined above. Do not  
 use on food plants.

**CENTIPEDES AND MILLIPEDES:** To prevent their entry  
 into the house, apply F-T-C-SPRAY at the 0.3% to 0.50%  
 rate to openings or cracks around doors, windows, founda-  
 tions, steps, and other entrances into the house.

**SPIDERS:** Most species of spiders that occur in the United  
 States are harmless. They destroy many injurious house-  
 hold insects, and are, therefore, beneficial. The Black  
 Widow spider is dangerous and will bite instinctively when  
 touched or threatened. The female inflicts a seriously

1-2-71 (Rev.)

poisonous bite which can be fatal. A person bitten by this spider should consult a doctor immediately. The female is about 1/2 inch long and shiny black. The front and back parts of the body are globular, the legs are long and slender. It usually has a red, orange or brownish yellow mark shaped like an hourglass on the underside, but sometimes this marking is of a different shape or is absent. Black Widow spiders seldom appear in living quarters of homes, but often are found in basements, window wells, beneath lawn benches or porches, and in garages, tool sheds, old lumber piles, trash piles, sand boxes, and other outdoor buildings. Spray suspected areas with 0.30% to 0.50% solution of F-T-C-SPRAY frequently. This same treatment may be used to control all spiders around the home.

**SCORPIONS:** Occasionally they enter the home. The best way to keep scorpions out of the house is to use a surface spray that will control scorpions. Apply F-T-C-SPRAY at the 0.50% rate to the lower part of tree trunks, stumps, piles of lumber, and rock walls. Also under structure that is set on piers.

**DILUTION TABLE - Use Water**

Use the Following Amounts of F-T-C (Emulsifiable Concentrate):

| Strength of Spray to Make | 0.025%     | 0.25%     | 0.50%  |
|---------------------------|------------|-----------|--------|
| 1 gallon                  | 1 1/2 Tsp. | 2 1/2 Oz. | 5 Oz.  |
| 5 gallons                 | 2 Tbs.     | 13 Oz.    | 26 Oz. |
| 15 gallons                | 3 Oz.      | 39 Oz.    | 78 Oz. |

**CAUTION:** May be absorbed through skin. Avoid inhalation and skin contact. In case of contact wash immediately with soap and water. Avoid contamination of feed and foodstuff. Do not use on household pets or humans. Toxic to fish and wild life. Do not contaminate any body of water nor apply to any area not specified on the label. **HARMFUL IF SWALLOWED. KEEP OUT OF THE REACH OF CHILDREN.** Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** Dispose of empty container by burying.

**JAMIESON CHEMICAL COMPANY**

4011 Florida Ave., Tampa, Fla. 33603, Ph. 232-0111



# F-T-C SPRAY

**AN EMULSIFIABLE CONCENTRATE  
DILUTE WITH WATER  
FOR OUTDOOR USE**

**CONTROL FLEAS - TICKS - CHIGGERS**  
Also controls Spiders, Scorpions, Centipedes and Millipeds

|                                      |       |
|--------------------------------------|-------|
| <b>ACTIVE INGREDIENTS</b> .....      | 95.6% |
| Gamma Isomer of Benzene Hexachloride |       |
| From Lindane .....                   | 12%   |
| Xylene .....                         | 83.6% |
| <b>INERT INGREDIENTS</b> .....       | 4.4%  |

**CAUTION:** KEEP OUT OF THE REACH OF CHILDREN. See side panel for additional precautions.

FLA. DA REG. NO. 5490-1

**DIRECTIONS FOR USE**

**FLEAS:** To destroy fleas in yards, and lawns, under houses, spray with F-T-C-SPRAY using the 0.25% solution. Apply the spray at the rate of 2 gallons per 1,000 sq. ft. Apply with tank sprayer, garden hose sprayer or other similar means. To control fleas inside the home use our A-M-P-EX SPRAY.

**FLEAS ON DOGS:** Wet the animal thoroughly with an application of 0.025% solution of F-T-C-SPRAY. **CAUTION:** Do not apply to puppies under two months old or to any cats. Do not use more than once a week.

**TICKS:** Control of ticks can be obtained by the use of 0.50% solution of F-T-C Spray applied as a spot treatment where ticks are found. Adequate coverage should be made to obtain complete control. Treatment will control infestation for about thirty days or more. Warning do not spray in streams, ponds or other water courses or their margins. Do not use on food plants.

**CHIGGER CONTROL (Mites):** Control of chiggers can be helped by clearing and removing underbrush and keeping weeds and grass cut which will reduce infestation. Treat the areas the same as for ticks outlined above. Do not use on food plants.

**CENTIPEDES AND MILLIPEDS:** To prevent their entry into the house, apply F-T-C-SPRAY at the 0.3% to 0.50% rate to openings or cracks around doors, windows, foundations, steps, and other entrances into the house.

**SPIDERS:** Most species of spiders that occur in the United States are harmless. They destroy many injurious household insects, and are, therefore, beneficial. The Black Widow spider is dangerous and will bite instinctively when touched or threatened. The female inflicts a seriously

1m3/71 (Rev.)